

Banda Arc

	Neo- proterozoic	Meso- proteroz



1	1	PROVENANCE OF CRETACEOUS SANDSTONES IN THE BANDA
2 3	2	ARC AND THEIR TECTONIC SIGNIFICANCE
т 5 б	3	
7 8 9	4	Sebastian Zimmermann ^{1,2*} , Robert Hall ²
10 11 12	5	
13 14 15	6	¹ iCRAG @ School of Earth Sciences, University College Dublin, Dublin 4, Ireland.
16 17	7	
18 19 20	8	² SE Asia Research Group, Department of Earth Sciences, Royal Holloway University of
21 22	9	London, Egham, Surrey, TW20 0EX, UK.
23 24 25	10	
26 27 28	11	* Corresponding author (Sebastian.zimmermann@ucd.ie)
29 30	12	
31 32	13	
33 34 25	14	# Provenance, Palaeogeography, Banda Arc, Tectonic reconstruction, Heavy minerals,
36 77	15	Zircon geochronology #
38	16	
40	17	
41 42	18	
43 44		
45		
46		
±/ 48		
49		
50 51		
52		
53		
54 55		
56		
57 58		
59		
50		
51 52		
53		
54		
55		

19 ABSTRACT

The provenance of Cretaceous sandstones in the Banda Arc islands differs from west to east. Sandstones in Sumba and West Timor contain significant amounts of feldspar (K-feldspar and plagioclase) and lithic fragments, suggesting a recycled to magmatic arc origin. In comparison, East Timor and Tanimbar sandstones are quartz rich, and suggest a recycled origin and/or continental affinity. Heavy mineral assemblages in Sumba and West Timor indicate metamorphic and minor acidic igneous sources and include a mixture of rounded and angular zircon and tourmaline grains. In East Timor, Babar and Tanimbar, an ultimate origin from a mainly acid igneous and minor metamorphic source is interpreted, containing a mixture of rounded and angular zircon and tourmaline grains.

Detrital zircon ages in all sandstones range from Archean to Mesozoic, but variations in age populations indicate local differences in source areas. Sumba and West Timor are characterised by zircon age peaks at 80-100 Ma, 200-240 Ma, 550 Ma, 1.2 Ga, 1.5 Ga and 1.8 Ma. East Timor and Tanimbar contain 80-100 Ma, 160-200 Ma, 240-280 Ma, 550 Ma and 1.5 Ga zircon peaks. Most populations are also common in Triassic and Jurassic formations along the Outer Banda Arc and in many other areas of SE Asia. However, the abundance of Jurassic and Cretaceous populations was unexpected. We interpret Cretaceous sandstones from Sumba, Timor and Tanimbar to have been deposited in SE Sundaland. Syn-sedimentary Cretaceous (68-140 Ma) sources are suggested to include the Schwaner Mountains in SW Borneo and Sumba. Material derived mainly from older recycled sediments that had their main sources in the Bird's Head, Western and Central Australia, and local sources close to Timor.

1. Introduction

SE Asia is composed of fragments of continental crust, oceanic crust and volcanic arcs, as a result of subduction-related processes in the region, that now form a complex tectonic assemblage (e.g. Hamilton, 1979; Metcalfe, 1998; Hutchison, 1989; Bowin et al., 1980; Hall, 1996, 2011, 2012; Packham, 1996; Charlton, 2001; Hinschberger et al., 2005). Since the Palaeozoic, rifting led to separation of fragments from the Gondwana margin, that were later successively accreted to Sundaland at different times (Fig. 1) (Audley-Charles et al., 1988; Hall, 2012; Barber and Crow, 2009; Metcalfe, 2011, 2013).

During the Mesozoic, large rivers drained the Australian continent and filled the major offshore basins of the NW Shelf, within a fluvial to marginal marine setting (Bishop, 1999; Barber et al., 2003), creating important offshore hydrocarbon reservoirs. The southern Outer Banda Arc islands of Sumba, Timor, Babar and Tanimbar (Fig. 2) are assumed to include the onshore equivalents of these sediments. Mesozoic sandstones have been exposed on these islands due to subduction and collision processes in a complicated tectonic history. Based on heavy minerals and detrital zircon geochronology, Triassic and Jurassic sandstones were interpreted to contain detritus derived from Western and Central Australia, but also an important component previously not recognised, from the Bird's Head region (Fig. 3), and some material from local Jurassic volcanism within the Inner Banda Block (Zimmermann and Hall, 2016).

Late Jurassic rifting of continental slivers from the Australian margin has been described by various researchers (e.g. von Rad et al., 1982; Longley et al., 2002; Heine and Müller, 2005; Hall et al., 2009; Hall, 2012; Heine et al., 2012; Gibbons et al., 2012). Tectonic blocks rifted from areas between the Exmouth

Plateau in the west and the Arafura Sea in the east, where separation of the continental blocks formed the Banda embayment north of Australia (Hall, 2002, 2012; Charlton, 2012; Spakman and Hall, 2010), and left the Sula Spur (Klompé, 1954) northeast of Australia. The clastic sediments deposited in the Outer Banda Arc islands before the Late Jurassic rifting were discussed by Zimmermann and Hall (2016). During the Cretaceous, the Australian-derived continental fragments moved northwards and were accreted to the Sundaland (Fig. 1) margin (e.g. Audley-Charles, 1968; Hamilton, 1979; Wensink, 1994; Wakita et al., 1996; Parkinson et al., 1998; Abdullah et al., 2000; Harris, 2006; Hall et al., 2009; Metcalfe, 2011; Hall, 2012) and clastic sediments were deposited on this continental crust during their northward movement and after their arrival in SE Asia. During Australia-SE Asia collision, since the Early Miocene, slices of the SE Asian margin were thrust back onto the Australian continental margin and are now found in the Outer Banda Arc islands.

The Cretaceous sandstones discussed in this paper were deposited after separation of blocks from the Gondwana margin in the Late Jurassic and before their Neogene thrusting onto the Australian margin in the Outer Banda Arc islands. Details are still not completely understood and are partly the subject of on-going research. We present petrology, heavy mineral analyses and U-Pb ages of detrital zircons from sandstones, siltstones and meta-sandstones. These data help to identify the origin and possible sources of detrital material and to consider previously debated tectonic fragmentation in the Banda Arc.

2. Post-Triassic Sundaland margin history

The continental blocks rifted in the Late Jurassic were added to Sundaland in the Cretaceous to form what is now much of Borneo, East Java and West Sulawesi. Three principal blocks have been identified (Hall et al., 2009; Metcalfe, 2009; Hall, 2012): the Banda, Argo and Inner Banda blocks, which rifted to leave the oceanic Banda embayment in the northwest Australian margin.

The Banda block was the first to be accreted to the Sundaland margin and is now identified with SW Borneo. For many years SW Borneo was interpreted as a fragment of Asian/Cathaysian origin (e.g. Hutchison, 1989; Metcalfe, 1988, 1990, 1996). The area of the Schwaner Mountains and further south was described **100** as the 'Basement Complex', 'Continental Core' or 'Sunda Shield' (van Bemmelen, ²⁷ **101** 1949) or the West Borneo Basement (Haile, 1974). It was considered to be an ⁻₃₀ 102 ancient continental area, but metamorphic rocks suggested to be Triassic or older **103** (e.g. van Bemmelen, 1949; Haile, 1974; Tate, 1991; Tate and Hon, 1991) were undated and known only to be intruded by Cretaceous granitoids. SW Borneo includes Cretaceous and Jurassic granites in the Schwaner Mountains (Haile et al., **105** ³⁹ 106 1977; Davies et al., 2014) and the metamorphic rocks are now known to be **107** Cretaceous (Davies et al., 2014). Recent work in Sarawak has identified the ⁴⁴ 108 boundary of Triassic Sundaland and suggests the Banda/SW Borneo block arrived in 47¹⁰109 the Early Cretaceous at c. 130 Ma (Hennig et al., 2017).

110 SW Borneo is bounded on its southeast side by the Meratus suture, which can be traced from SW to NE from West Java to the Meratus Mountains of SE **111** ⁵⁴ -- **112** Borneo, considered by Hamilton (1979) to be the approximate southeastern boundary of Cretaceous continental crust. The Meratus suture separates SW Borneo **113** ⁵⁹ 114 from other continental fragments added to the Sundaland margin later in the

Cretaceous. In the suture zone in Java and SE Borneo are ophiolitic, arc rocks and high pressure-low temperature metamorphic rocks which record subduction beneath Sundaland in the Early Cretaceous. Accretionary-collision complexes in SE Borneo ⁷ 118 and Java (Katili, 1971, 1973; Sukamto, 1975a,b; Sikumbang, 1986, 1990; Schiller et ⁹ 119 al., 1991; Wakita et al., 1994a,b, 1998; Parkinson et al., 1998; Wakita, 2000; ¹² **120** Clements et al., 2009) include rocks formed by oceanic spreading, arc volcanism, oceanic and forearc sedimentation, and subduction-related metamorphism. K-Ar **122** ages from metamorphic rocks summarised by Parkinson et al. (1998) indicate high pressure-low temperature metamorphism between 117 and 124 Ma, and radiolaria **124** associated with pillow lavas in Java are Early Cretaceous (Wakita et al., 1994b). ²⁴ 125 Based on evidence from SE Borneo, Sikumbang (1986, 1990) and Wakita et al. **126** (1998) concluded that ophiolite emplacement and arc-continent collision was ²⁹ 127 completed by about 90 Ma.

Several authors (e.g. Luyendyk, 1974; Ricou, 1994; Wakita et al., 1996; Parkinson et al., 1998; Sribudiyani et al., 2003; van Leeuwen et al., 2007; Smyth et al., 2007) have suggested that continental fragments accreted to Sundaland in the Cretaceous, outboard of the Meratus suture, had a Gondwana origin. The identification of different blocks in Java, SE Borneo and Sulawesi and interpretation of their former position on the Australian margin are based largely on zircon age data from different areas.

The Argo block forms the area that now includes much of East Java and West Sulawesi. Studies in East Java show that the southern part of the island is underlain by continental crust (Smyth et al., 2007) and suggest that there is similar crust beneath the Java Sea and in the forearc south of East Java (Deighton et al., 2011; Granath et al., 2011; Nugraha and Hall, 2012). Inherited zircons in Cenozoic

sedimentary and igneous rocks of East Java range in age from Archean to Cenozoic. The distribution of zircons reveals two different sources. Clastic rocks in north and west parts of East Java contain Cretaceous zircons, which probably came from the ⁷ 143 Meratus suture or from SW Borneo. In contrast, the Early Cenozoic Southern Mountains volcanic arc of East Java includes abundant acid volcanic and intrusive ¹² **145** rocks which contain only Archaean to Cambrian zircons. These indicate deep continental Gondwana crust below East Java which originated close to western Australia (Smyth et al., 2007, 2008). Australian-origin continental crust is also **147** considered to underlie parts of the southern Makassar Straits and East Java Sea **149** between Borneo and Java based on basement rocks encountered in exploration ²⁴ 150 wells (e.g. Manur and Barraclough, 1994; Satyana, 2015). Deep seismic data **151** suggest there is similar crust beneath the Java Sea south of Pulau Laut in SE ²⁹ 152 Borneo (Emmet et al., 2009; Granath et al., 2011) and south of East Java (Deighton et al., 2011; Nugraha and Hall, 2012).

Initial reconstructions (Hall et al., 2009; Hall, 2012) of the rifted blocks **155** suggested that the Argo block included all of present-day East Java-West Sulawesi. Hennig et al. (2016) have since shown that zircon age data from NW Sulawesi **157** indicate that the underlying continental crust originated in a position east of the Argo ⁴⁴ 158 block, and formed part of an Inner Banda block. There are other indications that a **159** single Argo block may be an over-simplification of the deep continental crust east of ⁴⁹ 160 the Meratus suture, but reconstruction is difficult because basement rocks are limited 52 **161** to small areas and mainly overlain by Cenozoic rocks. Jurassic ammonites and bivalves reported from South Sulawesi (Sukamto et al., 1990; Sukamto and **162** Westermann, 1993) suggest an Australian continental fragment at depth. Elsewhere in western Sulawesi there is evidence from inherited zircons, and from chemical **164**

б

10

characteristics of Cenozoic igneous rocks, of underlying continental basement (Priadi et al., 1993, 1994; Bergman et al., 1996; Polvé et al., 1997, 2001; Elburg and Foden, 1999; Elburg et al., 2003). There are blueschists and other high pressure-low temperature metamorphic rocks known from inliers in the Bantimala and Barru areas in South Sulawesi (Sukamto and Supriatna, 1982; Miyazaki et al., 1996, 1998; **170** Parkinson et al., 1998; Maulana et al., 2010, 2013) suggesting sutures between continental blocks. Neogene potassic volcanics in SW Sulawesi do not show the **172** Australian continental isotopic signatures shown by similar volcanic rocks further north in Sulawesi (Elburg et al., 2003) which may indicate an underlying suture. All ₂₂ 174 the supposed suture rocks are very far east of the Meratus suture and cannot be ²⁴ 175 connected to it; they could indicate the Argo block is actually made up of multiple **176** continental fragments separated by sutures or alternatively the observations could ²⁹ **177** be reconciled by postulating the Argo block was hyper-extended continental crust within which were zones of exhumed mantle and deep marine sediments.

The character of the deep crust remains uncertain because there are few areas exposing rocks older than Cenozoic. Geochemistry and SO palaeomagnetic studies suggest that Sumba formed part of the Sundaland margin by the Late Cretaceous (Wensink, 1994; Abdullah et al., 2000) but the character of the deep crust is unknown. ³He/⁴He ratios suggest that Australian continental crust was involved in genesis of magmas throughout the inner Banda arc from the Banda Ridges to Flores (Hilton et al., 1992). Similar isotope geochemical studies could help identify if the deep crust beneath Sumba is ancient continental or of younger arc origin.

After the arrival of the Argo and Inner Banda blocks, subduction **188** ⁵⁹ 189 ceased around the Sundaland margin at c.90 Ma. Thus, the outer part of the SE

б

10

Sundaland margin was underlain largely by Australian-origin continental crust (Fig. 4). Subsequently, there was a short-lived episode of subduction between the latest Cretaceous and Eocene. Extension in the Eocene formed the Makassar Straits, although the amount of extension in the South Makassar Straits was small (e.g. Situmorang, 1982; Johansen, 2007; Kupecz et al., 2013; Armandita et al., 2015), and **195** there has been major extension of the eastern part of the region which includes Sulawesi and Sumba during Banda rollback from the Middle Miocene (Rigg and Hall, 2012; Camplin and Hall, 2014; Nugraha and Hall, 2018). **197**

10

3. Stratigraphic background

Fig. 2 shows a simplified stratigraphy of the Cretaceous siliciclastic sedimentary rocks investigated (sampling locations in supplementary data file 1).

Sumba 3.1

Upper Cretaceous (Coniacian to Campanian) turbidites of the Lasipu Formation (Fig. 5) in Sumba were described by Burrolet and Salle (1982) and von der Borch et al. (1983). Cretaceous ages were determined by fossil fragments, bivalves, molluscs and gastropods (*Exogyra sp., Mytilidae sp., Parainoceramus sp.,* Platyceramus sp. and Actaeonella sp.) (von der Borch, 1983). Abdullah et al. (2000) described three magmatic episodes during the Cretaceous to Paleogene, generating pyroclastic rocks, basaltic-andesitic lava flows and granodioritic intrusions.

⁵³ 210 Rocks collected in Sumba during this study were assigned to the Upper 5₆ 211 Cretaceous Lasipu Formation. Localities sampled are divided into the central, south-⁵⁸ 212 central, western and eastern regions. This newly proposed subdivision highlights

important differences in diagenetic processes and low-grade metamorphism of the ² 214 sediments: the key differences are massive highly indurated metamorphosed rocks, with the absence of mudstone in the north, thick siltstone beds interbedded with thin mudstone layers in the west, and cyclic siltstone-mudstone interbedded sequences, with intercalated sandstone beds in the south and east. A sedimentary log in Fig. 5 shows an example of the well-bedded siltstone-sandstone intercalations (Fig. 5A) at Konda beach in south central Sumba, locally with channel structures (Fig. 5B) and layers up to 10-centimetre-thick of medium to coarse-grained sandstone (Fig. 5C). 13 samples from various locations in Sumba were analysed petrologically, 9 for heavy minerals and 7 for zircon geochronology. SUM 06 contains recrystallised Late Cretaceous foraminifera (Globotruncana sp.) from a shallow inner neritic environment (M. BouDagher-Fadel, pers. comm., 2012).

3.2 Timor

Timor consists predominantly of Mesozoic sedimentary rocks and exotic fragments, that were described as tectonically distributed over the island (Audley-Charles, 1986). An important feature is the suggested subdivision of units into 1) autochthonous components derived from the Australian continent (lower nappes) and 2) allochthonous units of non-Australian origin (upper nappes). These have been discussed by many researchers and termed the Banda Allochthon (e.g. Audley-Charles & Harris, 1990; Harris, 1991; Audley-Charles, 2011) and commonly explained by an alpine-style "overthrust model" (e.g. Carter et al., 1976; Barber, 1979; Norvick, 1979; Brown and Earle, 1983; Audley-Charles, 1986; Harris, 1989; Audley-Charles and Harris, 1990).

Cretaceous sandstones in West Timor belong to the Oe Baat Formation (Sawyer et al., 1993) (Fig. 1) which is a massive calcareous sandstone of grey-green colour (Samples SZ 26, SZ 27 and SZ 47). In East Timor, Cretaceous sandstones have been assigned to the evenly bedded and finely laminated Seical Formation (ET 17). The Seical Formation was not previously recognised at the sampling location, and sandstones there had been interpreted to be Triassic as they resemble rocks of the Babulu Formation in West Timor (Audley-Charles (1968). However, the samples collected contain Late Albian-Early Cenomanian foraminifera (Hedbergella sp., Favusella washitensis, Favusella sp.) from a shallow inner neritic environment (M. Boudagher-Fadel, pers. comm., 2012) and Cretaceous zircons.

3 3.3 Babar

The island of Babar is a typical mud volcano. Before this research no Cretaceous rocks had been reported from Babar. However, in central Babar there is a steep cliff of hard and dense fine-grained siltstone/meta-sandstone with common thick bedding (sample BAB 25). No fossils have been found. The hard and dense meta-sedimentary character of the siltstone closely resembles Cretaceous lithologies seen in Sumba and, like the Lasipu Formation, lithologies were associated with igneous intrusions. As discussed later, detrital zircon ages proved the maximum depositional age of these rocks to be Late Cretaceous.

3.4 Tanimbar

Mesozoic sandstones in Tanimbar are exposed on the western islands (Fig. 6A) that are tectonically separated from the main island to the south-southeast

261 (Kaye, 1989; Charlton et al., 1991). Cretaceous sandstones were assigned to the ² 262 Ungar Formation and divided informally into two members (Charlton et al., 1991). We recognise the same sandstone members, but include a third intermediate red 263 264 shale member, which contains radiolarian cherts dated by Jasin and Haile (1996) as uppermost Jurassic to Lower Cretaceous (Zimmermann and Hall, 2016). This member was informally named the Arumit Member (Charlton, pers. comm., 1996, in Jasin and Haile, 1996). The Ungar Formation members are: (1) Lower Sandstone 267 Member, (2) Arumit Member, and (3) Upper Sandstone Member. Neither of the 269 sandstone members has previously been dated. Palynomorphs indicate the Lower Sandstone Member is Upper Jurassic (Zimmermann and Hall, 2016). The Upper Sandstone Member must be Early Cretaceous or younger, based on the work of Jasin and Haile (1996).

The Arumit Member forms an estimated 60m thick clear marker throughout 6B). It consists of well-bedded thin the islands (Fig. red siltstone-mudstone interbeds that dip steeply to the southwest (Fig. 6C and Fig. 6D). Four distinctive chert horizons were recognised within the member and are highlighted in the sedimentary log in Fig. 7. Cherts are between 7 and 15 cm thick and yield radiolaria that were dated by Jasin and Haile (1996). Two horizons were 279 identified as Upper Jurassic (upper Tithonian-Berriasian) and Lower Cretaceous (Valanginian-Barremian) age. The contact of the Arumit Member with the Upper Sandstone Member is indicated in Fig. 6C, which is characterised by mud lenses and imbricated mud clasts within the sandstones.

The Upper Sandstone Member of the Ungar Formation includes fine to medium-grained arkosic sandstones. Rocks are commonly massive bright yellowishbrown. INPEX (pers. comm., 2012) provided comprehensive information for some

65

sandstones from the islands, based on microfossil analyses (foraminifera,
nanoplankton, palynomorphs and radiolaria). The rocks collected for this research
were correlated with these previously dated (Upper Jurassic - Lower Cretaceous)
samples. 8 samples from various islets were analysed petrologically, 5 for heavy
minerals and 4 for zircon geochronology.

4. Methodology

4.1 Petrology

Point counting of at least 300 grains of quartz, feldspar and lithic rock fragments (>0.0625 mm) by standard methods (Galehouse, 1971) was used to acquire light mineral modes, shown on ternary plots for each unit (Dickinson and Suczek, 1979; Dickinson et al., 1983). The fields in the diagrams for QFL (Quartz-Feldspar-Lithics) and QmFLt (Quartz monocrystalline-Feldspar-Lithic total) indicate possible derivation from a 'continental block', 'recycled orogen' or 'magmatic arc'.

300 Textural categories from 1 to 4 were assigned to qualitatively show 301 sorting and roundness of grains. Sorting categories are (1) poorly sorted, (2) 302 moderately sorted, (3) well sorted and (4) very well sorted. Rounding categories are 303 (1) angular, (2) sub-angular, (3) sub-rounded and (4) rounded.

5 4.2 Heavy Minerals

Detrital heavy minerals were analysed using standard methods after Mange and Maurer (1992). Samples collected were crushed, decarbonated in 10% acetic acid, sieved and washed (meshes of 0.063mm and 0.250mm) and separated

in a funnel using sodium polytungstate (SPT: 3Na₂WO₄,9WO₃,H₂O) or the lithium equivalent lithium polytungstate (LST), which have densities between 2.82-2.95 g/ml at room temperature. Identification of heavy minerals was performed manually by ⁷ 312 using an optical polarising microscope (NIKON Eclipse Lv 100) and additional SEM ₁₀ 313 analyses were performed to confirm selected grains. The ribbon count method used **314** was described by Galehouse (1971).

Common heavy minerals were grouped by their most likely protoliths, **315** based on suggested source rock associations (Feo-Codecido, 1956; Mange, 2002; 20 317 Nichols, 2009). Zircon, tourmaline, anatase, monazite, topaz and xenotime are ²² 318 considered to indicate acid igneous (granitic) sources. Pyroxene (Ortho-OPX and **319** Clino-CPX), titanite (sphene) and chromium spinel represent basic igneous and ²⁷ **320** ultrabasic (commonly arc-related) sources. Rutile, garnet, epidote, andalusite, ⁻⁾₃₀ 321 sillimanite, kyanite, chlorite, staurolite and corundum are interpreted to indicate **322** metamorphic sources, mainly of continental character. Other minerals, such as amphibole, baryte, brookite, zoisite, clinozoisite, sphalerite, prehnite, chloritoid, **324** cassiterite, allanite and vesuvianite are present, either in very low percentages or can be assigned to more than one group. Apatite is a very common mineral and **326** abundant in all samples of this study (up to 50%). Since it can be found in different ⁴⁴ 327 groups (acid igneous, granite pegmatite, contact metamorphic and basic igneous), it **328** is treated separately.

Varietal studies of zircon (colourless: euhedral, subhedral, subrounded, ⁵² 330 rounded, anhedral, elongate, zoned; purple: rounded, euhedral; brown, matrixattached) and tourmaline (brown: rounded, euhedral; blue: rounded, euhedral; green: all shapes) were performed during counting. Three types of grain shapes were **332** recognised: 1) euhedral, subhedral, anhedral, elongated and zoned zircons were

б

grouped into an 'euhedral' group; 2) rounded and subrounded zircons were grouped into a 'rounded' group; 3) grains with matrix attached represent the third group. Tourmaline grains were classified as either rounded or euhedral.

4.3 Zircon geochronology

Geochronology, using detrital zircons, is a powerful method to assess provenance and correlate sedimentary units (e.g. Goldstein et al., 1997; Cawood et al., 1999, 2003; Fedo et al., 2003; Gehrels et al., 2006; Sevastjanova et al., 2010; Schoene, 2014). The maximum depositional age (MDA) of sedimentary rocks can be determined (Dickinson and Gehrels, 2009) and geochronology is a valuable tool to improve tectonic models and palaeogeographic reconstructions (Murphy et al., 2004).

Selected samples were imaged with scanning electron microscope cathodoluminescence (SEM-CL) at University College London. U-Pb ages were acquired at University College London using Laser Ablation Inductively Coupled Plasma Mass Spectrometry (LA-ICPMS). U and Pb isotopes were analysed, using the following parameters: spot sizes of the ablation pits: 20-35µm; pulse repetition: 8-10Hz; dwell time: 25s; warm-up: 10-15s; wash-out: 18s. The ablated material was carried in helium gas into the plasma. A quadrupole mass spectrometer (Agilent Technologies 7700 Series ICP-MS) was used. Standards that were used were the Plešovice zircon (337.13±0.37 Ma) by Sláma et al. (2008) and a reference glass NIST SRM 612 (Pearce et al., 1997).

987 selected zircons (Sumba 321, West Timor 133, East Timor 141, Tanimbar 392) were chosen to investigate the relationship of grain shapes and the

analysed ages. The aim was to distinguish optically between rounded grains with
recycled histories and euhedral grains, which could have formed close to the time of
deposition. A simplified classification scheme was applied, using CL images of the
mounted zircons. Morphologies were subdivided into four groups: 1) euhedral; 2)
subhedral; 3) subrounded; 4) rounded.

5. Results

5.1 Light Minerals: Textures and petrography

Fig. 8 shows the results of point counting and textural analysis of sandstones from the various islands (tables in supplementary data 2). In general, samples are dominated by quartz, with varying abundances of feldspar and lithic fragments. Sorting and rounding vary between the islands.

13 samples from the Cretaceous Lasipu Formation in **Sumba** were analysed. Grains are angular to subrounded (2-3) and moderately to very well-sorted (2-4). Compositions are dominated by quartz (24-67%), lithic fragments (17-59%) and feldspar (14-26%). However, samples show slight variations of the modal composition between the different areas (Fig. 8). Central and western Sumba are characterised by high volcanic quartz and K-feldspar contents, while south central and eastern Sumba sandstones are dominated by polycrystalline quartz. Samples studied, commonly show a 'recycled orogen' to 'magmatic arc' modal composition on the QFL diagram and a strong magmatic arc origin on the QmFLt diagram (Fig. 8). Textures contain high rounding and sorting values, which plot across the mature field.

In West Timor sandstones, the grains are sub-angular and moderately to well sorted. Compositions are dominated by guartz (50-66%), with varying content of feldspar (16-27%) and lithic fragments (18-23%). Quartz types are dominated by polycrystalline guartz (QFL-total: 50% Qp vs 29% Qv vs 21% Qm). As shown in Fig. 8, samples plot in the recycled orogen field on the QFL diagram and within the dissected magmatic arc on the QmFLt diagram. On the textures plot, samples scatter across the immature to mature field boundary (Fig.8).

East Timor sandstones contain well sorted (3) sub-angular to rounded (2-4) grains. Marked predominance of guartz (80-89%) is characteristic (Fig. 8). There are also feldspars (5-11%) and lithic fragments (6-9%). The QFL and QmFLt diagrams indicate a 'quartzose recycled orogen' modal composition.

Cretaceous samples from the Upper Sandstone Member of the Ungar Formation in Tanimbar are quartz-rich (77-97%) arenites (Fig. 8). Grains are commonly sub-rounded to rounded (3-4) and moderately to very well sorted (2-4). Lithic fragments (max. 9%) and feldspar (max. 14%) are insignificant. Modal compositions plot in the 'recycled orogen'/'continental block' (QFL) to a 'quartzose recycled' (QmFLt) fields. Rounding and sorting indicate a texturally mature polycyclic character (Fig. 8).

Heavy Minerals and their protoliths 5.2

Heavy minerals are mainly ultra-stable minerals zircon, tourmaline and rutile, accompanied by apatite, garnet, subordinate and alusite and minor chlorite (tables in supplementary files 3). Fig. 9 summarises heavy mineral assemblages, protoliths and zircon and tourmaline morphology types.

405 The Lasipu Formation in **Sumba** has a significant metamorphic signal 2 406 (39-82%) that reflects the abundance of mainly garnet and andalusite (Fig. 9). 4 ₅ 407 Sandstones contain on average 14% acid igneous and 2% basic igneous grains. ⁷ 408 There is a mixture of rounded (17-67%) and euhedral zircon grains (12-48%). An 10 **409** average of 36% of zircon grains are attached to a matrix. Tourmaline is generally 11 12 **410** less abundant, but dominated by euhedral grains. 13

15 **411** The **West Timor** sandstone contains a metamorphic (38%) and acidic ¹⁷ **412** igneous (29%) signal indicated by chlorite and zircon (Fig. 9). Zircon grains are a 20 413 mixture of rounded (48%), euhedral (25%) and attached to matrix (27%).

23 **414** The East Timor sandstone is dominated by grains from an acidic ²⁵ 415 igneous protolith (49%), based on zircon and tourmaline, and a metamorphic source ₂₈ 416 (33%) mainly based on rutile and andalusite (Fig. 9). Zircon grains are mainly ³⁰ 417 rounded (~65%) and tourmalines are dominated by euhedral grain shapes (~74%).

³³ 418 The Babar sandstone does not contain a significant amount of ultra-₃₆ 419 stable heavy minerals. The strong metamorphic signal (Fig. 9) is based on chlorite 38 420 which represents 81% of the sample.

Tanimbar sandstones consist on average of 52% acid igneous grains 41 **421** ⁴³ 422 (mainly zircon and tourmaline). 32% metamorphic sources are mainly based on 46 **423** rutile, and alusite and garnet. Morphologies of zircons are dominantly euhedral (42-⁴⁸ **424** 66%) and rounded (34-54%). Tourmalines on average are 57% euhedral and 43% ₅₁ 425 rounded.

53 ₅₄ 426

1

3

6

8 9

14

16

19

21 22

24

26 27

29

31 32

34 35

37

39 40

42

44 45

47

49 50

52

- 62 63 64
- 65

427 5.3 Zircon Geochronology

The numerical ages assigned here to periods, epochs and stages are based on Gradstein et al. (2012). Data tables of LA-ICP-MS analyses are provided in the supplementary data 4, which contains ²⁰⁷Pb/²³⁵U ratios, ²⁰⁶Pb/²³⁸U ratios, calculated ages and preferred ages, considering exclusion of discordant grains. Histograms of detrital zircon ages and according grain morphologies of Cretaceous sandstones in the Banda Arc show generally a mixture of different age populations (Fig. 10), with similar proportions of Precambrian to Phanerozoic grains.

In Sumba 583 concordant LA-ICP-MS U-Pb detrital zircon ages were ²³ 436 obtained from samples SUM 1, SUM 6, SUM 10, SUM 21, SUM 22, SUM 24 and 26 **437** SUM 30. The youngest zircon ages (71.9±1 Ma in SUM 6 to 84.7±1.6 Ma in SUM 10) ²⁸ 438 constrain the maximum depositional age (MDA) as Campanian to Maastrichtian ₃₁ 439 (mean age of three youngest overlapping zircon ages 69.3 Ma). Samples contain zircons with 48.5% Phanerozoic, 47.9% Proterozoic and 3.6% Archean ages. Most abundant age populations (Fig. 10A) are Neoproterozoic (20.6%), Cretaceous (19.9%) and Mesoproterozoic (14.9%). The main peaks are at 80-100, 230, 550, ⁴⁰ **443** 1200 and 1600 Ma. Zircon grain morphologies of Cretaceous grains are dominated by euhedral and subhedral grains, Permian-Triassic zircons are euhedral and subhedral, Cambrian to Carboniferous zircons are subhedral to subrounded and Precambrian grains are mainly subrounded and rounded (Fig. 10B).

447 Sample SZ 47 from **West Timor** was sampled at a location mapped as 448 the Oe Baat Formation which has a Tithonian to Berriasian age (Charlton, 1987; 449 Sawyer et al., 1993). It contains 132 concordant zircon ages. However, the sample 450 contains 16 Cretaceous grains and the youngest zircon age (75.5±1.4 Ma) indicates

65

451 a Campanian (Late Cretaceous) MDA, which means that it cannot be part of the Oe ² 452 Baat Formation. This shows the value of the detrital zircon ages which indicate that a ₅ 453 sandstone mapped previously as an older formation belongs to a different, so far ⁷ 454 unnamed, formation. SZ 47 contains 40.9% Phanerozoic, 55.3% Proterozoic, and 10 **455** 3.8% Archean ages. There are no Jurassic grains. The most abundant age ¹² **456** populations are Neoproterozoic (22.7%),Paleoproterozoic (16.7%)and 457 Mesoproterozoic (15.9%). The main peaks are at 85, 220-240, 550 and 1600 Ma 15 (Fig. 10A). Zircon grains with Cretaceous ages are dominated by euhedral and 17 **458** ¹⁹₂₀ **459** subhedral grains. Paleozoic zircons show a mix of subrounded/rounded and 22 **460** euhedral/subhedral grain morphologies. Proterozoic zircons are mainly rounded and ²⁴ 461 subrounded (Fig. 10B). It is not clear to which formation this sandstone should be ₂₇ **462** assigned, but, there are clearly Cretaceous or younger sandstones in southern West ²⁹ **463** Timor. Sample SZ 37 was collected in the Kolbano area and also mapped as the Oe ³¹₃₂ 464 Baat Formation and contains belemnites supporting this assignment. However, it contains a single Cretaceous zircon, with one Permian grain, and abundant 34 **465** 466 Proterozoic zircons, but also contains Eocene foraminifera (M. BouDagher-Fadel, 39 **467** pers. comm., 2015). The belemnites must therefore be reworked. The Eocene rocks 468 in Kolbano were all previously assumed to be carbonates only and clearly need remapping (A.J. Barber, pers. comm., 2018). 44 469

ET 17 in **East Timor** provided 144 concordant analyses. The youngest 47 **470** ⁴⁹ **471** zircon age (97.3±1.3 Ma) constrains the MDA to the Late Cretaceous (Cenomanian). 52 **472** Zircon populations include 58.9% Phanerozoic, 39.7% Proterozoic and 1.4% ⁵⁴ **473** Archean ages. The most abundant age populations are Cambrian to Carboniferous 57 **474** (23.6%), Neoproterozoic (22.2%) and Permo-Triassic (18.8%). There is a small population of Cretaceous zircons (6.9%) and a greater number of Jurassic grains 59 **475**

63 64 65

1

3 4

6

8 9

11

13 14

16

18

21

23

25 26

28

30 31

33

35 36

37 38

40 41

42 43

45 46

48

50 51

476 (10.4%) The main peaks are at 95, 170, 260, 280 and 550 Ma (Fig. 10A). There is no 477 1800 Ma peak. It is striking that most zircons have subhedral and euhedral grain morphologies for most populations. The predominance of euhedral and subhedral 478 ⁷ 479 grains for Cretaceous, Jurassic, Permo-Triassic and Cambrian-Carboniferous 10⁴⁸⁰ suggests little recycling of most zircons (Fig. 10B).

13 **481** No Cretaceous rocks have previously been described from Babar. As ¹⁵ 482 mentioned earlier, sandstones are unfossiliferous and undated. The youngest zircon 483 (104.7±1Ma) in sample BAB 25 from Babar indicates an Early Cretaceous (Albian) 20 484 MDA. Sample BAB 25 yielded only 17 concordant analyses and the ages must be ²² 485 interpreted with care as some populations are likely to have been missed but the 25 **486** ages are comparable to populations in Cretaceous sandstones from Tanimbar and ²⁷ **487** East Timor (Fig. 10A). The sample consists of 70.6% Phanerozoic, 23.5% ₃₀ **488** Proterozoic and 5.9% Archean ages. Most abundant ages are Permo-Triassic ³² 489 (41.2%), Paleoproterozoic (23.5%) and Cretaceous (11.8%).

³⁵ 490 Cretaceous samples from Tanimbar belong to the Upper Sandstone 38 **491** Member of the Ungar Formation. Charlton et al. (2009) suggested an Early 40 492 Cretaceous age which is supported by palynomorph analyses (INPEX, pers. comm., 493 2012). Combined Cretaceous samples yielded 429 concordant analyses from 45 **494** samples TAN 11 (MDA 84.6±1 Ma), TAN 28 (MDA 83.7±1 Ma), TAN 31 (89.5±1 Ma) ⁴⁷ 495 and TAN 45 (103.6±2 Ma). The youngest zircon ages within this group indicate an 50 **496** MDA of Santonian (Late Cretaceous), and an age range of Santonian to Albian, ⁵² **497** which is younger than previously suggested. Cretaceous samples contain zircons of 55 **498** 54.3% Phanerozoic, 43.8% Proterozoic and 1.9% Archean ages. The most abundant 57 **499** age populations are Permo-Triassic (21.2%), Neoproterozoic (20.3%) and Cambrian ₆₀ 500 to Carboniferous (15.6%). Important Cretaceous (10.3%) and Jurassic (~8%)

1 2

3 4

5

б

8 9

11 12

14

16 17

18 19

21

23 24

26

28 29

31

33 34

36 37

39

41 42

43 44

46

48 49

51

53 54

56

58

501 populations are also present. The main peaks are at 100, 150-170, 240-280, 550, 502 1200 and 1800 Ma (Fig. 10A). It is striking that most zircons show subhedral and 5 **503** euhedral grain morphologies for Mesozoic and Paleozoic populations (Fig. 10B).

6 7

1 2

3 4

6. Discussion

Quartz-rich sandstones from Sumba, West Timor, East Timor, Babar and Tanimbar vary in light and heavy mineral compositions, which generally indicate mixed sources (Fig. 8). Slight differences in the modal compositions and textural analyses suggest derivation from a magmatic arc for Sumba and West Timor (immature to mature character with a mix of euhedral and rounded grains) and a recycled orogen for East Timor and Tanimbar, where morphologies are predominantly rounded.

Heavy mineral compositions are dominated by high concentrations of metamorphic minerals (mainly garnet and andalusite) in Sumba and West Timor (Fig. 9). In contrast, East Timor and Tanimbar contain high abundances of rounded zircon and tourmaline which supports the multiply recycled sedimentary character of grains from an acidic igneous source.

Zircon populations in Cretaceous sandstones of the Banda Arc islands resemble each other (Fig. 10). All islands contain Cretaceous zircons that are predominantly euhedral and subhedral, which suggests nearby volcanic sources at the time of deposition. The general distribution of age populations and corresponding grain morphologies suggest a mixture of different sources. However, differences between Sumba-West Timor (SWT) and Tanimbar-East Timor (TET) are noteworthy. SWT shows common Paleoproterozoic (1.5-1.8 Ga), Mesoproterozoic

525 (1.2 Ga) and Neoproterozoic (500-650 Ma) peaks, but in TET the 1.8 Ga population 526 is significantly lower. The most striking feature in TET is the abundance of Jurassic 5 **527** zircons that are missing in SWT and in Jurassic sandstones, such as the Lower 528 Sandstone Member of the Ungar Formation, shown in Fig. 11 (from Zimmermann ⁹ 529 and Hall, 2016). The 550 Ma peak is present in all Cretaceous samples, but missing ¹² **530** in most Triassic and Jurassic formations. The Cretaceous sandstone from Babar 15 **531** yielded only 17 concordant grains, including 2 Cretaceous ages. Lithologically it 17 **532** resembles the Sumba Lasipu Formation, but the small number of zircons includes a ¹⁹₂₀ 533 Jurassic grain, and this sample therefore resembles East Timor and Tanimbar 22 **534** sandstones more closely rather than those from Sumba and West Timor. ²⁴ 535 Furthermore, there are abundant Permian-Triassic and Cambrian to Carboniferous ₂₇ 536 populations in TET, but grains of this age are of minor significance in SWT (Fig. 10 ²⁹ **537** and Fig. 11).

³² **538**

1 2

3 4

б 7

8

11

13

14

16

18

21

23

25 26

28

30 31

33 34

36 37 38

40

42 43

44 45

47

49 50

52

54

³⁵ 539 Possible Sources 6.1

₃₉ 540 Fig. 12 shows zircon age histograms for SE Asian units that contain Cretaceous and Jurassic zircons, such as the Schwaner Mountains in Borneo 41 **541** 542 (Davies et al., 2014) and Central Sulawesi, including inherited ages from 46 **543** metamorphic and S-type granitoids (Hennig et al., 2016). In contrast, autochthonous ⁴⁸ 544 Cretaceous deposits that remained in Australia, such as well samples from the ₅₁ 545 Exmouth Plateau and Caswell Plateau (Southgate et al., 2011; Lewis and Sircombe, ⁵³ 546 2013), do not contain any, or very few, Phanerozoic zircons.

- 55 ⁵⁶ 547
- 57 58
- 59
- 60 61
- 62
- 63 64
- 65

6.1.1 Pre-Breakup sources

549 Provenance and geochronology studies of Triassic and Jurassic sandstones from autochthonous formations in the Banda Arc, shown in Fig. 11 [.] 8 **551** (Zimmermann and Hall, 2016), together with studies along the Australian NW Shelf 10 552 (Southgate et al., 2011; Lewis and Sircombe, 2013) (Fig. 12), the Bird's Head (Gunawan et al., 2012) and Central Sulawesi (Hennig et al., 2016) are a useful guide to linking sandstones to likely sources in Western Australia, Central Australia and the Bird's Head region (Fig. 13).

6.1.2 Post- Breakup sources

Continental fragments rifted from the northern Australian continent in the Late Jurassic (Fig. 3). Collision of these allochthonous fragments in the Cretaceous added the Argo, Banda and Inner Banda Blocks to the Sundaland margin (Fig. 4). After their arrival, the siliciclastic sediments deposited had sources in SE Asia including igneous rocks and reworked older crust. Australian material (igneous, metamorphic and sedimentary) that had been transported with the fragments and included inherited Precambrian zircons from continental crust with a west Australian origin (e.g. beneath East Java), as reported by Smyth et al. (2007, 2008) and Hall et al. (2009). Jurassic volcanic activity was suggested by Zimmermann and Hall (2016), based on zircons in West Timor sandstones, with a source along the NW Shelf of Australia within the southern Banda Block-Inner Banda Block (Hennig et al., 2016). Volcanic activity was driven by the break-up of Gondwana and the subsequent fragmentation and drift of continental Australian blocks.

572 The most common Cretaceous age peaks in this study are different in ² 573 SWT and TET; they are c. 90-80 Ma in Sumba and West Timor (SWT), and c. 110-⁴ 5 574 90 Ma (minor 130 Ma) in East Timor, Babar and Tanimbar (TET). Gunawan et al. ⁷ 575 (2012) interpreted Late Cretaceous ages from the Sirga Formation (c. 88 Ma) in the ⁹ 576 Bird's Head as most likely to have been derived from a local source. Abundant ¹² **577** granites of Cretaceous age in the Borneo Schwaner Mountains (Banda Block) were 578 interpreted as having been derived from pulses of magmatic activity around 112, 98, 84 and 76 Ma (Davies et al., 2014). Common populations in the granitoids are 17 **579** ¹⁹ 20 **580** 120-100 Ma and 90-80 Ma. In contrast, Cretaceous granodiorite intrusions from Sumba are younger and were dated at 86-77 Ma and 71-56 Ma (Abdullah et al., 22 **581** ²⁴ 582 2000).

Fig. 13 shows a map that is based on previous studies in greater Australia and SE Asia, highlighting the islands investigated and possible main regions (i.e. granitoid bodies, cratons and fragments) that could have supplied material to the Banda region.

6.2 Provenance of Cretaceous sandstones in the Banda Arc

The Schwaner Mountains in Borneo (granites generally older than c. 86 44 589 ⁴⁶ 590 Ma) and granodiorite intrusions from Sumba (younger than c. 86 Ma) are the most 49 **591** probable sources of contemporaneous Cretaceous zircons (Fig. 10 and Fig. 12). ⁵¹ **592** However, older zircons must have come from other places since pre-Cretaceous 5₄ 593 zircons are absent in the north Schwaner Mountains, although a few Jurassic ⁵⁶ 594 granites are known form the south Schwaner Mountains, and pre-Cretaceous rocks are not known from Sumba. Significant Archean, Neo and Meso-Proterozoic

1

3

б

596 populations show similarities to autochthonous Triassic-Jurassic and Cretaceous ² 597 sandstones in the Banda Arc (Fig. 11) and the Australian NW Shelf (Fig. 12). These 5 **598** were mainly derived from Western and North/Central Australia and the Bird's Head. ⁷ 599 This also explains abundant Permian-Triassic and Cambrian to Carboniferous populations in TET and SWT. Cretaceous zircons are missing in the autochthonous Australian units and support a SE Asian setting for the Banda Arc Cretaceous ¹⁴ 602 sandstones that form part of the Banda Allochthon (BA) or Banda Terrane (Audley-Charles, 2011) for the individual islands: BA West Timor, BA East Timor, BA Babar 17 **603** and BA Tanimbar.

In order to estimate the possible contribution of different sources, a cumulative percentage plot in Fig. 14A highlights relative abundances of ages and probable sources. TET contains abundant Jurassic zircons that are missing in SWT and in Jurassic sediments within the Argo Block (e.g. origin of the Lower Sandstone Member of the Ungar Formation in Fig. 11). Hence, a geographical separation with different provenance is necessary (Fig. 14B). The ultimate Jurassic source was probably located within the Banda Block/Inner Banda Block.

A Cretaceous palaeogeographic reconstruction with major sediment ⁴² 613 transport directions (Fig. 14B) shows suggested sources for Cretaceous formations in the Banda Arc that are now part of the Banda Allochthon, including West Timor ⁴⁷ 615 (BA-WT), East Timor (BA-ET), Babar (BA-B), and Tanimbar (BA-T). The heavy ₅₀ 616 mineral and zircon age data suggest a syn-sedimentary Cretaceous source in the Schwaner Mountains and Sumba, together with recycling of Permian-Triassic and Jurassic pre-rift units.

Data from the Northwest Shelf of Australia (Southgate et al., 2011; Lewis and Sircombe, 2013), Java (Smyth et al., 2003, 2007), Borneo (Davies et al.,

65

1

2014) and Sulawesi (Hennig et al., 2016) were included (Fig. 14A) to highlight age similarities and assess possible reconstructions. Trends can be recognized by **623** decreasing or increasing percentages of single zircon populations. Ultimate main ⁷ 624 sources for the Banda Allochthon sandstones are a mix of Cretaceous populations ⁹ 625 derived from Sundaland (Schwaner Mountains) and zircons derived from pre-**626** Cretaceous rocks formed, or deposited before separation from greater Australia. Single grain morphologies of zircons analysed (Fig. 10B) support interpretations of contemporaneous first cycle Cretaceous grains and a Permian-Triassic signal. Rounded Proterozoic (Australian) grains are also mixed with subhedral grains indicating differences in recycling of Australian cratons.

In **Sumba**, the principal zircon populations indicate sources in Western Australia (33%), North/Central Australia (26%), Borneo/Sumba (24%) and the Bird's Head region (16%) (Fig. 14A). 57% of Cretaceous zircons have ages less than c. 86 Ma, indicating a relationship to the Sumba granodiorites and/or the later eruptive stages of the Schwaner Mountains. Based on previous research (Sallé, 1982; von der Borch et al., 1983) and new field observations the rocks are interpreted as turbidites deposited in deep water. The variation in rocks in the central part of the island and around the south and southwest coast suggest facies changes along the shelf and slope. West Timor (BA-WT) contains similar population percentages as Sumba, mainly from Western Australia (36%), North/Central Australia (33%), Borneo/Sumba (12%) and the Bird's Head (19%) (Fig. 14A). 31% of the Cretaceous population are younger than 86 Ma indicating input from Sumba granodiorites and/or the later eruptive stages of the Schwaner Mountains.

644 East Timor (BA-ET) is sourced by Western Australia (35%), the Bird's Head (31%), North/Central Australia (17%), the Banda Block (10%) and Borneo

б

646 (7%), of which >100 Ma old zircons dominate (80%), indicating a probable source in 647 the Schwaner Mountains, whereas younger (<86 Ma) Sumba ages are absent.

In **Tanimbar** (BA-T), the main sources are similar to East Timor, 648 ₈ 649 deriving from Western Australia (31%), the Bird's Head (28%), North/Central Australia (23%), the Banda Block (8%) and Borneo (10%) (Fig. 14A). The Cretaceous population is strongly dominated by >86 Ma zircons (95%), indicating a likely source in the Schwaner Mountains with very minor possible contribution from Sumba. Thus, we suggest that sediment was transported to the site of deposition mainly from the west but also from the east (Fig. 14B).

As noted before, one Cretaceous sample from **Babar** (BA-B) yields only 17 concordant U-Pb zircon ages and it is not likely that such a small number of grains provides a representative sample. However, the few ages do suggest possible provenance trends. The main features are similarities to Cretaceous samples from East Timor and Tanimbar, suggesting the geographical location of the BA-B fragment was where it is now, between BA-ET and BA-T (Fig. 14B).

Tectonic Evolution of the Banda Allochthon 6.3

Previous researchers (e.g. Wensink, 1994; Abdullah et al., 2000; Hall 663 et al., 2009; Hall, 2012; Metcalfe, 2013) proposed models with fragments derived from Australia colliding with Sundaland. Modifications of previous palaeogeographic and tectonic reconstructions (Hall, 2012) are suggested here to locations of individual fragments and evolution of sediment supply, following Triassic and Jurassic reconstructions by Zimmermann and Hall (2016). Fig. 15A displays the rifting of fragments (Argo, Banda and Inner Banda) from the Early Cretaceous (130

65

Ma) and the situation within the Sundaland margin in the Late Cretaceous (75 Ma). The figures show the drift history to the north and collision with Sundaland with interpreted locations of the Banda Allochthon fragments in the Sundaland margin at 75 Ma. Sumba and BA-WT (SWT) are situated at the southern edge of Sundaland and were supplied by sediment from the north (Borneo) and east by local erosion of the Argo Block. BA-ET, BA-B and BA-T (TET) were situated at the eastern margin. Sediment was mainly derived from the east (Borneo and West Sulawesi) and from the north (reworking of the southern edge of the inner Banda Block).

Interpreted Cenozoic movements to the present-day locations of different fragments are shown in Fig. 15B. At around 23 Ma, the most northerly part of Australia (Sula Spur) collided with Sundaland, followed by subduction hinge rollback into the Banda Embayment (Spakman and Hall, 2010) and initial fragmentation of the Sula Spur (from c. 15 Ma).

The 5 Ma palaeogeographic map shows the setting shortly before the collision of the Banda volcanic arc with the Australian continent and the resulting emplacement of the Banda Allochthon. The present-day configuration was reached by overthrusting of these units on top of Mesozoic and pre-Triassic autochthonous (Gondwanan) sediments and basement, coupled with rapid uplift that created the Outer Banda Arc Islands.

0 7. Conclusions

Earlier studies of the Northwest Shelf of Australia identified sources for Permian-Triassic and Jurassic sandstones in Central and Western Australia and the Bird's Head, depositing the pre-breakup sequence. From the Late Jurassic, rifting of

65

the Australian Gondwana margin caused fragmentation and separation into blocks which were accreted to Sundaland in the Early and Late Cretaceous.

Cretaceous sandstones were deposited unconformably on top of the ₈ 697 pre-breakup sequence at the southeastern margin of Sundaland in what was to 10 698 become the Banda Allochthon. The sandstones commonly contain significant ⁻⁻₁₃ 699 contents of volcanic quartz, feldspar and lithic fragments, indicating mixed sources **700** for the fragments that later became Sumba, BA-West Timor, BA-East Timor, BA-Babar and BA-Tanimbar. The principal source rocks are suggested to be of **702** metamorphic origin in Sumba and West Timor (SWT), and acidic igneous and ²² **703** recycled sedimentary rocks in East Timor and Tanimbar (TET). Contemporaneous **704** Cretaceous igneous sources include Sumba and the Schwaner Mountains within ²⁷ **705** Sundaland. Variations between the SWT and TET fragments suggest differences in ₃₀ 706 provenance, most likely due to geographical separation along the southern shelf of **707** Sundaland. Neogene extension further isolated these Cretaceous sandstones and Neogene collision processes moved these fragments into the Outer Banda Arc Islands, where they are overthrust and now re-incorporated in the Australia margin.

Acknowledgements

This study was funded by the SE Asia Research Group at Royal ⁴⁹ 713 Holloway University of London supported by a consortium of oil companies including **714** BP, ENI, GDF Suez, Inpex, Murphy, Niko, Repsol, Shell and Statoil. Great thanks for assistance during fieldwork to Afif Saputra (Inpex), Institut Teknologi Bandung (ITB) **716** and the State Ministry of Research and Technology (RISTEK). We thank Martin Rittner and Andy Beard (UCL/Birkbeck College) for help and support at the LA-ICP-

MS and CL facility and Anna Bird and Dominique Tanner for help with heavy minerals and SEM work at Royal Holloway. Also, many thanks to Marcelle BouDagher-Fadel for microfossil analyses at University College London. We are grateful for access to unpublished material from the Tanimbar Islands that was provided by INPEX and JOGMEC.

Appendix A. Supplementary data

Supplementary data to this article can be found online. Supplementary Data File 1 contains data tables of sampling locations, formations and lithologies. Supplementary Data File 2 contains data tables of modal compositions and textural analyses. Supplementary Data File 3 contains data tables of heavy mineral analyses. Supplementary Data File 4 contains data tables of LA-ICP-MS analyses.

29 References

- Abdullah, C.I., Rampnoux, J.-P., Bellon, H., Maury, R.C., Soeria-Atmadja, R., 2000.
 The evolution of Sumba Island (Indonesia) revisited in the light of new data on the geochronology and geochemistry of the magmatic rocks. Journal of Asian Earth Sciences 18, 533-546.
- Armandita, C., Morley, C. K., & Rowell, P., 2015. Origin, structural geometry, and
 development of a giant coherent slide: The South Makassar Strait mass
 transport complex. Geosphere, 11(2), 376-403.
 - Audley-Charles, M.G., 1968. The geology of Portuguese Timor. Geological Society
 of London, Memoir 4, 76 pp.
- 49739Audley-Charles, M.G., 1986. Timor–Tanimbar Trough: the foreland basin to the51740evolving Banda orogen. In: Allen, P.A., Homewood, P. (Eds.), Foreland5253741Basins. International Association of Sedimentologists, Special Publication 8,5455742pp. 91-102.
- Audley-Charles, M.G., 2011. Tectonic post-collision processes in Timor. In: Hall, R., Cottam, M.A. & Wilson, M.E.J. (Eds.), The SE Asian Gateway: History and

- 745 Tectonics of the Australia-Asia collision. Geological Society of London Special 1 2 **746** Publication, 355, 241-266
- 3 747 Audley-Charles, M.G., Ballantyne, P.D., Hall, R., 1988. Mesozoic-Cenozoic rift drift 4 5 sequence of Asian Fragments from Gondwanaland. Tectonophysics 155, 317-748 б ⁷ 749 330. 8
- 9 750 Audley-Charles, M.G., Harris, R.A., 1990. Allochthonous terranes of the southwest 10 11 **751** Pacific and Indonesia. Philosophical Transactions Royal Society of London 12 ₁₃ 752 331, 571-587.
- 15¹¹753 14 Barber, A.J., 1979. Structural interpretation of the island of Timor, east Indonesia. ¹⁶ 754 United Nations ESCAP CCOP / SEATAR, Ad Hoc meeting on the geology 17 ¹⁸ 755 and tectonics of Eastern Indonesia, Bandung. 19
- 20 756 Barber, A.J., Crow, M.J., 2009. The structure of Sumatra and its implications for the 21 22 **757** tectonic assembly of Southeast Asia and the destruction of Paleotethys. 23 ⁻³₂₄ 758 Island Arc 18, 3-20.
- 25 Barber, P., Carter, P., Fraser, T., Baillie, P. & Myers, K. 2003. Paleozoic and 759 26 ²⁷ **760** Mesozoic petroleum systems in the Timor and Arafura Seas, Eastern 28 ²⁹ 761 Indonesia. Indonesian Petroleum Association, Proceedings 29th Annual 30 31 **762** Convention 485-500.
- 32 33 **763** Bergman, S. C., Coffield, D. Q., Talbot, J. P., & Garrard, R. J., 1996. Tertiary tectonic 34 35 **764** and magmatic evolution of Western Sulawesi and the Makassar Strait, 36 Indonesia: Evidence for a Miocene continent-continent collision. In R. Hall & 765 37 ³⁸ 766 D. J. Blundell (Eds.), Tectonic Evolution of SE Asia (Vol. 106, pp. 391-430). 39
- ⁴⁰ 767 Bishop, M.G., 1999. Total petroleum systems of the Bonaparte Gulf Basin area, 41 42 **768** Australia; Jurassic, Early Cretaceous-Mesozoic; Keyling, Hyland Bay-43 44 **769** Permian; Milligans-Carboniferous, Permian (No. 99-50-P). US Dept. of the 45 46 **770** Interior, US Geological Survey.
- 47 771 Bowin, C., Purdy, G.M., Johnston, C., Shor, G., Lawver, L., Hartono, H.M.S., Jezek, 48 ⁴⁹ 772 P., 1980. Arc-continent collision in the Banda Sea region. American 50 Association of Petroleum Geologists Bulletin 64, 868-918. 51 **773** 52
- 53 **774** Brown, M., Earle, M.M., 1983. Cordierite-bearing schists and gneisses from Timor, 55 **775** eastern Indonesia: P-T implications of metamorphism and tectonic 57 **776** 56 implications. Journal of Metamorphic Geology 1, 183-203.
- 777 Burollet, P.F., Salle, C., 1982. Histoire geologique de l'ile de Sumba (Indonesie). ⁶⁰ 778 Bulletin de la Société géologique de France 24, 573-580.
- 63 64 65

58

59

- 779 Camplin, D. J., & Hall, R., 2014. Neogene history of Bone Gulf, Sulawesi, Indonesia. ² 780 Marine and Petroleum Geology, 57, 88-108.
- 3 781 Carter, D.J., Audley-Charles, M.G., Barber, A.J., 1976. Stratigraphical analysis of 4 5 782 island arc-continental margin collision in eastern Indonesia. Journal of the б ⁷ 783 Geological Society 132, 179-198. 8
- 9 784 Cawood, P.A., Nemchin, A.A., Freeman, M., Sircombe, K., 2003. Linking source and 10 11 **785** sedimentary basin: Detrital zircon record of sediment flux along a modern 12 ₁₃ 786 river system and implications for provenance studies. Earth and Planetary 14 787 Science Letters 210, 259-268. 15
- 16 Cawood, P.A., Nemchin, A.A., Leverenz, A., Saeed, A., Ballance, P.F., 1999. U/Pb 788 17 ¹⁸ 789 dating of detrital zircons: Implications for the provenance record of Gondwana 19 20 **790** margin terranes. Geological Society of America Bulletin 111, 1107-1119. 21
- 22 **791** Charlton, T.R., 1987. The tectonic evolution of the Kolbano-Timor Trough ⁻³₂₄ 792 Accretionary Complex, Timor, Indonesia. PhD Thesis, University of London.
- 793 Charlton, T.R., Barber, A.J., McGowan, A.J., Nicoll, R.S., Roniewicz, E., Cook, S.E., ²⁷ **794** Barkham, S.T., Bird, P.R., 2009. The Triassic of Timor: Lithostratigraphy, ²⁹ 795 chronostratigraphy and palaeogeography. Journal of Asian Earth Sciences 31 **796** 36, 341-363.
- 33 **797** Charlton, T.R., De Smet, M.E.M., Samodra, H., Kaye, S.J., 1991. The stratigraphic 35 **798** and structural evolution of the Tanimbar islands, eastern Indonesia. Journal of 799 Southeast Asian Earth Sciences 6, 343-358.
- ³⁸ 800 Charlton, T.R., 2001. Permo-Triassic evolution of Gondwanan eastern Indonesia, ⁴⁰ 801 and the final Mesozoic separation of SE Asia from Australia. Journal of Asian 42 **802** Earth Sciences 19, 595-617.
- 43 44 803 Charlton, T.R., 2012. Permian-Jurassic Palaeogeography of the SE Banda Arc 45 4₆¹⁰ 804 Region. Berita Sedimentologi No. 24, 5-17.
- 47 805 Clements, B., Hall, R., Smyth, H. R., & Cottam, M. A., 2009. Thrusting of a volcanic 48 ⁴⁹ 806 arc: A new structural model for Java. Petroleum Geoscience, 15, 159-174. 50
- 51 **807** Davies, L., Hall, R., Armstrong, R., 2014. Cretaceous crust in SW Borneo: 52 petrological, geochemical and geochronological constraints from the 53 **808** 54 55 **809** Schwaner Mountains. Proceedings Indonesian Petroleum Association, 38th 56 57 **810** Annual Convention and Exhibition, IPA14-G-025.
- 58 Deighton, I., Hancock, T., Hudson, G., Tamannai, M., Conn, P., & Oh, K., 2011. Infill 811 59 ⁶⁰ 812 seismic in the southeast Java forearc basin: Implications for petroleum 61
- 63 64 65

1

23

25

26

28

30

32

34

36

37

39

- 813prospectivity. Proceedings Indonesian Petroleum Association 35th Annual $\frac{1}{2}$ 814Convention, IPA11-G-068.
- ³/₄815 Dickinson, W.R., Beard, L.S., Brakenridge, G.R., Erjavic, J.L., Ferguson, R.C.,
 ⁵/₆816 Inman, K.F., Knepp, R.A., Lindeberg, F.A., Ryberg, P.T., 1983. Provenance of
 ⁷/₈817 North American Phanerozoic sandstones in relation to tectonic setting.
 ⁹/₈818 Geological Society of America Bulletin 94, 222-235.
- 11 819Dickinson, W.R., Gehrels, G.E., 2009. Use of U–Pb ages of detrital zircons to infer12
13 820maximum depositional ages of strata: a test against a Colorado Plateau14
15Mesozoic database. Earth and Planetary Science Letters 288, 115-125.
- ¹⁶ 822 Dickinson, W.R., Suczek, C.A., 1979. Plate tectonics and sandstone composition.
 ¹⁸ 823 American Association of Petroleum Geologists Bulletin 63, 2164-2182.
- Elburg, M., & Foden, J., 1999. Sources for magmatism in central Sulawesi:
 geochemical and Sr- Nd-Pb isotopic constraints. Chemical Geology, 156(1-4),
 67-93.
- Elburg, M., van Leeuwen, T., Foden, J., & Muhardjo., 2003. Spatial and temporal isotopic domains of contrasting igneous suites in Western and Northern Sulawesi, Indonesia. Chemical Geology, 199(3-4), 243-276.
- Emmet, P. A., Granath, J. W., & Dinkelman, M. G., 2009. Pre-Tertiary sedimentary (keels" provide insights into tectonic assembly of basement terranes and present-day petroleum systems of the East Java Sea. Proceedings Indonesian Petroleum Association 33rd Annual Convention, IPA09-G-046 (Note: 10.11) (Not:
- ⁴⁰ 835 Fedo, C.M., Sircombe, K.N., Rainbird, R.H., 2003. Detrital zircon analysis of the
 ⁴¹ sedimentary record. Reviews in Mineralogy and Geochemistry 53, 277-303.
- Feo-Codecido, G., 1956. Heavy-mineral techniques and their application to
 Venezuelan stratigraphy. AAPG Bulletin 40, 984-1000.
- 47
48839Galehouse, J.S., 1971. Point counting. In: Carver, R.E. (Ed.), Procedures in49
50840sedimentary petrology, Wiley Interscience, New York, pp. 385-407.
- S1 841 Gehrels, G., Valencia, V., Pullen, A., 2006. Detrital zircon geochronology by laser ablation multicollector ICPMS at the Arizona LaserChron Center.
 Paleontological Society Papers 12, 67.
- Gibbons, A.D., Barckhausen, U., van den Bogaard, P., Hoernle, K., Werner, R., Whittaker, J.M. & Müller, R.D. 2012. Constraining the Jurassic extent of
- 60 61 62

19

- 846 Greater India: Tectonic evolution of the West Australian margin. 1 2 **847** Geochemistry, Geophysics, Geosystems 13 (5), Q05W13.
- 3 848 Goldstein, S., Arndt, N.T., Stallard, R.F., 1997. The history of a continent from U-Pb 4 5 849 ages of zircons from Orinoco River sand and Sm-Nd isotopes in Orinoco 6 7 850 basin river sediments. Chemical Geology 139, 271-286. 8
- Gradstein, F.M., Ogg, J.G., Schmitz, M., Ogg, G. 2012. The Geologic Time Scale 9 **851** 10 11 **852** 2012, Elsevier.
- 12 ₁₃ 853 Granath, J. W., Christ, J. M., Emmet, P. A., & Dinkelman, M. G., 2011. Pre-Cenozoic 14 15 **854** sedimentary section and structure as reflected in the JavaSPAN crustal-scale ¹⁶ 855 PSDM seismic survey, and its implications regarding the basement terranes in 17 ¹⁸ 856 the East Java Sea. In R. Hall, M. A. Cottam, & M. E. J. Wilson (Eds.), The SE 19 Asian Gateway: History and Tectonics of the Australia-Asia collision (Vol. 355, 20 **857** 21 22 **858** pp. 53-74).
- 23 ⁻⁰₂₄ 859 Gunawan, I., Hall, R. & Sevastjanova, I., 2012. Age, character and provenance of 25 the Tipuma Formation, West Papua: New insights from detrital zircon dating. 860 26 ²⁷ 861 Proceedings Indonesian Petroleum Association 36th Annual Convention, 28 ²⁹ 862 IPA12-G-027 1-14. 30
- Haile, N. S., 1974. Borneo. In A. M. Spencer (Ed.), Mesozoic-Cenozoic Orogenic 31 **863** 33 **864** Belts (Vol. 4, pp. 333-347).
- 34 35 **865** Haile, N. S., McElhinny, M. W., & McDougall, I., 1977. Palaeomagnetic data and 36 radiometric ages from the Cretaceous of West Kalimantan (Borneo), and their 866 37 ³⁸ 867 significance in interpreting regional structure. Journal of the Geological 39 ⁴⁰ 868 Society of London, 133, 133-144. 41
- 42 **869** Hall, R., 1996. Reconstructing Cenozoic SE Asia. In: Hall, R. & Blundell, D.J. (Eds.), 43 44 **870** Tectonic Evolution of SE Asia. Geological Society of London Special 45 46 **871** Publication 106, 153-184.
- 47 Hall, R., 2002. Cenozoic geological and plate tectonic evolution of SE Asia and the 872 48 ⁴⁹ 873 SW Pacific: computer-based reconstructions, model and animations. Journal 50 51 **874** of Asian Earth Sciences 20, 353-431. 52
- 53 **875** Hall, R., 2011. Australia-SE Asia collision: plate tectonics and crustal flow. In: Hall, 54 55 **876** R., Cottam, M.A., Wilson, M.E.J. (Eds.), The SE Asian Gateway: History and 56 57 **877** Tectonics of the Australia-Asia collision. Geological Society of London Special 58 878 Publication 355, 75-109. 59
- 60

- 61 62
- 63 64

- Hall, R., 2012. Late Jurassic-Cenozoic reconstructions of the Indonesian region and 879 ² 880 the Indian Ocean. Tectonophysics 570, 1-41.
- 3 881 Hall, R., Clements, B., Smyth, H.R., 2009. Sundaland: Basement character, structure 4 5 882 and plate tectonic development. Proceedings Indonesian Petroleum б ⁷ 883 Association, 33rd Annual Convention. 8
- Hall, R., Cottam, M.A., Wilson, M.E.J., 2011. The SE Asian gateway: history and 9 884 10 11 885tectonics of Australia-Asia collision. In: Hall, R., Cottam, M.A., Wilson, M.E.J. 12 ₁₃ 886 (Eds.), The SE Asian Gateway: History and Tectonics of the Australia-Asia 15 **887** 14 collision. Geological Society of London Special Publication 355, 1-6.
- ¹⁶ 888 Hall, R., Sevastjanova, I., 2012. Australian crust in Indonesia. Australian Journal of 17 ¹⁸ 889 Earth Sciences 59, 827-844. 19
- Hamilton, W., 1979. Tectonics of the Indonesian region. U.S.G.S. Prof. Paper 1078, 20 **890** 21 22 **891** 345pp.
- 23 ⁻⁰₂₄ 892 Harris, R.A., 1989. Processes of allochthon emplacement with special reference to 25 893 the Brooks Range ophiolite. Alaska and Timor, Indonesia. Ph.D. thesis, 26 ²⁷ 894 University of London. 28
- ²⁹ 895 Harris, R.A., 1991. Temporal distribution of strain in the active Banda orogen: a 31 **896** reconciliation of rival hypotheses. Journal of Southeast Asian Earth Sciences, 33 **897** 6, 373-386.
- 34 35 **898** Harris, R., 2006. Rise and fall of the Eastern Great Indonesian arc recorded by the 36 899 assembly, dispersion and accretion of the Banda Terrane, Timor. Gondwana 37 ³⁸ 900 Research 10, 207-231. 39
- 40 **901** Hasan, K. 1991. Upper Cretaceous flysch succession of the Balangbaru Formation, 42 **902** Southwest Sulawesi. Indonesian Petroleum Association, Proceedings 20th 44 903 Annual Convention, Jakarta, 183-208.
- 45 46 **904** Heine, C., Müller, R.D., 2005. Late Jurassic rifting along the Australian North West 47 Shelf: margin geometry and spreading ridge configuration. Australian Journal 905 48 ⁴⁹ 906 of Earth Sciences 52, 27-39. 50
- Heine, C., Quevedo, L., McKay, H., Müller, R.D., 2012. Plate tectonic consequences 51 **907** 52 53 **908** of competing models for the origin and history of the Banda Sea subducted 54 55 **909** oceanic lithosphere. arXiv preprint arXiv:1210.4958.
- 56 50 57 **910** Hennig, J., Hall, R., Armstrong, R.A., 2016. U-Pb zircon geochronology of rocks from ⁵⁸ 911 west Central Sulawesi, Indonesia: Extension-related metamorphism and
- 59 60 61

30

32

41

43

62 63 64

- 912 magmatism during the early stages of mountain building. Gondwana 1 ² 913 Research 32, 41–63.
- ³₄ 914 Hennig, J., Breitfeld, H. T., Hall, R., & Surya Nugraha, A., 2017. The Mesozoic ⁵ 915 tectono-magmatic evolution at the Paleo-Pacific subduction zone in West б ⁷ 916 Borneo. Gondwana Research, 48, 292-310. 8
- Hilton, D. R., Hoogewerff, J. A., van Bergen, M. J., & Hammerschmidt, K., 1992. 9 917 10 $11 \ 918$ Mapping magma sources in the east Sunda-Banda arcs, Indonesia: 12 13 **919** constraints from Helium isotopes. Geochimica and Cosmochimica Acta, 56(2), ¹⁴ 920 851-859.
- ¹⁶ 921 Hinschberger, F., Malod, J.-A., Rehault, J.-P., Villeneuve, M., Royer, J.-Y., 17 ¹⁸ 922 Burhanuddin, S., 2005. Late Cenozoic geodynamic evolution of eastern 19 Indonesia. Tectonophysics 404, 91-118. 20 **923** 21
- 22 **924** Hutchison, C.S., 1989. Geological evolution of South-east Asia (Vol. 13, p. 368). 23 24 **925** Oxford: Clarendon Press.
- ²⁵₂₆ 926 25 Jasin, B. and Haile, N., 1996. Uppermost Jurassic-Lower Cretaceous radiolarian ²⁷ 927 chert from the Tanimbar Islands (Banda Arc), Indonesia. Journal of Southeast 28 ²⁹ 928 Asian Earth Sciences, 14(1-2), pp.91-100. 30
- 31 **929** Johansen, K. B., Maingarm, S., & Pickard, A., 2007. Hydrocarbon potential of the 33 **930** South Makassar Basin. Proceedings of the 2007 South East Asia Petroleum 35 **931** Exploration Society Conference, 1-46.
- ³⁶ 37 **932** Katili, J. A., 1971. A review of the geotectonic theories and tectonic maps of ³⁸ 933 Indonesia. Earth-Science Reviews, 7, 143-163. 39
- ⁴⁰ **934** Katili, J. A., 1973. Geochronology of west Indonesia and its implication on plate 42 **935** tectonics. Tectonophysics, 19, 195-212.
- 43 44 936 Kaye, S.J., 1989. The structure of Eastern Indonesia. PhD thesis, University of 45 46 **937** London.
- 47 938 Klompé, T.H.F., 1954. The structural importance of the Sula Spur (Indonesia). 48 ⁴⁹ 939 Indonesian Journal of Natural Sciences 110, 21-40. 50
- 51 **940** Kupecz, J., Sayers, I., Tognini, P., Hilman, A., Tanos, C., & Ariyono, D., 2013. New 52 53 **941** insights into the tectono-stratigraphic evolution of the South Makassar Basin. 54 55 **942** Indonesian Petroleum Association, Proceedings 37th Annual Convention, 50 57 **943** 56 IPA13-G-158 151-141.
- ⁵⁸ 944 Lewis, C.J., Sircombe, K., 2013. Use of U-Pb geochronology to delineate 59 ⁶⁰ 945 provenance of North West Shelf sediments, Australia. In: Keep, M., Moss, 61
- 63 64 65

32

34

- 946 S.J., (Editors), The Sedimentary Basins of Western Australia IV, Proceedings 1 2 **947** of the Petroleum Exploration Society of Australia Symposium, Perth, WA, 3 948 2013, 1-27. 4
- 5 Longley, I.M., Buessenschuett, C., Clydsdale, L., Cubitt, C.J., Davis, R.C., Johnson, 949 б ⁷ 950 M.K., Marshall, N.M., Murr, A.P., 2002. The North West Shelf of Australia - a 8 Woodside perspective. In: Keep, M., Moss, S.J. (Eds.), The Sedimentary ⁹ 951 10 11 **952** Basins of Western Australia III, Proceedings West Australian Basins 12 13 **953** Symposium, Perth 2002, Petroleum Exploration Society of Australia, pp. 28-14 15 **954** 86.
- ¹⁶ 955 Luyendyk, B. P., 1974. Gondwanaland dispersal and the early formation of the 17 ¹⁸ 956 Indian Ocean. In B. P. Luyendyk & T. A. Davies (Eds.), Deep Sea Drilling 19 Project, Initial Reports (Vol. 26, pp. 945-952). Washington: US Government 20 **957** 21 22 **958** Printing Office.
- 23 24 **959** Mange, M.A., 2002. New look at heavy minerals - a short course. 16th International 25 960 Sedimentological Congress, Rand Afrikaans University, South Africa, 34 pp. 26
- ²⁷ 961 Mange, M.A., Maurer, H.F.W., 1992. Heavy minerals in colour. Chapman & Hall, ²⁹ 962 London, 147 pp.
- 31 **963** Manur, H., & Barraclough, R., 1994. Structural control on hydrocarbon habitat in the 33 **964** Bawean area, East Java Sea. Proceedings Indonesian Petroleum Association 35 **965** 23rd Annual Convention, 129-144.
- 36 Maulana, A., Christy, A. G., Ellis, D. J., Imai, A., & Watanabe, K., 2013. 966 37 ³⁸ 967 Geochemistry of eclogite- and blueschist-facies rocks from the Bantimala 39 40 968 Complex, South Sulawesi, Indonesia: Protolith origin and tectonic setting. 41 42 **969** Island Arc, 22(4), 427-452. 43
- 44 970 Maulana, A., Ellis, D. J., & Christy, A. G., 2010. Petrology, geochemistry and tectonic 45 46 **971** evolution of the South Sulawesi basement rocks, Indonesia. Proceedings 47 972 Indonesian Petroleum Association 34th Annual Convention, IPA10-G-192 48 ⁴⁹ 973 191-126. 50
- 51 **974** Metcalfe, I., 1988. Origin and assembly of Southeast Asian continental terranes. In 52 53 **975** M. G. Audley-Charles & A. Hallam (Eds.), Gondwana and Tethys (Vol. 37, pp. 54 55 **976** 101-118).
- 57 **977** Metcalfe, I., 1990. Allochthonous terrane processes in Southeast Asia. Philosophical ⁵⁸ 978 Transactions Royal Society of London, A331, 625-640.
- 60 61 62

56

28

30

32

- 63
- 64 65

- 979 Metcalfe, I., 1996. Pre-Cretaceous evolution of SE Asian terranes. In R. Hall & D. J. ² 980 Blundell (Eds.), Tectonic Evolution of SE Asia (Vol. 106, pp. 97-122).
- 3 981 Metcalfe, I. 1998. Palaeozoic and Mesozoic geological evolution of the SE Asian 4 5 982 region: multidisciplinary constraints and implications for biogeography. In Hall, б ⁷ 983 R. & Holloway, J. D. (eds.) Biogeography and Geological Evolution of SE 8 Asia. Backhuys Publishers, Leiden, The Netherlands, 25-41. 9 984
- Metcalfe, I., 2009. Late Palaeozoic and Mesozoic tectonic and palaeogeographical 11 985 ₁₃ 986 evolution of SE Asia. In E. Buffetaut, G. Cuny, J. Le Loeuff, & V. Suteethorn 15 **987** (Eds.), Late Palaeozoic and Mesozoic Ecosystems in SE Asia (Vol. 315, pp. ¹⁶ 988 7-23).
- ¹⁸ 989 Metcalfe, I., 2011. Tectonic framework and Phanerozoic evolution of Sundaland. Gondwana Research 19, 3-21. 20 **990**
- 21 22 **991** Metcalfe, I., 2013. Gondwana dispersion and Asian accretion: Tectonic and 23 ⁻³₂₄ 992 palaeogeographic evolution of eastern Tethys. Journal of Asian Earth ²³₂₆ 993 25 Sciences 66, 1-33.
- ²⁷ 994 Miyazaki, K., Sopaheluwakan, J., Zulkarnain, I., & Wakita, K., 1998. A jadeite-quartz-²⁹ 995 glaucophane rock from Karangsambung, central Java, Indonesia. Island Arc, 31 **996** 7(1-2), 223-230.
- 33 **997** Miyazaki, K., Zulkarnain, I., Sopaheluwakan, J., & Wakita, K., 1996. Pressure-35 **998** temperature conditions and retrograde paths of eclogites, garnet-glaucophane ³⁶ 37 **999** rocks and schists from South Sulawesi, Indonesia. Journal of Metamorphic ³⁸1000 39 Geology, 14, 549-563.
- 401001 Murphy, J.B., Fernández-Suárez, J., Keppie, J.D., Jeffries, T.E., 2004. Contiguous 41 42**1002** rather than discrete Paleozoic histories for the Avalon and Meguma terranes 43 44**1003** based on detrital zircon data. Geology 32, 585-588.

45 46**1004** Nichols, G., 2009. Sedimentology and stratigraphy. John Wiley & Sons.

- ⁴⁷481005 Norvick, M.S., 1979. The tectonic history of the Banda Arcs, eastern Indonesia: a ⁴⁹1006 review. Journal of the Geological Society 136, 519-526.
- 51**1007** Nugraha, A. M. S., & Hall, R., 2012. Cenozoic history of the East Java Forearc. 52 53**1008** Proceedings Indonesian Petroleum Association 36th Annual Convention, 54 55**1009** IPA12-G-028 021-021.
- ⁵⁶ 57**1010** Nugraha, A. M. S., & Hall, R., 2018. Late Cenozoic palaeogeography of Sulawesi, ⁵⁸1011 Indonesia. Palaeogeography, Palaeoclimatology, Palaeoecology, 490, 191-⁶⁰1012 209.
- 61 62 63 64 65

1

10

12

17

19

28

30

32

- 1013Packham, G. 1996. Cenozoic SE Asia: Reconstructing its aggregation and
reorganisation. In: Hall, R. & Blundell, D.J. (Eds.), Tectonic Evolution of SE $^{1}_{2}$ 1014Asia. Geological Society of London Special Publication 106, 123-152.
- ⁵1016 Parkinson, C.D., Miyazaki, K., Wakita, K., Barber, A.J., Carswell., D.A., 1998. An ⁷1017 overview and tectonic synthesis of the pre-Tertiary very-high-pressure ⁸1018 metamorphic and associated rocks of Java, Sulawesi and Kalimantan, ¹⁰11019 Indonesia. The Island Arc 7, 184-200.
- Pearce, N.J.G., Perkins, W.T., Westgate, J.A., Gorton, M.P., Jackson, S.E., Neal, C.R., Chenery, S.P., 1997. A compilation of new and published major and trace element data for NIST SRM 610 and NIST SRM 612 glass reference materials. Geostandards newsletter 21, 115-144.
- Polvé, M., Maury, R., Vidal, P., Priadi, B., Bellon, H., Soeria-Atmadja, R., . . . Cotten,
 J., 2001. Melting of lower continental crust in a young post-collision setting; a
 geochemical study of Plio-Quaternary acidic magmatism from central
 Sulawesi (Indonesia) (Vol. 172).
- Polvé, M., Maury, R. C., Bellon, H., Rangin, C., Priadi, B., Yuwono, S., . . . Atmadja,
 R. S., 1997. Magmatic evolution of Sulawesi (Indonesia): constraints on the
 Cenozoic geodynamic history of the Sundaland active margin.
 Tectonophysics, 272(1), 69-92.
- ³⁴₃₅1032 Priadi, B., 1993. Geodynamic implications of Neogene potassic calc-alkaline magmatism in Central Sulawesi : Geochemical and isotopic constraints.
 ³⁶₃₇1033 Proceedings of the 22nd Annual Convention of the Indonesian Association of Geologists, 1993, 59-81.
- Priadi, B., Polve, M., Maury, R. C., Bellon, H., Soeria-Atmadia, R., Joron, J. L., &
 Cotten, J., 1994. Tertiary and Quaternary magmatism in Central Sulawesi:
 chronological and petrological constraints. Journal of Southeast Asian Earth
 Sciences, 9(1/2), 81-93.
- Rangin, C., Jolivet, L., Pubellier, M., 1990. A simple model for the tectonic evolution
 of southeast Asia and Indonesia region for the past 43 m.y. Bulletin de la
 Société géologique de France 8, 889-905.
- ⁵⁴
551043Ricou, L. E., 1994. Tethys reconstructed plates, continental fragments and their⁵⁶
571044boundaries since 260 Ma from Central America to Southeastern Asia.⁵⁸
591045Geodinamica Acta, 7(4), 169-218.
- 60 61
- 62 63 64

- 1046Rigg, J. W. D., & Hall, R., 2012. Neogene development of the Savu Forearc Basin,1121047Indonesia. Marine and Petroleum Geology, 32, 76-94.
- ³A1048 Satyana, A. H., 2015. Rifting history of the Makassar Straits: New constraints from ⁵B1049 wells penetrating the basement and oils discovered in Eocene section -⁷B1050 implications for further exploration of West Sulawesi offshore. Indonesian ⁸B1051 Petroleum Association, Proceedings 39th Annual Convention, IPA15-G-104 ¹⁰D11052 101-134.
- ¹²₁₃1053 Sawyer, R.K., Sani, K., Brown, S., 1993. The stratigraphy and sedimentology of
 ¹⁴₁₅1054 West Timor, Indonesia. Indonesian Petroleum Association, Proceedings 22nd
 ¹⁶₁₇1055 annual convention, Jakarta, 1993 1, 533-574.
- 181056
19Schiller, D. M., Garrad, R. A., & Prasetyo, L., 1991. Eocene submarine fan
sedimentation in southwest Java. Proceedings Indonesian Petroleum
Association 20th Annual Convention, 125-182.
- ²²¹OSG Association 20th Annual Convention, 125-102.
 ²³241059 Schoene, B., 2014. 4.10 U–Th–Pb Geochronology, Treatise on Geochemistry (Second Edition), Elsevier, Oxford, pp. 341-378.
 ²⁷1061 Sevastjanova, I., Clements, B., Hall, R., Belousova, E.A., Griffin, W.L. & Pearson, N.
- Sevastjanova, I., Clements, B., Hall, R., Belousova, E.A., Griffin, W.L. & Pearson, N.
 291062 2011. Granitic magmatism, basement ages, and provenance indicators in the
 311063 Malay Peninsula: Insights from detrital zircon U-Pb and Hf-isotope data.
 Gondwana Research, 19 (4), 1024-1039.
- ³⁴₃₅1065 Sikumbang, N., 1986. Geology and tectonics of Pre-Tertiary rocks in the Meratus
 ³⁶₃₇1066 Mountains, South-East Kalimantan, Indonesia. PhD Thesis, University of
 ³⁸₃₉1067 London.
- 401068
41Sikumbang, N., 1990. The Geology and tectonics of the Meratus Mountains, South
Kalimantan, Indonesia. Geologi Indonesia. Journal of the Indonesian
Association of Geologists, 13(2), 1-31.
- ⁴⁵₄₆1071 Situmorang, B., 1982. The formation and evolution of the Makassar Basin,
 ⁴⁷₄₈1072 Indonesia. PhD Thesis, University of London.
- ⁴⁹1073 Sláma, J., Košler, J., Condon, D.J., Crowley, J.L., Gerdes, A., Hanchar, J.M., ⁵¹¹⁰⁷⁴ Horstwood, M.S.A., Morris, G.A., Nasdala, L., Norberg, N., Schaltegger, U., ⁵² Schoene, B., Tubrett, M.N., Whitehouse, M.J., 2008. Plešovice zircon — A ⁵⁴ new natural reference material for U–Pb and Hf isotopic microanalysis. ⁵⁶ Chemical Geology 249, 1-35.
- 58 59
- 59 60
- 61
- 62 63 64

- 1078 Smyth, H., Hall, R., Hamilton, J., Kinny, P., 2003. Volcanic origin of quartz-rich 1 2**1079** sediments in East Java. Indonesian Petroleum Association, Proceedings 29th ³₄1080 Annual Convention, Jakarta, 541-559.
- ⁵1081 Smyth, H.R., Hall, R., Nichols, G.J., 2008. Significant volcanic contribution to some 7**1082** 8 quartz-rich sandstones, east Java, Indonesia. Journal of Sedimentary গ083 Research 78, 335-356. 10
- 111084 Smyth, H.R., Hamilton, P.J., Hall, R., Kinny, P.D., 2007. The deep crust beneath ¹² 13**1085** island arcs: inherited zircons reveal a Gondwana continental fragment ¹⁴₁₅1086 beneath East Java, Indonesia. Earth and Planetary Science Letters 258, 269-¹⁶1087 282.
- 18**1088** 19 Southgate, P., Sircombe, K.N., Lewis, C.J., 2011. New insights into reservoir sand 20**1089** provenance in the Exmouth Plateau and Browse Basin. APPEA Perth, 21 22**1090** Western Australia.
- ²³₂₄1091 ²⁵₂₆1092 Spakman, W., Hall, R., 2010. Surface deformation and slab-mantle interaction during Banda arc subduction rollback. Nature Geoscience 3, 562-566.
- ²⁷1093 28 Sribudiyani, Muchsin, N., Ryacudu, R., Kunto, T., Astono, P., Prasetya, I., Yulianto, ²⁹1094 ³⁰ I., 2003. The collision of the East Java Microplate and its implication for 311095 hydrocarbon occurrences in the East Java Basin. Indonesian Petroleum 32 33**1096** Association, Proceedings 29th Annual Convention, IPA03-G-085 1-12.
- ³⁴ 35</sub>1097 Sukamto, R., 1975a. Geological map of Indonesia, Ujung Pandang 1:1,000,000. ³⁶1098 Geological Survey of Indonesia, Directorate of Mineral Resources, Geological ³⁸1099 Research and Development Centre, Bandung, Indonesia.
- 40**1100** Sukamto, R., 1975b. The structure of Sulawesi in the light of plate tectonics. Paper 41 42**1101** presented at the Regional Conference on the Geology and Mineral Resources ⁴³ 44**1102** of Southeast Asia, Jakarta, 4 - 7 August 1975.
- ⁴⁵₄₆1103 Sukamto, R., Setyogroho, B., Atmawinata, S., Aziz, S., Jamal, B., Suharsono, & ⁴⁷481104 Mangga, S. A., 1990. The Jurassic rocks in Indonesia. Bulletin Geological ⁴⁹1105 Research and Development Centre Bandung Indonesia, 14, 1-16.
- 51**1106** Sukamto, R., & Supriatna, S., 1982. The geology of the Ujung Pandang, Benteng 53**1107** and Sinjai quadrangles, Sulawesi 1:250, 000.
- ⁵⁴ 55**1108** Sukamto, R., & Westermann, G. E. G., 1993. Indonesia and Papua New Guinea. In ⁵⁶57**1109** G. E. G. Westermann (Ed.), The Jurassic of the Circum-Pacific (pp. 181-193): ⁵⁸1110 Cambridge University Press.
- 60 61 62

- 63 64
- 65

- 1111Tate, R. B., 1991. Cross-border correlation of geological formations in Sarawak and $_{2}^{1}$ 112Kalimantan. Bulletin of the Geological Society of Malaysia, 28, 63-96.
- ³₄1113 Tate, R. B., & Hon, V., 1991. The oldest rocks in Borneo; a note on the Tuang ⁵₆1114 Formation, West Sarawak and its importance in relation to the presence of a ⁷₁115 "basement" in West Borneo. Warta Geologi, Geological Society of Malaysia ⁹₁116 Newsletter, 17(5), 221-224.
- 11117van Bemmelen, R. W., 1949. The Geology of Indonesia. Nijhoff, The Hague:12
13Government Printing Office.
- van Leeuwen, T. M., Allen, C. M., Kadarusman, A., Elburg, M., Michael Palin, J., Muhardjo, & Suwijanto., 2007. Petrologic, isotopic, and radiometric age constraints on the origin and tectonic history of the Malino Metamorphic Complex, NW Sulawesi, Indonesia. Journal of Asian Earth Sciences, 29(5-6), 21123 751-777.
- Complex, NW Sulawesi, Indonesia. Journal of Asian Earth Sciences, 29(5-6), Complex, NW Sulawesi, Indonesia. Journal of Asian Earth Sciences, 29(5-6), 751-777. von der Borch, C.C., Grady, A.E., Hardjoprawiro, S., Prasetyo, H., Hadiwisastra, S., 1983. Mesozoic and Late Tertiary submarine fan sequences and their tectonic significance, Sumba, Indonesia. Sedimentary Geology 37, 113-132.
- von Rad, U. & Exon, N.F. 1982. Mesozoic-Cenozoic sedimentary and volcanic
 evolution of the starved passive continental margin off Northwest Australia. In:
 Studies in continental margin geology (Watkins, J.S. & Drake, C.L., Eds.)
 American Association of Petroleum Geologists Memoir, 34, 253-281.
- ³⁶₃₇131 Wakita, K., Munasari, S., J., Zulkarnain, I., & Miyazaki, K., 1994. Early Cretaceous tectonic events implied in the time-lag between age of radiolarian chert and its metamorphic basement in the Bantimala area, South Sulawesi, Indonesia.
 ⁴⁰133 Island Arc, 3, 90-102.
- Wakita, K., Munasri, & Widoyoko, B., 1994. Cretaceous radiolarians from the Luk Ulo Melange Complex in the Karangsambung area, Central Java, Indonesia.
 Journal of Southeast Asian Earth Sciences, 9(1/2), 29-43.
- ⁴⁹138 Wakita, K., Sopaheluwakan, J., Miyazaki, K., Zulkarnain, I. & Munasri 1996. Tectonic
 ⁵¹¹139 evolution of the Bantimala Complex, South Sulawesi, Indonesia. In: Hall, R. &
 ⁵²531140 Blundell, D.J. (Eds.), Tectonic Evolution of SE Asia. Geological Society of
 ⁵⁴551141 London Special Publication, 106, 353-364.
- ⁵⁶₅₇1142 Wakita, K., Miyazaki, K., Zulkarnain, I., Sopaheluwakan, J., & Sanyoto, P., 1998.
 ⁵⁸₅₉1143 Tectonic implications of new age data for the Meratus complex of south Kalimantan, Indonesia. Island Arc, 7(1-2), 202-222.
- 63 64

- 1145Wakita, K., 2000. Cretaceous accretionary-collision complexes in central Indonesia. $_{2}^{1}$ 146Journal of Asian Earth Sciences, 18(6), 739-749.
- ³₄1147 Wensink, H., 1994. Paleomagnetism of rocks from Sumba: tectonic implications ⁵₆1148 since the Late Cretaceous. Journal of Southeast Asian Earth Sciences 9, 51-⁷₈149 65.
- 9150Zimmermann, S., Hall, R., 2016. Provenance of Triassic and Jurassic sandstones in10111151111151the Banda Arc: Petrography, heavy minerals and zircon geochronology.12131152Gondwana Research 37, 1–19.

21153 **Figure Captions**

1

3

25

Fig. 1: Simplified map of SE Asia showing the Banda Arc Islands and blocks that rifted from Gondwana and were added by accretion to Sundaland (modified from Hall and Sevastjanova, 2012).

Fig. 2: The Banda Arc Islands discussed in this paper: Sumba, Timor, Babar and Tanimbar, showing key features of the stratigraphy of each island and the Mesozoic formations (Fm=Formation) that were analysed, with sample locations. Deep marine samples are highlighted in yellow and shallow marine samples in blue.

Fig. 3: Pre-breakup reconstruction in the Triassic showing the principal blocks rifted in the Jurassic with interpreted major sediment transport directions and suggested sources (from Zimmermann and Hall, 2016). Dark coloured areas are potential sources of zircon age populations.

Fig. 4: Reconstruction of Sundaland margin in the Cretaceous at c. 90 Ma after accretion of continental blocks rifted from the Banda embayment, based on Smyth et al. (2007), Hall et al. (2009), Hall (2012), and Hennig et al. (2016). The Banda block is present-day SW Borneo, the Argo block includes East Java and parts of West Sulawesi, and the Inner Banda block includes NW Sulawesi, and parts of NE Borneo.

³⁵/₃₆1170 Borneo.
³⁷/₃₈1171 Fig. 5: Sedimentary log of the well-bedded siltstone-sandstone intercalations at Konda Beach in south central Sumba; A) Interbedded sandstones and siltstones with up to 5 centimetres thick sandstone layers and internal fining upwards units; B)
⁴³/₄₁173 Small channel in siltstone sequence; C) Layer of a 10 centimetres thick medium to coarse-grained sandstone.

Fig. 6: A) Simplified geological map of the Tanimbar Islands, based on data collected in the field and structural interpretations modified from Kaye (1989); B) Panoramic view of the Arumit Member on Ungar Island; C) Well-bedded interbedded siltstones and mudstones; D) Contact of red shales of the Arumit Member with the Upper Sandstone Member.

Fig. 7: Sedimentary log of the Arumit Member of the Ungar Formation, showing 537181 details of the succession on Ungar Island and its interpreted relationship to the

46

62 63

1183 Lower and Upper Sandstone Members. Chert layers are highlighted in red and ${}_{2}^{1}$ 184 radiolarian ages are based on Jasin and Haile (1996).

⁴¹¹⁸⁵ Fig. 8: Summary of light mineral point counting of Cretaceous sandstones from the various islands. Ternary plots after Dickinson et al. (1983) showing possible provenance affiliation (Q – Quartz, F – Feldspar, L – Lithic fragments, Qm – Monocrystalline quartz, Lt – Total lithic fragments). Textures show assessments of sorting and rounding using simple number schemes which were used to estimate maturity.

Fig. 9: Summary plots of heavy mineral percentages, interpreted protoliths and varietal morphology of Cretaceous sandstones from the various islands.

19 20**1193** Fig. 10: Histograms showing grouped zircon ages for Cretaceous formations from ²¹₂₂1194 ²³₂₄1195 islands in the Banda Arc with possible sources. Bin width on the left (0-500 Ma) is 10 Ma, on the right (500-4000 Ma) is 50 Ma. Total numbers of zircons for each group ²⁵1196 26 are highlighted (red indicates the greater numbers, green the smaller). Percentages 27**1197** of Precambrian zircon grains that are older than 541 Ma are indicated; B) Bar charts 28 29**1198** for samples grouped according to age showing zircon morphology types for different ³⁰ 311199 ³² 331200 ³⁴ 35 age groups (Cr = Cretaceous, J = Jurassic, P/T = Permian-Triassic, C/C = Cambrian-Carboniferous, Neo = Neoproterozoic, Meso = Mesoproterozoic, Paleo = Paleoproterozoic, Arch = Archean).

Fig. 11: Kernel density estimation plots of Cretaceous formations from this study compared to Triassic and Jurassic formations from Zimmermann and Hall (2016). Colours indicate individual population ages (Cr=Cretaceous, J=Jurassic, P-T=Permian-Triassic, C-C=Ordovician-Carboniferous).

Fig. 12: Histograms summarizing the main features of zircon ages for the Schwaner Mountains of Borneo, Central Sulawesi and offshore Australia with colour bands marking likely sources. Bin width on the left (0-500 Ma) is 10 Ma, on the right (500-4000 Ma) is 50 Ma. Total numbers of zircons for each group are highlighted (red indicates the greater numbers, green the smaller). Percentages of Precambrian zircon grains that are older than 541 Ma are indicated.

Fig. 13: Map showing possible source areas in Australia and SE Asia with ages that resemble age populations found in the Banda Arc Islands and the NW Shelf of Australia (modified from Zimmermann and Hall, 2016).

65

55

1215 Fig. 14: A) Summary of zircon ages in Cretaceous sandstones of SE Asia and ¹ 21216 equivalents in the Australian NW Shelf (below) and possible source areas in ³₄1217 Australia and SE Asia (above). B) Cretaceous palaeogeographic reconstruction with ⁵1218 tectonic elements and age provinces with major interpreted sediment transport ⁷1219 8 directions (arrows) showing suggested sources for Cretaceous formations in the 91220 Banda Arc that are now the Banda Allochthon (BA). BA-WT=Cretaceous West 10 11**1221** Timor, BA-ET=Cretaceous East Timor, BA-B=Cretaceous Babar, BA-T=Cretaceous ¹² 13**1222** Tanimbar. The provenance features suggest distant syn-sedimentary Cretaceous ¹⁴1223 sources in the Schwaner Mountains and Sulawesi, with additional proximal sources ¹⁶1224 in Sumba and other parts of the outer Sundaland margin, with recycling of Permian-¹⁸1225 Triassic and Jurassic pre-rift units (purple and blue).

Fig. 15: A) Tectonic reconstructions for SE Asia (based on Hall, 2012) with focus on the rifting Argo and Banda terranes from the Late Jurassic to Early Cretaceous; B) Tectonic reconstructions of the Banda Allochthon (BA) at 23 Ma showing the initial collision between the Sula Spur and Sundaland; at 15 Ma showing subduction hinge rollback and early stage of fragmentation of the Sula Spur; at 5 Ma showing location of allochthonous fragments before Timor arc-continent collision; and present-day configuration highlighting the outer Banda Arc Islands with overthrust fragments.



Volcanic arc accreted in the Cretaceous

Fig. 1: Simplified map of SE Asia showing the Banda Arc Islands and blocks that rifted from Gondwana and have been added by accretion to Sundaland (modified from Hall and Sevastjanova, 2012).



Fig. 2: The Banda Arc Islands discussed in this paper: Sumba, Timor, Babar and Tanimbar, showing key features of the stratigraphy of each island and the Mesozoic formations (Fm=Formation) that were analysed, with sample locations. Deep marine samples are highlighted in yellow and shallow marine samples in blue.



Fig. 3: Pre-breakup reconstruction in the Triassic showing the principal blocks rifted in the Jurassic with interpreted major sediment transport directions and suggested sources (from Zimmermann and Hall, 2016). Dark coloured areas are potential sources of zircon age populations.



Fig. 4: Reconstruction of Sundaland margin in the Cretaceous at c. 90 Ma after accretion of continental blocks rifted from the Banda embayment, based on Smyth et al. (2007), Hall et al. (2009), Hall (2012), and Hennig et al. (2016). The Banda block is present-day SW Borneo, the Argo block includes East Java and parts of West Sulawesi, and the Inner Banda block includes NW Sulawesi, and parts of NE Borneo.



Fig. 5: Sedimentary log of the well-bedded siltstone-sandstone intercalations at Konda Beach in south central Sumba; A) Interbedded sandstones and siltstones with up to 5 centimetres thick sandstone layers and internal fining upwards units; B) Small channel in siltstone sequence; C) Layer of a 10 centimetres thick medium to coarse-grained sandstone.



Fig. 6: A) Simplified geological map of the Tanimbar Islands, based on data collected in the field and structural interpretations modified from Kaye (1989); B) Panoramic view of the Arumit Member on Ungar Island; C) Well-bedded interbedded siltstones and mudstones; D) Contact of red shales of the Arumit Member with the Upper Sandstone Member.



Fig. 7: Sedimentary log of the Arumit Member of the Ungar Formation, showing details of the succession on Ungar Island and its interpreted relationship to the Lower and Upper Sandstone Members. Chert layers are highlighted in red and radiolarian ages are based on Jasin and Haile (1996).



Fig. 8: Summary of light mineral point counting of Cretaceous sandstones from the various islands. Ternary plots after Dickinson et al. (1983) showing possible provenance affiliation (Q – Quartz, F – Feldspar, L – Lithic fragments, Qm – Monocrystalline quartz, Lt – Total lithic fragments). Textures show assessments of sorting and rounding using simple number schemes which were used to estimate maturity.



Fig. 9: Summary plots of heavy mineral percentages, interpreted protoliths and varietal morphology of Cretaceous sandstones from the various islands.

Figure 10



Fig. 10: Histograms showing grouped zircon ages for Cretaceous formations from islands in the Banda Arc with possible sources. Bin width on the left (0-500 Ma) is 10 Ma, on the right (500-4000 Ma) is 50 Ma. Total numbers of zircons for each group are highlighted (red indicates the greater numbers, green the smaller). Percentages of Precambrian zircon grains that are older than 541 Ma are indicated; B) Bar charts for samples grouped according to age showing zircon morphology types for different age groups (Cr = Cretaceous, J = Jurassic, P/T = Permian-Triassic, C/C = Cambrian–Carboniferous, Neo = Neoproterozoic, Meso = Mesoproterozoic, Paleo = Paleoproterozoic, Arch = Archean).



Fig. 11: Kernel density estimation plots of Cretaceous formations from this study compared to Triassic and Jurassic formations from Zimmermann and Hall (2016). Colours indicate individual population ages (Cr=Cretaceous, J=Jurassic, P-T=Permian-Triassic, C-C=Ordovician-Carboniferous).



Fig. 12: Histograms summarizing the main features of zircon ages for the Schwaner Mountains of Borneo, Central Sulawesi and offshore Australia with colour bands marking likely sources. Bin width on the left (0-500 Ma) is 10 Ma, on the right (500-4000 Ma) is 50 Ma. Total numbers of zircons for each group are highlighted (red indicates the greater numbers, green the smaller). Percentages of Precambrian zircon grains that are older than 541 Ma are indicated.



Fig. 13: Map showing possible source areas in Australia and SE Asia with ages that resemble age populations found in the Banda Arc Islands and the NW Shelf of Australia (modified from Zimmermann and Hall, 2016).





Fig. 14: A) Summary of zircon ages in Cretaceous sandstones of SE Asia and equivalents in the Australian NW Shelf (below) and possible source areas in Australia and SE Asia (above). B) Cretaceous palaeogeographic reconstruction with tectonic elements and age provinces with major interpreted sediment transport directions (arrows) showing suggested sources for Cretaceous formations in the Banda Arc that are now the Banda Allochthon (BA). BA-WT=Cretaceous West Timor, BA-ET=Cretaceous East Timor, BA-B=Cretaceous Babar, BA-T=Cretaceous Tanimbar. The provenance features suggest distant syn-sedimentary Cretaceous sources in the Schwaner Mountains and Sulawesi, with additional proximal sources in Sumba and other parts of the outer Sundaland margin, with recycling of Permian-Triassic and Jurassic pre-rift units (purple and blue).



Fig. 15: A) Tectonic reconstructions for SE Asia (based on Hall, 2012) with focus on the rifting Argo and Banda terranes from the Late Jurassic to Early Cretaceous; B) Tectonic reconstructions of the Banda Allochthon (BA) at 23 Ma showing the initial collision between the Sula Spur and Sundaland; at 15 Ma showing subduction hinge rollback and early stage of fragmentation of the Sula Spur; at 5 Ma showing location of allochthonous fragments before Timor arc-continent collision; and present-day configuration highlighting the outer Banda Arc Islands with overthrust fragments.