

A CLASSIFICATION OF ALL 1-SALEM GRAPHS

LEE GUMBRELL AND JAMES MCKEE

ABSTRACT. One way to study certain classes of polynomials is by considering examples that are attached to combinatorial objects. Any graph G has an associated reciprocal polynomial R_G , and with two particular classes of reciprocal polynomials in mind one can ask the questions: (a) when is R_G a product of cyclotomic polynomials (giving the *cyclotomic graphs*)? (b) when does R_G have the minimal polynomial of a Salem number as its only non-cyclotomic factor (the non-trivial *Salem graphs*)? Cyclotomic graphs were classified by Smith in 1970; the maximal connected ones are known as Smith graphs. Salem graphs are ‘spectrally close’ to being cyclotomic, in that nearly all their eigenvalues are in the critical interval $[-2, 2]$. On the other hand Salem graphs do not need to be ‘combinatorially close’ to being cyclotomic: the largest cyclotomic induced subgraph might be comparatively tiny.

We define an m -Salem graph to be a connected Salem graph G for which m is minimal such that there exists an induced cyclotomic subgraph of G that has m fewer vertices than G . The 1-Salem subgraphs are both spectrally close and combinatorially close to being cyclotomic. Moreover, every Salem graph contains a 1-Salem graph as an induced subgraph, so these 1-Salem graphs provide some necessary substructure of all Salem graphs. The main result of this paper is a complete combinatorial description of all 1-Salem graphs: in the non-bipartite case there are 25 infinite families and 383 sporadic examples.

1. INTRODUCTION

Definition 1. *A graph is called cyclotomic if all its eigenvalues lie in the interval $[-2, 2]$.*

A bipartite graph G on n vertices is called a Salem graph if the largest eigenvalue λ_1 is greater than 2, and the remaining $n - 1$ eigenvalues are no greater than 2.

A non-bipartite graph G on n vertices is called a Salem graph if the largest eigenvalue λ_1 is greater than 2 and the remaining $n - 1$ eigenvalues are in the interval $[-2, 2]$.

Some motivation, and the reason for this nomenclature, will be explained in the next section. For the moment we move rapidly to the statement of the theorem to be proved in this paper.

A number of partial results concerning the structure of Salem graphs appear in [28, Section 3], and a complete description of all Salem trees in [28, Theorem 7.2], but a combinatorial description of all Salem graphs is still unknown. In this paper we increase our understanding of the set of Salem graphs by considering how close they are to being cyclotomic in the following combinatorial sense: how many vertices do we need to delete to produce an induced cyclotomic subgraph?

1991 *Mathematics Subject Classification.* 11R06, 05C50.

Key words and phrases. Salem graphs, graph spectra.

Definition 2. A Salem graph G is called an m -Salem graph if m is minimal such that there exists a set of m vertices in the graph whose deletion leaves a graph with all eigenvalues in the interval $[-2, 2]$.

The 1-Salem graphs are both spectrally and combinatorially close to being cyclotomic: there is a single eigenvalue greater than 2, and there exists some vertex whose deletion leaves a cyclotomic induced subgraph. Our main theorem is a complete classification of this especially interesting subset of the Salem graphs. We remark that any Salem graph contains a 1-Salem graph as an induced subgraph, so that our classification reveals something of the necessary substructure of an arbitrary Salem graph. It is worth observing that our definition requires an m -Salem graph to be connected, but that when considering induced cyclotomic subgraphs these subgraphs need not be connected.

We also remark that certainly not all Salem graphs are 1-Salem. For example, the complete graph on n vertices is a Salem graph for $n \geq 4$, but it is not 1-Salem if $n \geq 5$.

The description of bipartite 1-Salem graphs is rather trivial, and is essentially contained in [28]. For non-bipartite Salem graphs all eigenvalues are at least -2 , and Cameron *et al* ([3]) showed that the connected graphs with this property are either generalised line graphs or belong to a finite list of graphs represented in the root system E_8 . The latter are disposed of by a finite computation. The main work in our classification comes from a description of the 1-Salem graphs that are generalised line graphs. We summarise our results in the following Theorem.

Theorem 1. (i) All bipartite 1-Salem graphs are described by Theorem 5 (§4).
(ii) Every 1-Salem graph that is a generalised line graph is described by Theorem 6; there are 25 infinite families and 6 sporadic examples (§5).
(iii) There are 377 non-bipartite 1-Salem graphs that are not generalised line graphs: see Theorem 11 (§6).

In the next section we give some motivation, history, and discussion of related results. Then in Section 3 we state some definitions and results that we will need to prove the three components of Theorem 1. The proof of Theorem 1 is split over three sections: Section 4 for the bipartite graphs; Section 5 for the generalised line graphs; and Section 6 for the non-bipartite graphs that are not generalised line graphs.

2. MOTIVATION

2.1. The remarkable interval $[-2, 2]$. An algebraic integer θ is called *totally real* if all its Galois conjugates are real. One can then define the real interval $I_\theta = [\theta_{min}, \theta_{max}]$, where θ_{min} and θ_{max} are the smallest and largest of the Galois conjugates of θ .

The number 4 plays a peculiar role in the theory of totally real algebraic integers. If J is any interval of length strictly less than 4, then there exist only finitely many totally real θ such that $I_\theta \subseteq J$ (proved by Schur for J centred on the origin, and extended to the general case by Pólya in a footnote to Schur's paper [38]). On the other hand, if J is any interval of length strictly greater than 4, then there exist infinitely many totally real θ such that $I_\theta \subseteq J$ (Robinson, [37]). If J has length exactly 4 and the endpoints are integers, then there are infinitely many such θ . If $J = [a, a + 4]$ with a an integer, and $I_\theta \subseteq J$, then $I_{\theta-a-2} \subseteq [-2, 2]$, so

that essentially the only such J of interest is the special interval $[-2, 2]$. For other intervals of length 4 nothing is known.

By a theorem of Kronecker [20], the only θ for which $I_\theta \subseteq [-2, 2]$ are those of the form $\theta = \zeta + 1/\zeta$ for ζ some (complex) root of 1. For this reason one sometimes refers to this as the *cyclotomic* case.

2.2. Algebraic integers from combinatorial objects. One approach to the study of algebraic integers is to look for examples attached to combinatorial objects (graphs, knots, etc.). The eigenvalues of graphs are totally real, and provide a fruitful hunting-ground for such algebraic integers. Indeed Estes [12] showed that all totally real algebraic integers arise as eigenvalues of graphs (Hoffman [17, 18] having shown that the problem was equivalent to that for eigenvalues of integer symmetric matrices). On the other hand it is easy to find examples of totally real θ for which the minimal polynomial of θ is not the characteristic polynomial of a graph, and with more work one can find examples where the minimal polynomial of θ is not even the *minimal* polynomial of any integer symmetric matrix ([11, 26]).

If G is a graph, we let χ_G denote its characteristic polynomial, and define R_G to be the associated reciprocal polynomial $R_G(z) := z^{\deg \chi_G} \chi_G(z + 1/z)$. For any set S of reciprocal polynomials of interest, there is a correspondingly interesting set of graphs G for which $R_G \in S$.

2.3. Cyclotomic graphs. After §2.1, the question of which graphs have all eigenvalues in the interval $[-2, 2]$ acquires peculiar interest, and such graphs we call *cyclotomic*. (The reason for this name is that after Kronecker's theorem the associated reciprocal polynomial of such a graph is a product of cyclotomic polynomials.) It is enough to consider connected examples, and Smith [39] gave a complete classification. The maximal connected examples comprise two infinite families and three sporadic cases: see Figure 1.




FIGURE 1. The maximal connected cyclotomic graphs (Smith graphs). Note that each graph has one more vertex than the subscript in its name.

2.4. Salem numbers and Salem graphs: definitions. A *Salem number* is a real algebraic integer τ such that: (i) $\tau > 1$; (ii) all Galois conjugates of τ other than τ itself lie in the unit disc $\{z \in \mathbb{C} : |z| \leq 1\}$; (iii) at least of the Galois conjugates of τ has modulus exactly 1. It follows that the minimal polynomial of τ is reciprocal, τ has even degree at least 4, and all of the Galois conjugates of τ other than τ and $1/\tau$ lie on the unit circle $\{z \in \mathbb{C} : |z| = 1\}$.

It is not known whether there are Salem numbers arbitrarily close to 1. The smallest known Salem number is $1.17628\dots$, the larger real root of the reciprocal polynomial $L(z) = z^{10} + z^9 - z^7 - z^6 - z^5 - z^4 - z^3 + z + 1$. This polynomial was discovered by considering examples attached to combinatorial objects: $L(-z)$ is the Alexander polynomial of a pretzel knot.

One can ask when a graph has as its reciprocal polynomial the minimal polynomial of a Salem number, and one might call these ‘Salem graphs’. Since irreducibility of the reciprocal polynomial is not easy to capture combinatorially, it is natural to relax this definition slightly, and merely ask that the reciprocal polynomial has all but two roots on the unit circle. Such graphs cannot be bipartite: the eigenvalues of a bipartite graph are symmetric about 0, and the reciprocal polynomial as defined above cannot then have a unique root outside the unit disc. The closest that a non-cyclotomic bipartite graph can get to being cyclotomic (measured in terms of the number of eigenvalues outside the critical interval $[-2, 2]$) is to have a single root greater than 2 and hence also a root less than -2 . In such cases we can exploit the symmetry of the eigenvalues to recover a polynomial whose roots are in the right place to be the minimal polynomial of a Salem number (but with no guarantee of irreducibility): we redefine $R_G(z) = z^{\deg(\chi_G)/2} \chi_G(\sqrt{z} + 1/\sqrt{z})$ for bipartite graphs.

This discussion motivates the following extension of Definition 1, which might otherwise seem unnecessarily convoluted. We follow a common convention that if G has n vertices then its eigenvalues are labelled $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$.

Definition 3. *A bipartite graph G is called a Salem graph if the largest eigenvalue λ_1 is greater than 2, and the remaining $n - 1$ eigenvalues are no greater than 2. Then $\lambda_n = -\lambda_1 < -2$. A bipartite Salem graph is called trivial if $\lambda_1^2 \in \mathbb{Z}$. The associated number $\tau(G)$ is the larger root of $\sqrt{z} + 1/\sqrt{z} = \lambda_1$; this is a Salem number unless G is trivial.*

A non-bipartite graph G is called a Salem graph if $\lambda_1 > 2$ and the remaining $n - 1$ eigenvalues are in the interval $[-2, 2]$. A non-bipartite Salem graph is called trivial if $\lambda_1 \in \mathbb{Z}$. The associated number $\tau(G)$ is the larger root of $z + 1/z = \lambda_1$; this is a Salem number unless G is trivial.

This notion of a Salem graph was introduced in [28]. Special cases of Salem graphs (or associated Coxeter transformations) and their connection with Salem numbers were studied in earlier (and later) papers: [10], [32], [25], [21], [24], [22], [31], [27], [23]. They are also the subject of the first author’s thesis ([15]). The highlight of [28] was a proof that all limit points of the set of graph Salem numbers are Pisot numbers, confirming a conjecture of Boyd [2] in this setting. Limit points of special subsets of the set of graph Salem numbers are also considered in [25], [21], [24], [27] and [40]. Many examples of Salem graphs and their spectra have been considered in other contexts, for example [41], [7], [30].

Graphs with $\lambda_2 \leq 2$ are known as *reflexive* graphs. See, for example, [36] and references therein. Reflexive graphs with a cutvertex are neatly described in [34]. All Salem graphs are certainly reflexive, and bipartite reflexive graphs are either Salem or cyclotomic. Not all Salem graphs have a cutvertex; for example the complete bipartite graph $K_{3,2}$ does not. The main challenge in classifying non-bipartite Salem graphs is to control the smallest eigenvalue, as well as the largest two.

3. BACKGROUND

For convenience we list some standard definitions and results that we shall exploit in the proof of Theorem 1. We refer to the first result as ‘interlacing’. It allows us to bound the second-largest eigenvalue of a graph by the index of any induced subgraph obtained by deleting a single vertex.

Theorem 2 (Cauchy [5]; or see [14], Theorem 9.1.1). *Let G be an n -vertex graph with vertex set $V(G)$ and eigenvalues $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$. Also, let H be the induced graph on $V(G) \setminus \{v\}$ obtained from G by deleting the vertex v and its incident edges. Then the eigenvalues $\mu_1 \geq \mu_2 \geq \dots \geq \mu_{n-1}$ of H interlace with those of G ; that is*

$$\lambda_1 \geq \mu_1 \geq \lambda_2 \geq \mu_2 \geq \dots \geq \mu_{n-1} \geq \lambda_n.$$

We also need the following definitions from graph theory: see [9] or [14]. Let G be an n -vertex graph or multigraph. The *line graph* $L(G)$ is the graph whose vertices are the edges of G , with two vertices in $L(G)$ adjacent whenever the corresponding edges in G have exactly one vertex in common. The *root graph* of a line (multi)graph $L(G)$ is simply G itself (with a small number of exceptions, G is uniquely determined by $L(G)$). Now let a_1, \dots, a_n be non-negative integers. The *generalized line graph* $L(G; a_1, \dots, a_n)$ is the graph $L(\hat{G})$, where \hat{G} is the multigraph $G(a_1, \dots, a_n)$ obtained from G by adding a_i pendant 2-cycles at vertex v_i ($i = 1, \dots, n$). An *internal path* of a graph G is a sequence of vertices v_1, \dots, v_k of G such that all vertices are distinct (except possibly v_1 and v_k), v_i is adjacent to v_{i+1} (for $i = 1, \dots, k-1$), v_1 and v_k have degree at least 3, and all of v_2, \dots, v_{k-1} have degree 2.

We define a *generalized cocktail party graph* (*GCP*) as a graph isomorphic to a complete graph (or clique) but with some independent edges removed; that is a graph where all vertices have degree $n-1$ or $n-2$. We will use $GCP(n, m)$ to denote the GCP on n vertices with m edges removed, where $0 \leq m \leq \lfloor n/2 \rfloor$. In $GCP(n, m)$, we will refer to a vertex of degree $n-1$ as being ‘of maximal degree’.

Theorem 3 (see [9], Theorem 2.3.1 and Theorem 2.1.1). *A connected graph is a generalized line graph if and only if its edges can be partitioned into GCPs such that*

- (i) *two GCPs have at most one common vertex;*
- (ii) *each vertex is no more than two GCPs;*
- (iii) *if two GCPs have a common vertex, then it is of maximal degree in both of them.*

Also, a graph is a line graph if and only if its edges can be partitioned in such a way that every edge is in a clique and no vertex is in more than two cliques.

A further important tool in the proof of Theorem 6 is the following:

Proposition 4 ([28], Proposition 3.2). *Let G be a connected graph with $\lambda_1 > 2$ and $\lambda_2 \leq 2$, then the vertices V of G can be partitioned as $V = M \cup A \cup H$ where*

- (i) *the induced subgraph $G|_M$ is minimal subject to $\lambda_1(G|_M) > 2$;*
- (ii) *the set A consists of all vertices of $G|_{A \cup H}$ adjacent to some vertex of M ;*
- (iii) *the induced subgraph $G|_H$ is cyclotomic.*

4. ALL BIPARTITE 1-SALEM GRAPHS

We now begin our proof of Theorem 1, starting with part (i); the bipartite 1-Salem graphs.

Theorem 5. *Let H_1, \dots, H_s be a finite set of connected cyclotomic bipartite graphs (so excluding odd cycles from the connected induced subgraphs of the graphs in Figure 1), and for each i ($1 \leq i \leq s$) let S_i be a non-empty subset of the vertices of H_i such that all the vertices of S_i fall in the same subset of the bipartition of H_i . Form the graph G by taking the union of all the H_i along with a new vertex v , and with edges joining v to each vertex in each S_i . Then*

- (i) *unless G is cyclotomic, it is 1-Salem;*
- (ii) *all connected bipartite 1-Salem graphs arise in this way.*

Proof. The second part is clear: if G is a bipartite 1-Salem graph, then there exists a vertex v whose deletion leaves a cyclotomic induced subgraph, and the components of this give the H_i . The first part is a consequence of interlacing: G has at most one eigenvalue greater than 2, and being bipartite we are done. \square

One can be more explicit and describe precisely which combinations of H_i and S_i result in G being cyclotomic, simply by running through the possibilities for cyclotomic G given by the connected induced subgraphs of the graphs in Figure 1.

5. ALL 1-SALEM GENERALISED LINE GRAPHS

Theorem 6. *The 1-Salem generalised line graphs are precisely the graphs in Figures 2, 3 and 4.*

A quick notational point: in Figures 2–4, a dashed edge indicates a path between the endpoints of the dashed edge, having an arbitrary number of edges (perhaps even none, or perhaps with a lower bound shown); the parameter attached to a dashed edge gives the number of edges on this path. Dotted edges and vertices are used to indicate edges and vertices that may or may not be there. Here the dotted edges and vertices always form a ‘snake’s tongue’ shape. The 1-Salem generalised line graphs of Theorem 6 are presented as 25 infinite families and six sporadic graphs, but there are in fact 60 non-isomorphic infinite families when we consider all the possible options of paths with snake’s tongues attached.

Also, as a simple corollary to Theorem 6 we can easily note which of the graphs in Figures 2–4 are line graphs (rather than generalised line graphs) based on the characterization in Theorem 3. These are the graphs where the edges can be partitioned into cliques rather than GCPs and there are 12 infinite families and 3 sporadic graphs.

The following trivial extension of [28, Theorem 3.4] can be used to show that all the graphs in Figures 2–4 are Salem graphs.

Lemma 7. *Suppose that G is a non-cyclotomic non-bipartite graph containing a vertex v such that the induced subgraph on $V(G) \setminus \{v\}$ is cyclotomic. Also suppose that G is in the family of graphs with least eigenvalue greater than -2 , then G is a Salem graph.*

To apply Lemma 7 to the graphs in Figures 2–4, we note that all the graphs in that figure are generalised line graphs (using the characterisation in Theorem 3, for example), and in each case one readily spots a vertex v whose deletion leaves a cyclotomic induced subgraph.

We now prove Theorem 6 using the structure given in Proposition 4; that is we will grow our graphs starting with the vertices in M , then $M \cup A$, then $M \cup A \cup H$. We quickly note a simple lemma that will be referred to frequently in the proof.




FIGURE 2. Twelve of the 25 infinite families of 1-Salem generalised line graphs. The parameters a, b, c, d are ≥ 0 unless specified. For G_1 , recall that $K_{1,4}$ is the star on five vertices, and e here is a single edge between any pair of the four vertices of degree 1.

Lemma 8. *A 1-Salem graph G may not contain an induced K_5 . Also, if it contains an induced K_4 , then only one of the four vertices in that K_4 may be attached to any other vertices in G and the vertex we remove to make G cyclotomic must be this distinguished vertex in K_4 .*

Proof. The first sentence is clear since if we remove any one of the vertices of K_5 we obtain a K_4 which is not cyclotomic. For the second sentence, removing a vertex v of K_4 leaves a K_3 so if any of the other vertices of the K_4 were attached to any vertices of G , so would the K_3 be after removing v , and no connected supergraph of K_3 is cyclotomic. \square

5.1. M — **the minimal graphs.** We begin by considering what our minimal graphs may look like.

Proposition 9. *All Salem generalized line graphs must contain a K_3 .*

Proof. We know from Theorem 3 that generalised line graphs are built from GCPs and it is easy to see that the only GCPs that do not contain a K_3 are $GCP(1, 0)$,




FIGURE 3. Thirteen of the 25 infinite families of 1-Salem generalised line graphs. The parameters a, b are ≥ 0 unless specified.




FIGURE 4. The six sporadic 1-Salem generalised line graphs.

$GCP(2,0)$, $GCP(2,1)$, $GCP(3,1)$ and $GCP(4,2)$. These graphs are certainly all cyclotomic and, moreover, all the generalised line graphs that can be made using them (observing the rules in Theorem 3) will be subgraphs of either \hat{A}_n or \hat{D}_n . However, by definition Salem graphs are non-cyclotomic, so a Salem generalised line graph must include at least one GCP that contains a K_3 . \square

As a corollary to this we get that the minimal graphs we are after must be the three ways of attaching a single vertex to a K_3 .

Corollary 10. *The minimal graphs with respect to the property of being a Salem generalized line graph are the three graphs in Figure 5.*




FIGURE 5. The three minimal graphs in Corollary 10.

We now know what our minimal graphs look like but before proceeding we make one more observation. For $G \setminus \{v\}$ to be cyclotomic we must have $v \in M$; that is, the vertex we are removing to induce a cyclotomic graph must be one of the vertices of the minimal graph M .

5.2. A — the adjacent vertices. Growing our graph from $G|_M$ to $G|_{M \cup A}$ is the most difficult part, however the restriction to 1-Salem generalized line graphs reduces this to a finite search, after some work. By Theorem 3 we have a highly-constrained structure: we cannot simply add vertices and edges anywhere. First let us consider the ways we can partition M_1 , M_2 and M_3 into GCPs; this will reveal where we can add vertices. The generalised line graph M_1 can be seen uniquely as a K_3 and a K_2 ; it then has three vertices of maximal degree that are not already in two GCPs. And M_3 can be seen uniquely as a K_4 , but by Lemma 8 we know that we can attach further vertices to only one of its original vertices. The graph M_2 , however, is one of the seven graphs that has two non-isomorphic root multigraphs (see [9, Theorem 2.3.4]) so we need to consider both versions. Let $M_{2,1}$ be M_2 partitioned as a K_3 with two K_2 's attached and both joined at their other ends; here we only have one vertex that is of maximal degree and not already in two GCPs. Let $M_{2,2}$ be M_2 partitioned as a $GCP(4, 1)$ where we then have two vertices of maximal degree.

The set of vertices A are those in the graph $G|_{A \cup H}$ that are adjacent to a vertex in M . By Theorem 3, to grow from $G|_M$ to $G|_{M \cup A}$ we can:

- expand a GCP to a larger one that contains the original one (taking care of the degrees of vertices);
- attach a new GCP to a vertex of maximal degree that is only in one GCP (attaching only at vertices of maximal degree);
- do both of the above.

Once we have done this we can then also add an edge between any two vertices of A of maximal degree in their GCPs (in effect, adding a K_2) or take two maximal degree vertices of A that are in separate GCPs and only in one GCP each and ‘merge’ them together (connect two disconnected GCPs by making them share an available maximal degree vertex).

Lemma 8 helps here as we know that we need not consider any GCPs that contain a K_5 . In fact, by studying the GCPs G that have this property and looking at the induced graph $G \setminus \{v\}$ for each $v \in V(G)$ we find the fairly short list of GCPs that we can attach or expand to in Table 1 below.

$GCP(2,0) = K_2$		2 vertices of maximal degree
$GCP(3,1)$		1 vertex of maximal degree
$GCP(3,0) = K_3$		3 vertices of maximal degree
$GCP(4,1)$		2 vertices of maximal degree
$GCP(4,0) = K_4$		4 vertices of maximal degree
$GCP(5,2)$		1 vertex of maximal degree

TABLE 1. The GCPs that we can attach or expand to in A .

Note that $GCP(4,2) = C_4$ is not included; when partitioned as $GCP(4,2)$ it has no vertices of maximal degree, so cannot be attached to anything and when partitioned as four K_2 's it has no vertices that are not already in two GCPs. Also, $GCP(3,1)$ and $GCP(5,2)$ are two of the seven graphs that have non-isomorphic root multi-graphs. However, when $GCP(5,2)$ is partitioned as two K_3 's and two K_2 's it has no vertices that are not already in two GCPs and when $GCP(3,1)$ is partitioned as two K_2 's one of its vertices will not be in A .

The process of going from $G|_M$ to $G|_{M \cup A}$ is then a finite one; we only have so many GCPs in the minimal graphs $M_1, M_{2,1}, M_{2,2}$ and M_3 to expand and only so many ways to attach these six GCPs to them. We also have a small number of cases where we can add in extra edges between vertices of A or merge them. In working through all these combinations we discard a number of graphs that are not 1-Salem as they require more than one vertex to be removed to make them cyclotomic.

The list of 1-Salem generalised line graphs $G|_{M \cup A}$ has been omitted for reasons of space. There are 224 that are distinct as graphs, although many of these graphs can arise in more than one way as $G|_{M \cup A}$. The largest has 11 vertices. Initially these were all found by hand; the list was then checked by a computer search.

5.3. H — the cyclotomic parts. We now look at the set H in Lemma 4. We can reduce our choices by observing that the only cyclotomic graphs that are also generalised line graphs are \tilde{D}_n and $\tilde{A}_n = C_{n+1}$. However, we will show that $G|_H$ cannot contain cycles and can only contain subgraphs of \tilde{D}_n . For $n > 3$ we note that \tilde{A}_n can be partitioned uniquely as $(n+1)$ K_2 's with each vertex in two GCPs, so we cannot simply attach vertices to it. The other option is then to expand a GCP to a larger one so that we now have vertices only in one GCP to attach to other things. The smallest case is to expand one of the $(n+1)$ K_2 's to a K_3 . Clearly, this new vertex must be in A rather than H as the graph is no longer cyclotomic. However, in order to make the graph cyclotomic we must remove a vertex but all of the vertices are in A or H and we have shown that the vertex we remove must be in M . Cycles of length 3 can be seen as both three K_2 's or one K_3 but by similar reasoning on the choice of vertex we are removing, we can exclude this case too. A similar argument holds again for cycles of length 4 for both ways of partitioning its edges.

The graph \tilde{D}_n can be uniquely partitioned as $(n - 4)$ K_2 's with a $GCP(2, 0)$ (or snake's tongue) at either end. However, in this graph each vertex of maximal degree within its GCP is already in two GCPs. If we remove one or both of the snake's tongues we are left with graphs we can work with—a path or a path with a snake's tongue on the end—each with at least one vertex of maximal degree that is only in one GCP. With this in mind we can think of $G|_H$ simply consisting of paths of any length, possibly with a snake's tongue on one end.

The final step in growing these graphs is to go from $G|_{MUA}$ to $G|_{MUAUH} = G$. To any vertices in A of maximal degree and only in one GCP we can attach a single path of arbitrary length (remembering that Lemma 8 tells us that we can only attach things to one vertex of a K_4). If we so choose, we can join any two of these pendant paths together (equivalently, attach a path of arbitrary length to two different elements of A of maximal degree that are each only in one GCP). Furthermore, on the end of any pendant paths we can include a snake's tongue. These graphs are then precisely the graphs in Figures 2–4 and Lemma 7 tells us that they are all 1-Salem, completing this section of the proof.

We can consider letting the lengths of one or more pendant or internal paths tend to infinity. By [28, Corollary 4.4] the corresponding sequence of graph Salem numbers converges to a Pisot number. By studying the different ways we can let paths tend to infinity, we can produce several explicit families of Pisot numbers.

6. THE 1-SALEM GRAPHS REPRESENTED IN E_8

6.1. The root system and signed graph E_8 . There are various concrete descriptions of the root system E_8 (see [4, Chapter 3], for example). Here is one of them. Let $\mathbf{e}_1, \dots, \mathbf{e}_8$ be an orthogonal basis for \mathbb{R}^8 , with each \mathbf{e}_i having length $\sqrt{2}$. The root system E_8 contains 240 vectors: the 16 vectors $\pm\mathbf{e}_1, \dots, \pm\mathbf{e}_8$, and 224 vectors of the form $(\pm\mathbf{e}_i \pm \mathbf{e}_j \pm \mathbf{e}_k \pm \mathbf{e}_l)/2$, where $ijkl$ is one of the 14 strings 1234, 1256, 1278, 1357, 1368, 1467, 2358, 2367, 2457, 2468, 3456, 3478, 5678. The usual inner product of any two vectors from these 240 is one of $-2, -1, 0, 1$, or 2 ; and all the vectors have length 2. A graph G with adjacency matrix A is said to be represented in E_8 if $A + 2I$ is the Gram matrix of some subset of the vectors forming the root system E_8 (so vectors correspond to vertices, and adjacency of distinct vertices is given by the usual inner product of the corresponding vectors).

The *Weyl group* $W(E_8)$ is the group of isometries of E_8 generated by the involutions $\mathbf{v} \mapsto \mathbf{v} - (\mathbf{v} \cdot \mathbf{v}_i)\mathbf{v}_i$ for each of the 240 vectors \mathbf{v}_i (in fact there are only 120 of these involutions, since \mathbf{v}_i and $-\mathbf{v}_i$ yield the same involution).

6.2. Fast backtracking in E_8 . Let P be a property of graphs such that if G has property P then so does any induced subgraph. (Such a property is called *hereditary*.) The following natural process, familiar to computational graph-theorists, provides a computationally-efficient method for searching for all graphs that: (a) are represented in E_8 ; and (b) have property P .

Assign an ordering to the vectors of E_8 : $\mathbf{v}_1, \dots, \mathbf{v}_{240}$. A naive search for all graphs having property P that are represented in E_8 would proceed as follows. Start with an empty list of vectors; look through the vectors in order, adding them to the list so long as the vectors in the list represent a graph that has property P ; once stuck, backtrack. The number of graphs represented in E_8 is so large that this process is only practical for extremely restrictive properties P . To improve matters, we exploit the isometries of E_8 .




FIGURE 6. The two 11-vertex 1-Salem graphs that are not generalised line graphs.

Choose a $S \subseteq W(E_8)$. When considering if it is possible to add a new vector to ones list, apply each isometry in S to the potential enlarged list, and reject the new vector if any of the permuted lists appears earlier in the lexicographical ordering derived from our ordering of the vectors. Larger sets S make each update more expensive, but potentially reduce the backtracking by pruning certain branches at an early stage.

We used a few other programming tricks to speed up the process further, but the main idea was to exploit isometries of E_8 . In practice we used a set S of size 162, including the 120 involutions above, and 42 carefully selected others.

6.3. 1-Salem graphs represented in E_8 . The property of being a 1-Salem graph is not hereditary, but the property of being either 1-Salem or cyclotomic is. We used the fast backtracking approach of §6.2 to find all such graphs represented in E_8 , and then weeded out those that were cyclotomic, and also those that were generalised line graphs (and hence covered by part (ii) of Theorem 1). Table 2 summarises the results, showing the number of non-bipartite connected 1-Salem graphs that are not generalised line graphs by the number of vertices. The largest examples have 11 vertices; they are shown in Figure 6. These computations complete the final part of the proof of Theorem 1, which we state here as a theorem.

Theorem 11. *There are 377 non-bipartite 1-Salem graphs that are not generalised line graphs. The numbers of vertices for these graphs range between 6 and 11; the number of graphs for each of these numbers of vertices is shown in Table 2.*

Number of vertices	6	7	8	9	10	11
Number of graphs	10	43	111	153	58	2

TABLE 2. The number of non-bipartite 1-Salem graphs that are not generalised line graphs.

7. ACKNOWLEDGEMENT

We are grateful to Jonathan Cooley, who verified several of our computations independently, hugely increasing our confidence in their accuracy. We thank the referee for helpful suggestions about the structure of the paper, and for drawing our attention to some further related work.

REFERENCES

- [1] B. Bollobás, *Graph theory: an introductory course*, Graduate Texts in Mathematics **63**, Springer, New York, 1979.
- [2] D.W. Boyd, Small Salem numbers, *Duke Math. J.* **44**(2) (1977), 315–328.

- [3] P.J. Cameron, J.M. Goethals, J.J. Seidel, E.E. Shult, Line graphs, root systems, and elliptic geometry, *J. Algebra* **43** (1976), 305–327.
- [4] P.J. Cameron, J.H. van Lint, *Designs, graphs, codes and their links*, Cambridge University Press, Cambridge, 1991.
- [5] A.L. Cauchy, Sur l'équation à l'aide de laquelle on détermine les inégalités séculaires des mouvements des planètes, Oeuvres complètes, Iième Série, **9**, 174–195, Gauthier-Villars, 1829.
- [6] T. Chung, J. Koolen, Y. Sano, T. Taniguchi, The non-bipartite integral graphs with spectral radius three, *Linear Algebra and its Applications* **435**(10) (2011), 2544–2559.
- [7] D. Cvetković, M. Doob, I. Gutman, On graphs whose eigenvalues do not exceed $\sqrt{2 + \sqrt{5}}$, *Ars Comb.* **14** (1982), 225–239.
- [8] D. Cvetković, M. Doob, S. Simić, Generalised line graphs, *J. Graph Theory* **5** (1981), 385–399.
- [9] D. Cvetković, P. Rowlinson, S. Simić, *Spectral generalizations of line graphs: on graphs with least eigenvalue -2* , London Mathematical Society Lecture Note Series **314**, Cambridge University Press, Cambridge, 2004.
- [10] J.W. Cannon, Ph. Wagerich, Growth functions of surface groups, *Math. Ann.* **293** (1992), 239–257.
- [11] E. Dobrowolski, A note on integer symmetric matrices and Mahler's measure, *Canad. Math. Bull.* **51**(1) (2008), 57–59.
- [12] D.R. Estes, Eigenvalues of symmetric integer matrices, *J. Number Theory* **42** (1992), 292–296.
- [13] C. Godsil, B. McKay, A new graph product and its spectrum, *Bull. Australian Math. Soc.* **18**(01) (1978), 21–28.
- [14] C. Godsil, G. Royle, *Algebraic Graph Theory*, Springer, New York, 2001.
- [15] L. Gumbrell *On spectral constructions for Salem graphs*, PhD thesis, 2013.
- [16] F. Harary, *Graph Theory*, Addison-Wesley, 1969.
- [17] A.J. Hoffman, On limit points of spectral radii of non-negative symmetric matrices, in *Graph Theory and its Applications*, Lecture Notes in Math. 303, 165–172, Springer, Berlin, 1972.
- [18] A.J. Hoffman, Eigenvalues of graphs, in *Studies in Graph Theory*, 225–245, ed. D.R. Fulkerson, Math. Assoc. Amer., Washington, 1975.
- [19] A.J. Hoffman, J.H. Smith, On the spectral radii of topologically equivalent graphs, in *Recent advances in graph theory*, 273–281, ed. M. Fiedler, Academia Praha, 1975.
- [20] L. Kronecker, Zwei sätze über gleichungen mit ganzzahligen coefficienten, *J. Reine Angew. Math.* **53** (1857), 173–175.
- [21] P. Lakatos, On spectral radius of Coxeter transformation of trees, *Publicationes Mathematicae Debrecen*, **54** (1999), 181–187.
- [22] P. Lakatos, Salem numbers, PV numbers and spectral radii of Coxeter transformations, *Comptes Rendus Math. de L'Acad. des Sciences La Soc. Royale du Canada*, **23** (2001), 71–77.
- [23] P. Lakatos, Salem numbers defined by Coxeter transformation, *Linear Algebra and its Appl.*, **432** (2010), 144–154.
- [24] J. McKee, Families of Pisot numbers with negative trace, *Acta Arith.*, **93**(4), (2000), 373–385.
- [25] J. McKee, C. Smyth, P. Rowlinson, Salem numbers and Pisot numbers from stars, in *Number Theory in Progress*, vol. 1 (Zakopane-Kocielisko, 1997), ed. K. Györy, H. Iwaniec, J. Urbanowicz, 309–319, de Gruyter, Berlin, 1999.
- [26] J. McKee, Small-span characteristic polynomials of integer symmetric matrices, in *Algorithmic Number Theory*, ed. G. Hanrot, F. Morain, E. Thomé, 9th International Symposium, ANTS-IX, Nancy, France, Springer Lecture Notes in Computer Science **6197** (2010), 270–284.
- [27] J. McKee, C. Smyth, Salem numbers of trace -2 , and traces of totally positive algebraic integers, in *Algorithmic Number Theory*, ed. D.A. Buell, 6th International Symposium, ANTS-VI, Burlington, USA, Springer Lecture Notes in Computer Science **3076** (2004), 327–337.
- [28] J. McKee, C. Smyth, Salem numbers, Pisot numbers, Mahler measure, and graphs, *Experimental Mathematics* **14** (2005), 211–229.
- [29] J. McKee, C. Smyth, Integer symmetric matrices having all their eigenvalues in the interval $[-2, 2]$, *J. Algebra* **317** (2007), 260–290.
- [30] J. McKee, C. Smyth, Integer symmetric matrices of small spectral radius and small Mahler measure, *Int. Math. Research Notices* (2011); doi: 10.1093/imrn/rnr011.

- [31] C.T. McMullen, Coxeter groups, Salem numbers and the Hilbert metric, *Publ. Math. Inst. de Hautes Etudes Scientifiques* **95** (2002), 151–183.
- [32] W. Parry, Growth series of Coxeter groups and Salem numbers, *J. Algebra* **154** (1993), 406–415.
- [33] V. Prasolov, S. Ivanov, *Problems and theorems in linear algebra*, American Mathematical Society, 1994.
- [34] Z. Radosavljević, S. Simić, Which bicyclic graphs are reflexive, *Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat.* **14** (2003), 64–85.
- [35] S.B. Rao, N.M. Vinghi, K.S. Vijayan, The minimal forbidden subgraphs for generalized line-graphs, in *Combinatorics and Graph Theory, Proc. Symp. Indian Statistical Institute, Calcutta, 25–29 February 1980*, edited by S.B. Rao, 459–472, Lecture Notes in Math. **855**, Springer, Berlin, 1981.
- [36] M. Rašajski, Z. Radosavljević, B. Mihailović, Maximal reflexive cacti with four cycles: The approach via Smith graphs, *Linear Algebra and its Applications* **435** (2011), 2530–2543.
- [37] R.M. Robinson, Intervals containing infinitely many sets of conjugate algebraic integers, *Mathematical analysis and related topics: essays in honor of George Pólya*, 305–315, Stanford, 1962.
- [38] I. Schur, Über die Verteilung der Wurzeln bei gewissen algebraischen Gleichungen mit ganzzahligen Koeffizienten, *Math. Z.* **1** (1918), 377–402.
- [39] J.H. Smith, Some properties of the spectrum of a graph, in *Combinatorial structures and their applications (Proc. Calgary Internat. Conf., Calgary, Alta., 1969)*, edited by R. Guy, H. Hanani, H. Saver, J. Schönheim, 403–406, Gordon and Breach, New York, 1970.
- [40] R. Stekolshchik, *Notes on Coxeter transformations and the McKay correspondence*, Springer, Berlin Heidelberg, 2008.
- [41] Y. Zhang, Eigenvalues of Coxeter transformations and the structure of regular components of an Auslander-Reiten quiver, *Comm. Algebra* **17**(10) (1989), 2347–2362.

DEPARTMENT OF MATHEMATICS, ROYAL HOLLOWAY, UNIVERSITY OF LONDON, EGHAM HILL, EGHAM, SURREY, TW20 0EX, ENGLAND, UK.

E-mail address: Lee.Gumbrell@gmail.com

DEPARTMENT OF MATHEMATICS, ROYAL HOLLOWAY, UNIVERSITY OF LONDON, EGHAM HILL, EGHAM, SURREY, TW20 0EX, ENGLAND, UK.

E-mail address: James.McKee@rhul.ac.uk