

THE ACTION OF GENERALISED SYMMETRIC GROUPS ON SYMMETRIC AND EXTERIOR POWERS OF THEIR NATURAL REPRESENTATIONS

WILLIAM O'DONOVAN

ABSTRACT. We establish upper and lower bounds on the dimension of the space spanned by the symmetric powers of the natural character of generalised symmetric groups. We adapt the methods of Savitt and Stanley from [1] to obtain bounds both over the complex numbers and in prime characteristic.

1. INTRODUCTION

In [1], Savitt and Stanley proved that the space spanned by the symmetric powers of the standard n -dimensional complex representation of S_n has dimension asymptotic to $\frac{n^2}{2}$. In this paper, we look at some extensions of this, as well as demonstrate the wide applicability of the methods employed in [1] for upper bounds. The methods used in [1] to obtain lower bounds are not so readily applicable, but some partial results can be found. The study of the symmetric powers of the natural representation of S_n is motivated in part by the fact that if λ is a partition and $r \geq \sum_{i=1}^{\lambda_1} \binom{\lambda_i}{2}$, then $\text{Sym}^r V$ contains the Specht module S^λ .

In Section 2 of this paper, we consider the action of the generalised symmetric groups $C_k \wr S_n$, on the symmetric and exterior powers of their natural representations as $n \times n$ complex matrices. We prove that, to leading order, the space spanned by the characters of the symmetric powers of the natural representation of $C_k \wr S_n$ has dimension between $kn \log n$ and $\frac{kn^2}{2}$. If $\lambda = (\lambda_1, \dots, \lambda_s)$ is a partition, then we define $f_\lambda(x) := \prod_{j=1}^s (1 - x^{\lambda_j})^{-1}$.

Date: February 11, 2016.

2010 Mathematics Subject Classification. 20C30

Key words and phrases. Symmetric power, wreath product, exterior power, generalised symmetric group, Brauer character.

The identity (where $\sigma \in S_n$ has cycle type λ and π denotes the character of the standard representation of S_n)

$$\sum_r (\text{Sym}^r \pi)(\sigma) q^r = p_\lambda(1, q, q^2, \dots) = f_\lambda(q),$$

plays an important part of the analysis in [1]; in Section 2 we obtain a generalisation of this for wreath products. Furthermore, we show that if $k \geq 2$, then the space spanned by the characters of the exterior powers of this representation has maximum possible dimension, namely $n + 1$. In contrast, the space spanned by the exterior powers of the natural character of S_n (the case when $k = 1$) has dimension n .

In Section 3 of this paper, we examine the same issue explored in [1], but this time in positive characteristic, and prove that the space spanned by the symmetric powers of the natural representation of S_n in characteristic p is bounded (to leading order) by $\frac{p}{2p+2}n^2$. Although the argument for lower bounds given in [1] cannot be modified for this case, we use a different method to establish a lower bound of (again to leading order) $\frac{p-1}{p}n \log n$. Numerical data suggests that this lower bound can be improved upon and that the upper bound is closer to the truth. We conclude by putting both extensions together: we consider the symmetric powers of $C_k \wr S_n$ in characteristic p and prove that the dimension of the space spanned by the symmetric powers is bounded above, to leading order, by $r_p(k) \frac{p}{2p+2}n^2$, where $r_p(k)$ is the p' -part of k , as defined in Definition 9.

2. GENERALISED SYMMETRIC GROUPS

2.1. Symmetric Powers. We consider an extension of [1], namely to the generalised symmetric group $C_k \wr S_n$ for $k \in \mathbb{N}$; we think of this group as the subgroup of $GL_n(\mathbb{C})$ consisting of matrices whose entries are k^{th} roots of unity, and which have exactly one non-zero entry in each row and column.

Definition 1. Given $\sigma \in C_k \wr S_n$, we define $|\sigma|$ to be the element of S_n corresponding to the permutation matrix we obtain by setting all of the

non-zero entries of σ equal to one. We also define a map $\lambda : C_k \wr S_n \rightarrow \text{Par}(n)$ by sending $\sigma \in C_k \wr S_n$ to the cycle type of $|\sigma|$. If $\lambda(\sigma) = (\lambda_1, \dots, \lambda_s)$, let $t_i \in \mathbb{C}$ denote the product of the entries in σ which correspond to the cycle of $|\sigma|$ of length λ_i . Where $\lambda(\sigma)$ has a repeated part, say $\lambda_i = \lambda_{i+1}$, then we do not order t_i and t_{i+1} . In other words, we regard the two s -tuples $(t_1, \dots, t_i, t_{i+1}, \dots, t_s)$ and $(t_1, \dots, t_{i+1}, t_i, \dots, t_s)$ as equivalent.

For example, if $k = 2$ and

$$\sigma = \begin{pmatrix} 0 & -1 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 1 & 0 \end{pmatrix}$$

then $|\sigma| = (12)(3)(45) \in S_5$, $\lambda(\sigma) = (2, 2, 1)$, $t_1 = 1$, $t_2 = -1$ and $t_3 = 1$; equivalently, as $\lambda(\sigma)$ has a repeated part, we could take $t_1 = -1$, $t_2 = 1$ and $t_3 = 1$. The conjugacy class of $\sigma \in C_k \wr S_n$ is determined by $\lambda(\sigma) = (\lambda_1, \dots, \lambda_s)$ and a representative of the equivalence class of $t = (t_1, \dots, t_s)$ under the equivalence relation described above. Also, such pairs (λ, t) where $\lambda = (\lambda_1, \dots, \lambda_s) \in \text{Par}(n)$ and $t = (t_1, \dots, t_s)$, where each t_i is a k^{th} root of unity, parametrise the conjugacy classes. For details see [4, Section 4.2].

Let V be the natural representation of $C_k \wr S_n$ with basis e_1, \dots, e_n and let $\text{Sym}^r V$ be its r^{th} symmetric power with character χ_r . Let $\sigma \in C_k \wr S_n$ and consider the action of σ on $\text{Sym}^r V$. Now, $\text{Sym}^r V$ has a basis $\{e_1^{c_1} \dots e_n^{c_n} : \sum_i c_i = r\}$, and we want to know when a basis vector contributes to the trace of the matrix of σ acting on $\text{Sym}^r V$. Observe that, with this basis, σ always maps a basis vector to a multiple of another basis vector; consequently, the only contribution to the trace comes from those basis vectors which are eigenvectors for the action of σ .

Suppose, without loss of generality, that $|\sigma| = (1, 2, \dots, \lambda_1) \dots (\lambda_1 + \dots + \lambda_{s-1} + 1, \dots, n)$, so $\lambda(\sigma) = (\lambda_1, \dots, \lambda_s)$. Then the vector $v = e_1^{c_1} \dots e_n^{c_n}$ is

an eigenvector for σ if and only if $c_1 = \dots = c_{\lambda_1}$, $c_{\lambda_1+1} = \dots = c_{\lambda_1+\lambda_2}$ and so on. Hence the eigenvectors for σ are in a one-to-one correspondence with the set of non-negative integral solutions (c_1, \dots, c_s) to $\lambda_1 c_1 + \dots + \lambda_s c_s = r$.

Suppose that v is a basis vector which is an eigenvector for the action of σ . Then we have $\sigma.v = \prod_{i=1}^s t_i^{c_i} v$, and so $\chi_r(\sigma) = \sum \prod_{i=1}^s t_i^{c_i}$, where the sum runs over all the basis vectors of $\text{Sym}^r V$ which are eigenvectors for the action of σ . It follows that

$$\chi_r(\sigma) = \sum_{(c_1, \dots, c_s)} \prod_{i=1}^s t_i^{c_i},$$

where the sum is over all non-negative integral solutions to the equation $\lambda_1 c_1 + \dots + \lambda_s c_s = r$. Define the power series

$$f_\sigma(x) = \frac{1}{(1 - t_1 x^{\lambda_1}) \dots (1 - t_s x^{\lambda_s})}.$$

The coefficient of x^m in the power series $f_\sigma(x)$ is

$$\sum_{(c_1, \dots, c_s)} \prod_{i=1}^s t_i^{c_i} = \chi_m(\sigma),$$

where the sum is over all solutions to the equation $\lambda_1 c_1 + \dots + \lambda_s c_s = m$. Consequently, this power series is the generating function for the values of $\chi_r(\sigma)$.

Let $D(n, k)$ be the dimension of the space spanned by the characters of the symmetric powers of the natural representation for $C_k \wr S_n$, i.e. $D(n, k) = \dim(\text{Span}_{\mathbb{C}}\{\chi_r : r \in \mathbb{N}\})$. Consider the matrix of the characters χ_r for $r \in \mathbb{N}$; $D(n, k)$ is equal to the row-rank of this matrix. Since row-rank and column-rank are the same, $D(n, k)$ equals the dimension of the space spanned by the columns of the table. By the above argument, the generating function for the entries of the column corresponding to the conjugacy class of $\sigma \in C_k \wr S_n$ is $f_\sigma(x)$. We have therefore proved:

Proposition 2. $D(n, k) = \dim(\text{Span}_{\mathbb{C}}\{f_\sigma(x) : \sigma \in C_k \wr S_n\})$.

Proposition 3. We have $D(n, k) \leq \frac{kn^2}{2} + (\frac{k}{2} - 1)n + 1$.

Proof. Let $\zeta = \exp(\frac{2\pi i}{k})$ and consider $\{f_\sigma(x) : \sigma \in C_k \wr S_n\}$; we claim that a common denominator for these power series is

$$D(x) := \prod_{\substack{0 \leq i < k \\ 1 \leq j \leq n}} (1 - \zeta^i x^j).$$

Suppose that $\theta \in \mathbb{C}$ and $x - \theta$ divides $\frac{1}{f_\sigma(x)}$ in $\mathbb{C}[x]$, for some $\sigma \in C_k \wr S_n$. Let r be the minimal natural number such that, in $\mathbb{C}[x]$, $x - \theta$ divides $1 - \zeta^i x^r$ for some $0 \leq i < k$. Then, in a factorisation over \mathbb{C} into linear factors of a common denominator for the $f_\sigma(x)$, the factor $x - \theta$ appears exactly $\lfloor \frac{n}{r} \rfloor$ times by minimality of r . Since $\zeta^i \theta^r = 1$, it follows that for any $m \in \mathbb{N}$, $\zeta^{im} \theta^{rm} = 1$, whence $x - \theta$ divides $1 - \zeta^{im} x^{rm}$ in $\mathbb{C}[x]$. This is a factor of $D(x)$ for $m = 1, \dots, \lfloor \frac{n}{r} \rfloor$, so $(x - \theta)^{\lfloor \frac{n}{r} \rfloor}$ divides $D(x)$ in $\mathbb{C}[x]$, and this establishes the claim.

The elements $\frac{1}{D(x)}, \frac{x}{D(x)}, \dots, \frac{x^{\deg D - n}}{D(x)}$ form a spanning set for the vector space $\text{Span}_{\mathbb{C}}\{f_\sigma(x) : \sigma \in C_k \wr S_n\}$, giving $D(n, k) \leq \deg D + 1 - n$. Note that $D(x)$ has degree $k \sum_{j=1}^n j = \frac{kn}{2}(n+1)$, and the result follows. \square

We observe that the arguments given to prove the above two propositions are generalisations of those used in [1, Section 1] and [1, Proposition 2.1] respectively.

Although the method used in [1, Section 3] to obtain a lower bound is not adaptable to this case (see also Remark 13), we can establish a lower bound by other methods.

Proposition 4. *There exists a constant $c > 0$ such that $D(n, k) \geq \max(\frac{n^2}{2} - cn^{\frac{3}{2}}, k((n-1)\log(n-1) - 2(n-2)))$.*

Proof. Since $\{f_\lambda(x) : \lambda \in \text{Par}(n)\}$ is a subset of $\{f_\sigma(x) : \sigma \in C_k \wr S_n\}$, we get the lower bound $D(n, k) \geq \frac{n^2}{2} - cn^{\frac{3}{2}}$ for some constant c by [1, Proposition 3.3]. Let W_{n-1} denote the set of partitions of $n-1$ which have the form $(a^b, 1^{n-1-ab})$, where $a, b \in \mathbb{N}_0$ and $a > 1$, and let $\zeta = \exp(\frac{2\pi i}{k})$. We claim that $Y := \{\frac{1}{1-\zeta^r x} f_\lambda(x) : 0 \leq r < k, \lambda \in W_{n-1}\}$ is a linearly independent

set. We prove this by induction on the position of the partition λ in the lexicographic ordering on $\text{Par}(n-1)$.

Suppose that for some constants $\alpha_{\lambda,r}$, we have

$$\sum_{\lambda \in W_{n-1}} \sum_{r=0}^{k-1} \alpha_{\lambda,r} \frac{f_{\lambda}(x)}{1 - \zeta^r x} = 0.$$

Let μ_1 be the partition $(n-1) \in W_{n-1}$, then we can re-write this as

$$-\sum_{r=0}^{k-1} \frac{\alpha_{\mu_1,r}}{(1 - \zeta^r x)(1 - x^{n-1})} = \sum_{\lambda \in W_{n-1} \setminus \{\mu_1\}} \sum_{r=0}^{k-1} \alpha_{\lambda,r} \frac{f_{\lambda}(x)}{1 - \zeta^r x}.$$

The left-hand side has a singularity at $\exp(\frac{2\pi i}{n-1})$, whereas the right-hand side is holomorphic at this point. It follows that $\sum_{r=0}^{k-1} \frac{\alpha_{\mu_1,r}}{1 - \zeta^r x} = 0$. Since the function $\frac{1}{1 - \zeta^r x}$ has a pole at ζ^{-r} only, we see that the power series $\frac{1}{1 - \zeta^r x}$ for $0 \leq r < k$ are linearly independent. Hence $\alpha_{\mu_1,r} = 0$ for all r .

Let $\mu = (a^b, 1^{n-ab-1})$ and suppose that $\alpha_{\lambda,r} = 0$ for every partition $\lambda \in W_{n-1}$ which is greater than μ in the lexicographic ordering. Then we can write

$$-\sum_{r=0}^{k-1} \frac{\alpha_{\mu,r}}{(1 - \zeta^r x)(1 - x^a)^b(1 - x)^{n-ab-1}} = \sum_{\lambda < \mu} \sum_{r=0}^{k-1} \alpha_{\lambda,r} \frac{f_{\lambda}(x)}{1 - \zeta^r x}.$$

The left-hand side has a pole of order b at $\exp(\frac{2\pi i}{a})$. There are two cases. If the right-hand side is holomorphic at $\exp(\frac{2\pi i}{a})$, then we deduce that $\alpha_{\mu,r} = 0$ by the same argument given in the base case, and so we may suppose that the right-hand side has a pole at $\exp(\frac{2\pi i}{a})$. However, if the pole at $\exp(\frac{2\pi i}{a})$ on right-hand side were of order b or greater, then there would be a partition $\lambda \in W_{n-1}$ with the part a repeated at least b times. By construction of the set W_{n-1} , this is impossible since $\lambda < \mu$.

Hence the right-hand side has a pole at $\exp(\frac{2\pi i}{a})$ of order strictly less than b . Consequently, we deduce that $\alpha_{\mu,r} = 0$. It follows by induction that all the $\alpha_{\lambda,r}$ are equal to zero. This proves the claim.

Therefore, we have $D(n, k) \geq k|W_{n-1}|$. It is clear that

$$|W_m| = 1 + \sum_{t=2}^m \lfloor \frac{m}{t} \rfloor \geq 1 + \sum_{t=2}^m (\frac{m}{t} - 1).$$

Therefore, we have $|W_m| \geq 1 + mH_m - m - (m - 1)$, where H_m is the m^{th} harmonic number, and so, using the simple bound $H_m \geq \log m$, it follows that $|W_m| \geq m \log m - 2(m - 1)$. Consequently, we deduce that $D(n, k) \geq k((n - 1) \log(n - 1) - 2(n - 2))$, as required. \square

Remark 5. In the above proof we used that the power series $\frac{1}{1-\zeta^r x}$ for $r = 1, \dots, k$ are linearly independent, whence $D(1, k) = k$. In particular, the upper bound $D(n, k) \leq \frac{kn^2}{2} + (\frac{k}{2} - 1)n + 1$ is an equality when $n = 1$.

2.2. Exterior Powers. Let $\Lambda^r V$ be the r^{th} exterior power of the natural representation of $C_k \wr S_n$ with basis $\{e_{i_1} \wedge \dots \wedge e_{i_k} : 1 \leq i_1 < \dots < i_k \leq n\}$ and character ψ_r .

Note that with this basis of $\Lambda^r V$, each $\sigma \in C_k \wr S_n$ again sends a basis vector to a multiple of a basis vector, so the only contribution to the trace of the matrix of σ on $\Lambda^r V$ is from those basis vectors which are eigenvectors for the action of σ . Let a basis vector be $v = e_1^{c_1} \wedge e_2^{c_2} \wedge \dots \wedge e_n^{c_n}$, where $c_i \in \{0, 1\}$ and we interpret e_i^0 to mean that e_i does not appear (for example, $e_1^1 \wedge e_2^0 \wedge e_3^1 = e_1 \wedge e_3$). Without loss of generality, suppose that $|\sigma| = (1, \dots, \lambda_1) \dots (\lambda_1 + \dots + \lambda_{s-1} + 1, \dots, n)$, so $\lambda(\sigma) = (\lambda_1, \dots, \lambda_s)$.

Observe that, for every j such that $c_j = 1$, we must have $c_k = 1$ whenever k is in the same cycle as j in $|\sigma|$. Therefore, the eigenvectors correspond to solutions to the equation $\lambda_1 c_1 + \dots + \lambda_s c_s = r$ where each c_i is either 0 or 1. Moreover, a m -cycle in S_n with $c_i = 1$ on its support sends v to $(-1)^{m-1}v$ since an m -cycle has sign $(-1)^{m-1}$.

Then $\sigma.v = \prod_i (-1)^{\lambda_i - 1} t_i$, where the product is over $1 \leq i \leq s$ such that $c_i = 1$ and t_i is as in Definition 1. It follows that

$$\psi_r(\sigma) = \sum_{(c_1, \dots, c_s)} \prod_i (-1)^{\lambda_i - 1} t_i,$$

where the sum is over solutions (c_1, \dots, c_s) to the equation $\lambda_1 c_1 + \dots + \lambda_s c_s = r$ and the product is again over $1 \leq i \leq s$ such that $c_i = 1$.

Given $\sigma \in C_k \wr S_n$ with $\lambda(\sigma) = (\lambda_1, \dots, \lambda_s)$ and associated s -tuple of k^{th} roots of unity (t_1, \dots, t_s) , we define

$$g_\sigma(x) = \prod_{i=1}^s (1 + t_i (-1)^{\lambda_i - 1} x^{\lambda_i}).$$

A similar argument to that used to prove Proposition 2 shows that $g_\sigma(x)$ is the generating function for the values of $\psi_r(\sigma)$. Let $E(n, k)$ be the dimension of the space spanned by the characters of the exterior powers of the natural representation for $C_k \wr S_n$. Then we have proved:

Proposition 6. $E(n, k) = \dim(\text{Span}_{\mathbb{C}}\{g_\sigma(x) : \sigma \in C_k \wr S_n\})$.

From this, we are able to deduce the following:

Corollary 7. For any $k \geq 2$, $E(n, k) = n + 1$.

Proof. Let $\zeta = \exp(\frac{2\pi i}{k})$. We consider two cases; first suppose that n is even. Then for $j = 1, \dots, \frac{n}{2}$, we define the following $n + 1$ polynomials: $a_j(x) = (1 + (-1)^{n-j} x^{n-j})(1 + (-1)^j \zeta x^j)$, $b_j(x) = (1 + (-1)^{n-j} x^{n-j})(1 + (-1)^j \zeta^2 x^j)$ together with $1 - \zeta x^n$, giving $n + 1$ polynomials which have the form $g_\sigma(x)$ for some $\sigma \in C_k \wr S_n$. We claim these polynomials are linearly independent.

Suppose that for some constants $\alpha_j, \beta_j, \gamma$, we have

$$\sum_{j=1}^{\frac{n}{2}} \alpha_j a_j(x) + \sum_{j=1}^{\frac{n}{2}} \beta_j b_j(x) + \gamma(1 - \zeta x^n) = 0.$$

Equating coefficients of x and x^{n-1} gives us (after simplifying) the equations $\alpha_1 \zeta + \beta_1 \zeta^2 = 0$ and $\alpha_1 + \beta_1 = 0$, from which we deduce that $\alpha_1 = \beta_1 = 0$. We similarly deduce $\alpha_j = \beta_j = 0$ for $1 \leq j < \frac{n}{2}$ by equating coefficients of x^j and x^{n-j} . This leaves three equations in $\alpha_{\frac{n}{2}}, \beta_{\frac{n}{2}}, \gamma$, from which it is trivial to show that these constants are all zero, thus proving the claim.

If, however, n is odd, then we consider the polynomials $a_j(x), b_j(x)$ for $j = 0, 1, \dots, \lfloor \frac{n}{2} \rfloor$, together with $1 - \zeta x^n$ and $1 - \zeta^2 x^n$. Again this gives $n + 1$

polynomials which are of the form $g_\sigma(x)$, and a similar argument to that given for the previous case establishes that they are linearly independent. \square

3. SYMMETRIC GROUPS IN PRIME CHARACTERISTIC

3.1. Symmetric group. The arguments used in [1] worked over the complex numbers, and so it is natural to ask what happens in prime characteristic. To obtain a good theory of characters in this case, we need to work with Brauer characters. Let $\beta_{r,p}$ denote the Brauer character of the r^{th} symmetric power of the natural representation of S_n in characteristic p .

For a full account of Brauer characters see [2, Chapter 15]; all we need to know about Brauer characters for our purposes is the following:

Lemma 8. *If $g \in S_n$ has order coprime to p , then $\beta_{r,p}(g) = \chi_r(g)$.*

Proof. This is a special case of the fact that the Brauer character of the p -modular reduction of a characteristic zero representation affording χ equals χ on the conjugacy classes of p -regular elements. For a proof, see [2, Theorem 15.8]. \square

Let $B(p, n)$ denote the dimension of the space spanned by the Brauer characters of the symmetric powers of the natural representation of S_n in characteristic p . An element $g \in S_n$ has order coprime to p if and only if its cycle type has all its parts coprime to p . Let X_p denote the set of partitions of n whose parts are all coprime to p . The usual argument shows that $\dim(\text{Span}_{\mathbb{C}}\{f_\lambda : \lambda \in X_p\}) = B(p, n)$.

We aim to put the rational functions $f_\lambda(x)$ (for $\lambda \in X_p$) over their least common denominator, which we denote by $g_p(x)$. If $g_p(x)$ has degree δ_p , then the elements $\frac{1}{g_p(x)}, \frac{x}{g_p(x)}, \dots, \frac{x^{\delta_p-n}}{g_p(x)}$ are a spanning set for $\text{Span}_{\mathbb{C}}\{f_\lambda : \lambda \in X_p\}$, giving $B(p, n) \leq \delta_p + 1 - n$. Our strategy is as follows: to work out what $g_p(x)$ is (up to a scalar), and then to bound its degree.

Definition 9. Let $k \in \mathbb{N}$, and write $k = p^a m$ where m is coprime to p . Then we define $r_p(k) = m$.

Proposition 10. *We have $g_p(x) = \pm \prod_{k \leq n} (1 - x^{r_p(k)})$.*

Proof. For $\lambda \in X_p$, the denominator of $f_\lambda(x)$ is a polynomial of degree n which factorises over \mathbb{Z} as a product of cyclotomic polynomials. If p divides d , then the d^{th} cyclotomic polynomial $\Phi_d(x)$ does not appear in this factorisation, whereas if p does not divide d , then $\Phi_d(x)$ appears at most $\lfloor \frac{n}{d} \rfloor$ times in the common denominator. It follows that

$$g_p(x) = \prod_{\substack{1 \leq k \leq n, \\ (k,p)=1}} \Phi_k(x)^{\lfloor \frac{n}{k} \rfloor}.$$

Note that $\prod_{k \leq n} (1 - x^{r_p(k)})$ is also a product of cyclotomic polynomials and $\Phi_d(x)$ appears if and only if $r_p(k)$ is a multiple of d . If p divides d , then $r_p(k)$ is not a multiple of d ; conversely, if p does not divide d , then whenever k is a multiple of d , $r_p(k)$ is a multiple of d . Since there are $\lfloor \frac{n}{d} \rfloor$ multiples of d between 1 and n , it follows that the d^{th} cyclotomic polynomial appears $\lfloor \frac{n}{d} \rfloor$ times in $\prod_{k \leq n} (1 - x^{r_p(k)})$. \square

Proposition 11. *The degree of $g_p(x)$ is at most $\frac{p}{2p+2}n^2 + \frac{pn}{p-1}$.*

Proof. Let $m \in \mathbb{N}$ with $m \leq n$. The equation $r_p(k) = m$ has no solutions if m is divisible by p and $1 + \lfloor \log_p(\frac{n}{m}) \rfloor$ solutions if p does not divide m , giving

$$g_p(x) = \prod_{\substack{1 \leq k \leq n, \\ (k,p)=1}} (1 - x^k)^{1 + \lfloor \log_p(\frac{n}{k}) \rfloor}.$$

Hence the degree of $g_p(x)$, namely δ_p , is

$$\sum_{\substack{1 \leq k \leq n, \\ (k,p)=1}} k(1 + \lfloor \log_p(\frac{n}{k}) \rfloor).$$

Let $l = \lfloor \log_p n \rfloor$; we can rewrite the above as

$$\delta_p = \sum_{\substack{1 \leq k \leq n, \\ (k,p)=1}} k + \sum_{\substack{1 \leq k \leq \frac{n}{p}, \\ (k,p)=1}} k + \cdots + \sum_{\substack{1 \leq k \leq \frac{n}{p^l}, \\ (k,p)=1}} k.$$

It is easy to show that

$$\sum_{\substack{1 \leq k \leq x, \\ (k,p)=1}} k \leq \frac{p-1}{2p} x^2 + x.$$

Therefore, we have

$$\delta_p \leq \frac{p-1}{2p} \left(n^2 + \frac{n^2}{p^2} + \cdots + \frac{n^2}{p^{2l}} \right) + \left(n + \frac{n}{p} + \cdots + \frac{n}{p^l} \right).$$

Observe that the first bracketed sum is bounded above by the infinite geometric series $\sum_{i=0}^{\infty} \frac{n^2}{p^{2i}}$, and the second sum is bounded by $\sum_{i=0}^{\infty} \frac{n}{p^i}$. This gives $\delta_p \leq \frac{p-1}{2p} \frac{n^2}{1-p^{-2}} + \frac{pn}{p-1}$. After some simple algebra, we see that $\delta_p \leq \frac{p}{2p+2} n^2 + n(\log_p n + 1)$, as required. \square

Recall that just before Definition 9, we established that $B(p, n) \leq \delta_p + 1 - n$, and so the following result is immediate:

Corollary 12. *For every prime number p and $n \in \mathbb{N}$, we have $B(p, n) \leq \frac{p}{2p+2} n^2 + \frac{n}{p-1} + 1$.*

Hence we have obtained an upper bound on $B(p, n)$ which is asymptotic to $\frac{pn^2}{2p+2}$; this is always smaller than the $\frac{n^2}{2}$ obtained in the characteristic zero case in [1]. It is also worth noting that as p becomes very large, this upper bound becomes close to the one obtained for characteristic zero. We now aim to find a lower bound on $B(p, n)$.

Remark 13. It is, at first sight, natural to try to imitate the argument for a lower bound given in [1, Section 3] by considering the dimension of the vector space $\text{Span}_{\mathbb{C}}\{e_{\lambda}(1, x, x^2, \dots) : \lambda \in X_p\}$, where e_{λ} is the elementary symmetric function of type λ . However, the proof given relies on making a change of basis for the vector space Λ^n of symmetric functions of degree n . Unfortunately, this basis change is not compatible with the requirement that all parts of λ be coprime to p . In other words, while $\{e_{\lambda}(1, x, x^2, \dots) : \lambda \in \text{Par}(n)\}$ is a basis for the \mathbb{C} -vector space with basis $\{f_{\lambda}(x) : \lambda \in \text{Par}(n)\}$,

this statement is false if we replace $\text{Par}(n)$ by X_p . Therefore, our strategy is to exhibit a large linearly independent subset of $\text{Span}_{\mathbb{C}}\{f_\lambda : \lambda \in X_p\}$.

Given $n \in \mathbb{N}$ and a prime number p , we define $A_{p,n}$ to be the set of partitions of n which have the form $(r^a, 1^b)$, where r is coprime to p , and $a, b \in \mathbb{N}_0$.

Proposition 14. *We have $B(p, n) \geq \frac{p-1}{p}n \log n + n(\gamma - 1 - \frac{p-1}{p} - \frac{\gamma}{p})$, where γ denotes the Euler–Mascheroni constant.*

Proof. Suppose that in $\mathbb{Q}(x)$, we have $\sum_{\mu \in A_{p,n}} \alpha_\mu f_\mu(x) = 0$ for $\alpha_\mu \in \mathbb{Q}$. The set $A_{p,n}$ is totally ordered by the lexicographic order on partitions; let $\mu_1 = (r^a, 1^b)$ be the largest partition in $A_{p,n}$. Then in $\mathbb{Q}[x]$ we have that $\Phi_r(x)^a$ divides the denominator of $f_{\mu_1}(x)$, and $\Phi_r(x)^a$ does not divide the denominator of any other $f_\mu(x)$ because μ_1 is the largest partition in $A_{p,n}$.

Assume that $\alpha_{\mu_1} \neq 0$. Then we may write

$$\sum_{\mu \in A_{p,n} \setminus \{\mu_1\}} \alpha_\mu f_\mu(x) = -\alpha_{\mu_1} f_{\mu_1}(x).$$

Putting the left-hand side over a common denominator and multiplying both sides by $\frac{1}{f_{\mu_1}(x)\Phi_r(x)^a}$, we obtain the relation $\frac{F(x)}{G(x)} = \frac{-\alpha_{\mu_1}}{\Phi_r(x)^a}$, where $\Phi_r(x)^a$ does not divide $G(x)$. But this gives, in $\mathbb{Q}[x]$, the relation $F(x)\Phi_r(x)^a = -\alpha_{\mu_1}G(x)$ which contradicts uniqueness of factorisation in $\mathbb{Q}[x]$. Hence $\alpha_{\mu_1} = 0$, and by induction it follows that $\alpha_\mu = 0$ for every $\mu \in A_{p,n}$. We deduce that the set $\{f_\lambda(x) : \lambda \in A_{p,n}\}$ is a linearly independent subset of $\mathbb{Q}(x)$, and therefore that $B(p, n) \geq |A_{p,n}|$.

It is easy to see that

$$|A_{p,n}| = 1 + \sum_{\substack{1 < r \leq n, \\ (p,r)=1}} \lfloor \frac{n}{r} \rfloor.$$

Therefore, we have (where H_n denotes the n^{th} harmonic number)

$$|A_{p,n}| \geq 1 + \sum_{\substack{1 < d \leq n, \\ (p,d)=1}} \left(\frac{n}{d} - 1 \right) = 1 + n(H_n - 1 - \frac{1}{p}H_{\lfloor \frac{n}{p} \rfloor}).$$

It follows from [3, Theorem 1] that

$$|A_{p,n}| \geq 1 + \frac{p-1}{p}n \log n + n(\gamma - 1 - \frac{\gamma}{p}).$$

This gives the required result. \square

3.2. Wreath Products. We conclude by unifying our two extensions and considering the case of the Brauer characters of the generalised symmetric group. Let $B(p, n, k)$ denote the dimension of the space spanned by the Brauer characters of the symmetric powers of the natural representation of $C_k \wr S_n$ in characteristic p and let $\zeta = \exp(\frac{2\pi i}{k})$. An element $\sigma \in C_k \wr S_n$ has order coprime to p if and only if both $\lambda(\sigma) \in X_p$ and every t_i has order coprime to p . Let $R_{p,n,k}$ be the subset of $C_k \wr S_n$ consisting of elements of order coprime to p . The usual argument shows that $B(p, n, k) = \dim(\text{Span}_{\mathbb{C}}\{f_{\sigma}(x) : \sigma \in R_{p,n,k}\})$.

Proposition 15. *We have $B(p, n, k) \leq r_p(k)(\frac{p}{2p+2}n^2 + n(\log_p n + 1)) - n + 1$.*

Proof. Recall from the proof of Proposition 3 that a common denominator for the $f_{\sigma}(x)$ for $\sigma \in C_k \wr S_n$ is

$$D(x) = \prod_{\substack{0 \leq i < k \\ 1 \leq j \leq n}} (1 - \zeta^i x^j).$$

We say that we can cancel $p(x) \in \mathbb{C}[x]$ if $\frac{D(x)}{p(x)}$ is a common denominator for the $f_{\sigma}(x)$, where $\sigma \in R_{p,n,k}$. Our strategy is to cancel factors from $D(x)$. First, note that if p divides the order of ζ^i , then we may cancel $\prod_{j=1}^n (1 - \zeta^i x^j)$ from $D(x)$ by definition of $R_{p,n,k}$. Fix i such that ζ^i has order coprime to p . We need the following lemma.

Lemma 16. *If $\sigma \in R_{p,n,k}$ and $x - \theta$ divides $\frac{1}{f_\sigma(x)}$ in $\mathbb{C}[x]$, then θ has order coprime to p .*

Proof. Suppose that $x - \theta$ divides $\frac{1}{f_\sigma(x)}$ in $\mathbb{C}[x]$, then $\frac{1}{f_\sigma(\theta)} = 0$ and so $\prod_i (1 - t_i \theta^{\lambda_i}) = 0$. Hence there exist i, j such that $1 - \zeta^i \theta^{\lambda_j} = 0$. Since $\sigma \in R_{p,n,k}$, ζ^i has order coprime to p and λ_j is coprime to p . But the order of θ divides $\text{ord}(\zeta^i) \lambda_j$, and so is coprime to p . \square

Therefore, in a factorisation of $\prod_{j=1}^n (1 - \zeta^i x^j)$ into linear factors, we may cancel any factor $x - \theta$ where the order of θ is divisible by p . Consider the factor $1 - \zeta^i x^j$ where ζ^i has order d , and write $j = p^e r$ where p does not divide r . We claim that exactly r roots of the equation $1 - \zeta^i x^j = 0$ have order not divisible by p . Indeed, let η be a primitive j^{th} root of unity and let ν be a primitive $(dj)^{\text{th}}$ root of unity. Then the roots of $1 - \zeta^i x^j = 0$ form a $\langle \eta \rangle$ -coset in the group $\langle \nu \rangle$. Since a cyclic group of order $p^e r$ has exactly r elements of order coprime to p and $\langle \nu \rangle$ is abelian, this establishes our claim. It follows that we may cancel $j - r$ linear factors from the factorisation of $1 - \zeta^i x^j$, leaving a term of degree $r = r_p(j)$, as defined in Definition 9.

Consequently, cancelling factors from $\prod_{j=1}^n (1 - \zeta^i x^j)$, replaces $1 - \zeta^i x^j$ with a term of degree $r_p(j)$. This new polynomial has degree $\sum_{j=1}^n r_p(j)$, which is at most $\frac{p}{2p+2} n^2 + n(\log_p n + 1)$ by the proof of Proposition 11. We thus can put the power series $\{f_\sigma(x) : \sigma \in R_{p,n,k}\}$ over a common denominator of degree at most $r_p(k) (\frac{p}{2p+2} n^2 + n(\log_p n + 1))$, giving the desired result. \square

Acknowledgement. I am grateful to my advisor, Dr. Mark Wildon, for his many helpful comments and suggestions and to the referee for useful comments.

REFERENCES

- [1] DAVID SAVITT AND RICHARD P. STANLEY, ‘A note on the symmetric powers of the standard representation of S_n ’, *The Electronic Journal of Combinatorics* **7** (2000), R6
- [2] I. MARTIN ISAACS, ‘*Character Theory of Finite Groups*’, Dover Publications (1994)
- [3] BAI-NI GUO AND FENG QI, ‘Sharp bounds for harmonic numbers’, *Applied Mathematics and Computation* 218 (2011), no. 3, 991–995; Available online at <http://dx.doi.org/10.1016/j.amc.2011.01.089>
- [4] GORDON JAMES AND ADALBERT KERBER, ‘*The Representation Theory of the Symmetric Group*’, Cambridge University Press (1984)