

Exactly solvable interacting two-particle quantum graphs

Jens Bolte¹ and George Garforth²

Department of Mathematics
Royal Holloway, University of London
Egham, TW20 0EX
United Kingdom

Abstract

We construct models of exactly solvable two-particle quantum graphs with certain non-local two-particle interactions, establishing appropriate boundary conditions via suitable self-adjoint realisations of the two-particle Laplacian. Showing compatibility with the Bethe ansatz method, we calculate quantisation conditions in the form of secular equations from which the spectra can be deduced. We compare spectral statistics of some examples to well known results in random matrix theory, analysing the chaotic properties of their classical counterparts.

1 Introduction

In this paper we investigate the properties of two-particle quantum graphs with particular focus on the acquisition and analysis of their spectra. A quantum graph is a collection of vertices and edges of finite or infinite length equipped with a differential operator. The first theoretical model of a quantum graph was devised by Pauling [Pau36]. His motivation was to study the dynamics of free electrons in hydrocarbons by modelling carbon molecules as vertices and carbon-carbon bonds as edges. This idea was later adopted by Ruedenberg and Scherr [RS53] who used quantum graphs to describe free electrons donated by covalent bonds confined to entire quasi-one-dimensional molecules. Since then there have been multiple applications of quantum graphs in a variety of fields including quantum waveguides [FJK87], quantum chaos [KS97], quantum computation [Lov10] and mesoscopic systems [TM05]. For a review of quantum graphs, see [EKK⁺08, BK13a, ASV⁺16].

An important aspect of quantum graphs is that they may serve as models for quantum systems with corresponding complex classical dynamics. Kottos and Smilansky [KS97]

¹E-mail address: jens.bolte@rhul.ac.uk

²E-mail address: george.garforth.2012@live.rhul.ac.uk

demonstrated that eigenvalue correlations in quantum graphs can be described with random matrix models, therefore providing an example for the celebrated Bohigas-Giannoni-Schmit conjecture [BGS84] which is a central topic in quantum chaos. While the majority of quantum graphs literature is focussed on one-particle models, there have been a number of studies of many-particle quantum graphs. The first of these, by Melnikov and Pavlov [MP95], investigated the dynamics of two interacting particles on a connected graph with three infinite edges. Under certain restrictions of the system, they were able to find self-adjoint realisations of the two-body Laplacian corresponding to particle-particle and particle-vertex interactions. The resulting two-body wave function allowed the calculation of the conductivity of the system. More recently, Bolte and Kerner constructed two-particle quantum graphs with interactions localised at the vertices [BK13b] and later with singular contact interactions [BK13c]. Boundary conditions via suitable self-adjoint extensions of the two-particle Laplacian were ascertained using quadratic forms. These results were then used to study Bose-Einstein condensation [BK14].

To some extent, the success of one-particle quantum graph models relies upon the fact that their spectra are determined by a secular equation [KS97], i.e., quantum eigenvalues are given as zeros of a finite-dimensional determinant. This fact leads to very efficient methods to calculate eigenvalues, and also allows one to prove exact trace formulae for spectral densities [Rot83, KS97, BE09]. The reason behind this is the fact that, locally, the classical configuration space of a one-particle graph is one dimensional. For the quantum model this means that every eigenfunction must be a linear combination of left- and right-moving, one dimensional plane waves. Many-particle quantum graphs have higher dimensional classical configuration spaces, in general prohibiting a finite dimensional secular equation that determines the eigenvalues. This obstacle can be overcome under specific circumstances when symmetries lead to an exactly solvable model.

The first model of an exactly solvable many-body quantum system confined to a single dimension is due to Lieb and Liniger [LL63]. They determined the exact spectra of a repulsively δ -interacting Bose gas on a circle, a result which was later generalised to distinguishable particles by Yang [Yan67]. Gaudin [Gau71] later employed the Bethe ansatz to describe similar systems confined to an interval. These methods were formalised by the use of the Bethe ansatz, a sum of two-particle plane waves over possible particle configurations. Implicit in the use of the Bethe ansatz is the requirement for certain symmetries brought about by the interactions in the model. The consequence of increasing the complexity to systems of particles on two-edge graphs is that these symmetries are destroyed. By imposing certain non-local particle interactions, however, Caudrelier and Crampé [CC07] showed that compatibility with the Bethe ansatz is recovered. They were then able to calculate the exact spectra of these two-edge many-particle quantum graphs. Extending this method to general two-particle quantum graphs is the main aim of this paper.

In the next section we review the construction of general one-particle quantum graphs establishing appropriate boundary conditions by self-adjoint realisation of the one-particle Laplacian $-\Delta_1$. The boundary conditions lead to a quantisation condition from which the spectra can be calculated. In Section 3, we review the construction of two-particle quantum graphs introducing the notion of δ -interactions in terms of boundary conditions which char-

acterise a self-adjoint Laplacian $-\Delta_2$. We introduce the Bethe ansatz method for systems of two particles on a single edge, recovering historical results mentioned above. Systems of δ -interacting particles on graphs with more than a single edge are, in general, incompatible with the Bethe ansatz method. In Section 4, we construct exactly solvable systems of two particles on equilateral stars introducing the notion of $\tilde{\delta}$ -type interactions. These will be generalisations of the interactions imposed in [CC07] to ensure exact solvability in the two-edge setting. Establishing appropriate boundary conditions of the two-particle Laplacian, we prove exact solvability and calculate the spectra using the Bethe ansatz. In Section 5 we extend our argument from equilateral stars to general graphs. In Section 6 we analyse the spectral statistics of a number of examples of two-particle quantum graphs and comment on the nature of the corresponding classical dynamics.

2 One-particle quantum graphs

In this section we review [well known properties of one-particle quantum graphs and their spectra](#), see [EKK⁺08, BK13a, ASV⁺16]. A combinatorial, oriented graph $\Gamma(\mathcal{V}, \mathcal{I}, \mathcal{E}, f)$ is a collection of vertices $\mathcal{V} = \{v_1, \dots, v_{|\mathcal{V}|}\}$, connected by a set of internal edges $\mathcal{I} = \{i_1, \dots, i_{|\mathcal{I}|}\}$ and external edges $\mathcal{E} = \{e_1, \dots, e_{|\mathcal{E}|}\}$. The map f assigns to each external edge e_j a single vertex $f(e_j) = v_\eta$, and to each internal edge i_j an ordered pair of vertices $f(i_j) = (v_\gamma, v_\lambda)$ where $v_\gamma =: f_0(i_j)$ and $v_\lambda =: f_l(i_j)$ are initial and terminal vertices respectively. A pair of edges will be called distant if they have no common vertex and neighbouring if they have at least one common vertex. The set of distant and neighbouring edge couples will be denoted \mathcal{D} and \mathcal{N} , respectively. The degree d_η of a vertex $v_\eta \in \mathcal{V}$ is the number of edges connected to it. The combinatorial graph is turned into a metric graph by assigning a finite interval $[0, l_j]$ to each internal edge $i_j \in \mathcal{I}$ in such a way that $f_0(i_j)$ is identified with $x = 0$ and $f_l(i_j)$ with $x = l_j$. To each external edge $e_j \in \mathcal{E}$, a half-line $[0, \infty)$ is assigned such that $f(e_j)$ is identified with $x = 0$. A metric graph is called compact if there are no external edges, $\mathcal{E} = \emptyset$.

Let us consider the metric graph associated with $\Gamma = \Gamma(\mathcal{V}, \mathcal{I}, \mathcal{E}, f)$. The appropriate Hilbert space

$$\mathcal{H}_1 = \left(\bigoplus_{j=1}^{|\mathcal{I}|} L^2(0, l_j) \right) \oplus \left(\bigoplus_{j=|\mathcal{I}|+1}^{|\mathcal{I}|+|\mathcal{E}|} L^2(0, \infty) \right) \quad (2.1)$$

is the direct sum of the constituent Hilbert spaces on each edge. Vectors $\Psi = (\psi_j)_{j=1}^{|\mathcal{I}|+|\mathcal{E}|} \in \mathcal{H}_1$ are lists of square-integrable functions $\psi_j : (0, l_j) \rightarrow \mathbb{C}$, $j \in \{1, \dots, |\mathcal{I}|\}$ and $\psi_i : (0, \infty) \rightarrow \mathbb{C}$, $j \in \{|\mathcal{I}| + 1, \dots, |\mathcal{I}| + |\mathcal{E}|\}$. A quantum graph is a metric graph Γ with an associated Laplacian $-\Delta_1$ which acts according to

$$-\Delta_1 \Psi = \left(-\psi_j''(x) \right)_{j=1}^{|\mathcal{I}|+|\mathcal{E}|}, \quad (2.2)$$

where dashes denote ordinary, possibly weak, derivatives. In order for the Laplacian to be a one-particle quantum Hamiltonian it needs to be realised as a self-adjoint operator.

2.1 Self-adjoint realisation

One-particle observables on Γ are self-adjoint operators on \mathcal{H}_1 . We thus look for self-adjoint realisations of $-\Delta_1$ with domains characterised by boundary conditions at the vertices. To this end we define boundary vectors $\Psi_{bv}, \Psi'_{bv} \in \mathbb{C}^{2|\mathcal{I}|+|\mathcal{E}|}$ according to

$$\Psi_{bv} = \begin{pmatrix} (\psi_j(0))_{j=1}^{|\mathcal{I}|} \\ (\psi_j(l_j))_{j=1}^{|\mathcal{I}|} \\ (\psi_j(0))_{j=|\mathcal{I}|+1}^{|\mathcal{I}|+|\mathcal{E}|} \end{pmatrix} \quad \text{and} \quad \Psi'_{bv} = \begin{pmatrix} (\psi'_j(0))_{j=1}^{|\mathcal{I}|} \\ (-\psi'_j(l_j))_{j=1}^{|\mathcal{I}|} \\ (\psi'_j(0))_{j=|\mathcal{I}|+1}^{|\mathcal{I}|+|\mathcal{E}|} \end{pmatrix}. \quad (2.3)$$

Letting $H^2(\Gamma)$ be the set of all $\Psi \in \mathcal{H}_1$ such that $\psi_j \in H^2(0, l_j)$ for all $j \in \{1, \dots, |\mathcal{I}|\}$ and $\psi_j \in H^2(0, \infty)$ for all $j \in \{|\mathcal{I}| + 1, \dots, |\mathcal{I}| + |\mathcal{E}|\}$, we can state the following theorem [KS99].

Theorem 2.1. The Laplacian $-\Delta_1$ is self-adjoint on the set of all $\Psi \in H^2(\Gamma)$ which satisfy the boundary condition

$$A\Psi_{bv} + B\Psi'_{bv} = 0 \quad (2.4)$$

with $(2|\mathcal{I}|+|\mathcal{E}|) \times (2|\mathcal{I}|+|\mathcal{E}|)$ matrices A, B subject to the conditions $\text{rank}(A, B) = 2|\mathcal{I}|+|\mathcal{E}|$ and $AB^\dagger = BA^\dagger$.

2.2 Spectra of one-particle quantum graphs

The task is then to calculate the spectra of these quantum graphs by considering the eigenvalue equation

$$-\Delta_1\Psi = E\Psi \quad (2.5)$$

alongside boundary conditions prescribed by Theorem 2.1. We shall focus on compact graphs where the spectra are discrete.

The starting point is the observation that the components ψ_j of eigenfunctions $\Psi \in \mathcal{H}_1$ with real Laplace eigenvalues $E = k^2$ are necessarily of the form

$$\psi_j(x) = \alpha_j e^{ikx} + \beta_j e^{-ikx}, \quad (2.6)$$

where α_j and β_j are complex constants. Imposing boundary conditions (2.4) on functions $\Psi \in H^2(\Gamma)$, [one defines the scattering matrices](#)

$$S_v(k) = -(A + ikB)^{-1}(A - ikB) \quad (2.7)$$

and

$$T(k, \mathbf{l}) = \begin{pmatrix} 0 & e^{ik\mathbf{l}} \\ e^{ik\mathbf{l}} & 0 \end{pmatrix} \quad (2.8)$$

with

$$e^{ik\mathbf{l}} = \text{diag}(e^{ikl_j})_{j=1}^{|\mathcal{I}|}. \quad (2.9)$$

With the scattering matrices one can state the following theorem.

Theorem 2.2. [KS06] The non-zero eigenvalues of a self-adjoint Laplacian $-\Delta_1$ defined on Γ and with boundary conditions in the vertices specified through the matrices A, B are the values $E = k^2$ with multiplicity m , where $k \neq 0$ are solutions to the secular equation

$$\det [\mathbb{I}_{2|\mathcal{I}|} - S_v(k)T(k, \mathbf{l})] = 0 \quad (2.10)$$

with multiplicity m .

2.3 Star representation

We have seen that the spectrum of a compact quantum graph is given by the secular equation (2.10) which is a function of matrices $T(k, \mathbf{l})$ and $S_v(k)$. The former clearly contains the metric information. The latter contains information about the interactions at the vertices prescribed by A and B . In what follows we restrict our attention to *local* boundary conditions where boundary values of functions at different vertices are not related. The significance of this is that we can consider scattering at each vertex independently. We formalise this interpretation by dissecting the compact graph into a collection of star graphs with finitely many, infinite edges.

Definition 2.3. Consider a compact graph $\Gamma(\mathcal{V}, \mathcal{I}, f)$. Let the map g associate to each internal edge i_j an ordered pair of external edges $g(i_j) := (e_j, e_{j+|\mathcal{I}|})$. Here $e_j =: g_0(i_j)$ and $e_{j+|\mathcal{I}|} =: g_l(i_j)$ are external edges associated with initial and terminal vertices of i_j respectively so that $f(e_j) = f_0(i_j)$ and $f(e_{j+|\mathcal{I}|}) = f_l(i_j)$. The star representation of the compact graph Γ is the collection $\Gamma^{(s)}(\mathcal{V}, \mathcal{E}, f)$ of star graphs $\Gamma_\eta(v_\eta, \mathcal{E}_\eta, f)$ where \mathcal{E}_η is the set of edges e_j such that $f(e_j) = v_\eta$. Clearly we have that $2|\mathcal{I}| = |\mathcal{E}|$. The star graphs are turned into metric graphs by assigning half-lines $[0, \infty)$ to its edges.

Consider the star representation $\Gamma^{(s)}$ of a compact graph Γ . The Hilbert space associated with $\Gamma^{(s)}$ is

$$\mathcal{H}_1^{(s)} = \bigoplus_{j=1}^{|\mathcal{E}|} L^2(0, \infty), \quad (2.11)$$

and boundary values of vectors $\Psi = (\psi_j^{(s)})_{j=1}^{|\mathcal{E}|} \in \mathcal{H}_1^{(s)}$ are

$$\Psi_{\text{bv}}^{(s)} = \left(\psi_j^{(s)}(0) \right)_{j=1}^{|\mathcal{E}|} \quad \text{and} \quad \Psi_{\text{bv}}^{(s)'} = \left(\psi_j^{(s)'}(0) \right)_{j=1}^{|\mathcal{E}|}, \quad (2.12)$$

so that analogues of boundary conditions (2.4) are given by

$$A\Psi_{\text{bv}}^{(s)} + B\Psi_{\text{bv}}^{(s)'} = 0. \quad (2.13)$$

Let \mathbb{P} be an $|\mathcal{E}|$ -dimensional permutation matrix which reorders vectors Ψ according to

$$\mathbb{P}\Psi = (\Psi_\eta)_{\eta=1}^{|\mathcal{V}|}, \quad (2.14)$$

where each Ψ_η lists functions $\psi_j^{(s)}$ with $f(e_j) = v_\eta$. Local boundary conditions then imply the decomposition

$$A = \mathbb{P}^{-1} \left(\bigoplus_{v_\eta \in \mathcal{V}} A_\eta \right) \mathbb{P} \text{ and } B = \mathbb{P}^{-1} \left(\bigoplus_{v_\eta \in \mathcal{V}} B_\eta \right) \mathbb{P}, \quad (2.15)$$

where the matrices A_η, B_η characterise the boundary conditions in the vertices v_η of constituent star graphs Γ_η . The total scattering matrix can then be reconstructed according to

$$S_v(k) = \mathbb{P}^{-1} \left(\bigoplus_{\eta=1}^{|\mathcal{V}|} S_v^{(\eta)}(k) \right) \mathbb{P} \quad (2.16)$$

with

$$S_v^{(\eta)}(k) = -(A_\eta + ikB_\eta)^{-1}(A_\eta - ikB_\eta). \quad (2.17)$$

Let us retrieve the secular equation (2.10) by reconstructing the original compact graph from its star representation (see [KS97, KN05]). Consider the functions $\psi_j^{(s)}$ and $\psi_{j+|\mathcal{I}|}^{(s)}$ related to the external edges e_j and $e_{j+|\mathcal{I}|}$ respectively. Joining up the external edges to form a single internal edge of length l_j is imposing the relation

$$\psi_j^{(s)}(x) = \psi_{j+|\mathcal{I}|}^{(s)}(l_j - x). \quad (2.18)$$

Choosing the form (2.6) and defining vectors

$$\boldsymbol{\alpha} = (\alpha_j)_{j=1}^{|\mathcal{E}|} \text{ and } \boldsymbol{\beta} = (\beta_j)_{j=1}^{|\mathcal{E}|} \quad (2.19)$$

we have the relation

$$\boldsymbol{\beta} = T(k, \mathbf{l})\boldsymbol{\alpha}. \quad (2.20)$$

The total scattering matrix in this context acts according to

$$\boldsymbol{\alpha} = S_v(k)\boldsymbol{\beta}. \quad (2.21)$$

Applying (2.20) and (2.21) successively we recover the secular equation (2.10) as required.

3 Two-particle quantum graphs with δ -type interactions

The Hilbert space of a many-particle quantum system is given by the tensor product of one-particle Hilbert spaces. The appropriate two-particle Hilbert space for a compact two-particle quantum graph is

$$\mathcal{H}_2 = \left(\bigoplus_{j=1}^{|\mathcal{I}|} L^2(0, l_j) \right) \otimes \left(\bigoplus_{j=1}^{|\mathcal{I}|} L^2(0, l_j) \right) = \bigoplus_{(i_m, i_n) \in \mathcal{I} \times \mathcal{I}} L^2(D_{mn}), \quad (3.1)$$

where

$$D_{mn} = (0, l_m) \times (0, l_n). \quad (3.2)$$

The total classical configuration space for two particles on Γ is the union

$$D_\Gamma = \bigcup_{m,n=1}^{|\mathcal{I}|} D_{mn} \quad (3.3)$$

of rectangles. The two-particle Hilbert space can then be written $\mathcal{H}_2 = L^2(D_\Gamma)$. Vectors $\Psi = (\psi_{mn})_{m,n=1}^{|\mathcal{I}|}$ consist of functions $\psi_{mn} : D_{mn} \rightarrow \mathbb{C}$.

At this point let us introduce the two-particle Laplacian $-\Delta_2$ which acts according to

$$-\Delta_2 \Psi = \left(-\frac{\partial^2 \psi_{mn}}{\partial x_1^2} - \frac{\partial^2 \psi_{mn}}{\partial x_2^2} \right)_{m,n=1}^{|\mathcal{I}|}. \quad (3.4)$$

We wish to consider the two-particle eigenvalue equation

$$-\Delta_2 \Psi = E \Psi \quad (3.5)$$

alongside boundary conditions which prescribe single-particle interactions with vertices as well as singular contact interactions between particles. The latter interactions take place along the diagonals $x_1 = x_2$ of squares D_{mm} and are rigorous versions of δ -type interactions of the form

$$\alpha \delta(x_1 - x_2). \quad (3.6)$$

They naturally define the dissected configuration space

$$D_\Gamma^* = \left(\bigcup_{m,n=1|m \neq n}^{|\mathcal{I}|} D_{mn} \right) \cup \left(\bigcup_{m=1}^{|\mathcal{I}|} (D_{mm}^+ \cup D_{mm}^-) \right) \quad (3.7)$$

with subdomains of squares D_{mm} defined

$$D_{mm}^+ = \{(x_1, x_2) \in D_{mm}; x_1 > x_2\} \quad (3.8)$$

and

$$D_{mm}^- = \{(x_1, x_2) \in D_{mm}; x_1 < x_2\}. \quad (3.9)$$

The total dissected two-particle Hilbert space is then $\mathcal{H}_2^* = L^2(D_\Gamma^*)$. Thus two-particle wave functions $\Psi \in \mathcal{H}_2^*$ are lists

$$\Psi = \begin{pmatrix} (\psi_{mn})_{m,n=1|m \neq n}^{|\mathcal{I}|} \\ (\psi_{mm}^+)_{m=1}^{|\mathcal{I}|} \\ (\psi_{mm}^-)_{m=1}^{|\mathcal{I}|} \end{pmatrix} \quad (3.10)$$

of functions $\psi_{mn} : D_{mn} \rightarrow \mathbb{C}$, $m \neq n$, and $\psi_{mm}^\pm : D_{mm}^\pm \rightarrow \mathbb{C}$. The space $H^2(D_\Gamma^*)$ is defined in an analogous way.

The boundary conditions that one needs to impose on vectors $\Psi \in H^2(D_\Gamma^*)$ in order to generate single-particle interactions with the vertices as well as singular contact interactions (3.6) were established in [BK13c] and consist of two parts. The first set of boundary conditions is as described in Theorem 2.1 for each of the two variables separately. The boundary values of $\Psi \in H^2(D_\Gamma^*)$ at the vertices are

$$\Psi_{\text{bv}}^{(v)}(y) = \begin{pmatrix} (\psi_{mn}(0, l_n y))_{m,n=1}^{|\mathcal{I}|} \\ (\psi_{mn}(l_m, l_n y))_{m,n=1}^{|\mathcal{I}|} \\ (\psi_{mn}(l_m y, 0))_{n,m=1}^{|\mathcal{I}|} \\ (\psi_{mn}(l_m y, l_n))_{n,m=1}^{|\mathcal{I}|} \end{pmatrix} \quad \text{and} \quad \Psi_{\text{bv}}^{(v)'}(y) = \begin{pmatrix} (\psi_{mn,1}(0, l_n y))_{m,n=1}^{|\mathcal{I}|} \\ (\psi_{mn,1}(l_m, l_n y))_{m,n=1}^{|\mathcal{I}|} \\ (\psi_{mn,2}(l_m y, 0))_{n,m=1}^{|\mathcal{I}|} \\ (\psi_{mn,2}(l_m y, l_n))_{n,m=1}^{|\mathcal{I}|} \end{pmatrix} \quad (3.11)$$

for all $y \in (0, 1)$ where, for compactness the labels \pm are dropped. Boundary conditions at the vertices are then

$$(\mathbb{I}_2 \otimes A \otimes \mathbb{I}_{|\mathcal{I}|}) \Psi_{\text{bv}}^{(v)} + (\mathbb{I}_2 \otimes B \otimes \mathbb{I}_{|\mathcal{I}|}) \Psi_{\text{bv}}^{(v)'} = 0. \quad (3.12)$$

The δ -type interactions are characterised by the conditions

$$\psi_{mm}^+(x_1, x_2)_{x_1=x_2^+} = \psi_{mm}^-(x_1, x_2)_{x_1=x_2^-}; \quad (3.13)$$

$$\left(\frac{\partial}{\partial x_1} - \frac{\partial}{\partial x_2} - 2\alpha \right) \psi_{mm}^+(x_1, x_2)_{x_1=x_2^+} = \left(\frac{\partial}{\partial x_1} - \frac{\partial}{\partial x_2} \right) \psi_{mm}^-(x_1, x_2)_{x_1=x_2^-} \quad (3.14)$$

for every $x \in (0, l_m)$. Here, and in the remainder of this paper, we use the notation $x^\pm = \lim_{\delta \rightarrow 0} (x \pm \delta)$. In [BK13c] it was proven that the two-particle Laplacian with these boundary conditions is self-adjoint.

In Section 2, we calculated the spectra of one particle quantum graphs by specifying the form (2.6) of eigenfunctions of $-\Delta_1$ and applying boundary conditions (2.4). We would like to extend this approach to the two-particle quantum graph setting. As we are dealing with a two dimensional configuration space, the difficulty is that in general there does not exist a suitable analogue of the general form of an eigenfunction (2.6). It is well known, however, that in particular cases a Bethe ansatz can be used in this way.

Let us begin by considering systems of δ -interacting particles on the simplest metric graph, an interval $[0, l]$. Such systems were exactly solved in [Gau71, Yan67, LL63, Bet31] using the Bethe ansatz; a superposition of possible many-particle plane wave states as explicit eigenfunctions of the two-particle Laplacian. In this way exact spectra of many-particle systems can be calculated. To see how this works in the present setting we choose Dirichlet boundary conditions at the interval ends, $A = \mathbb{I}_2$ and $B = 0$. The Bethe ansatz method in this context is the assumption that the two components of eigenfunctions Ψ of $-\Delta_2$ take the form

$$\psi^\pm(x_1, x_2) = \sum_{P \in \mathcal{W}_2} \mathcal{A}^{(P, \pm)} e^{i(k_{P1}x_1 + k_{P2}x_2)}. \quad (3.15)$$

The amplitudes $\mathcal{A}^{(P,\pm)}$ are labelled by elements P of the Weyl group

$$\mathcal{W}_2 = (\mathbb{Z}/2\mathbb{Z})^2 \rtimes S_2 \quad (3.16)$$

of order 8, which is a semidirect product of its normal subgroup $(\mathbb{Z}/2\mathbb{Z})^2$ of order 4 with the symmetric group S_2 of order 2. This group is generated by the two elements R and T which act according to

1. $T(1, 2) = (2, 1)$;
2. $R(1, 2) = (-1, 2)$,

and satisfy the conditions

1. $TT = I$;
2. $RR = I$;
3. $TRTR = RTRT$.

The Bethe ansatz (3.15) can be seen to satisfy the eigenvalue equation (3.5) with Laplace eigenvalues $k_1^2 + k_2^2$.

We would first like to verify that the system is indeed exactly solvable, that is, boundary conditions (3.12)–(3.14) imposed on the ansatz (3.15) are compatible with these conditions. To this end let us define the 2-dimensional vector

$$\mathcal{A}^P = \begin{pmatrix} \mathcal{A}^{(P,-)} \\ \mathcal{A}^{(PT,+)} \end{pmatrix}. \quad (3.17)$$

The vertex boundary conditions (3.12) then imply

$$\mathcal{A}^{PR} = -\mathcal{A}^P \text{ and } \mathcal{A}^{PTRT} = -e^{2ik_{P2}l} \mathcal{A}^P \quad (3.18)$$

for all $P \in \mathcal{W}_2$. The δ -type interaction conditions (3.13) and (3.14) imply

$$\mathcal{A}^{PT} = S_p(k_{P1} - k_{P2}) \mathcal{A}^P \quad (3.19)$$

with

$$S_p(k) = \frac{1}{k + i\alpha} \begin{pmatrix} -i\alpha & k \\ k & -i\alpha \end{pmatrix}. \quad (3.20)$$

To prove exact solvability we need only show that relations (3.18) and (3.19) are consistent with the properties of \mathcal{W}_2 . This amounts to the requirements

1. $S_p(u)S_p(-u) = \mathbb{I}_2$,
2. $S_p(u)S_p(v) = S_p(v)S_p(u)$,

which are easily verified by the explicit form of $S_p(k)$. Now we have established that the system is exactly solvable, we would like to deduce the spectrum. Applying (3.18) and (3.19) successively, we arrive at the quantisation condition that

$$Z_{\text{interval}}(k_{P_1}, k_{P_2}) = 0 \quad (3.21)$$

with

$$Z_{\text{interval}}(k_1, k_2) = \det [\mathbb{I}_2 - e^{2ik_1l} S_p(k_1 - k_2) S_p(k_1 + k_2)] \quad (3.22)$$

is satisfied for all $P \in \mathcal{W}_2$. We note here that the form of $S_p(k)$ is such that if (3.21) is satisfied for some $P \in \mathcal{W}_2$, then it is necessarily satisfied for elements $PR, PTRT \in \mathcal{W}_2$. [This leads to the following result.](#)

Theorem 3.1. [\[Gau71\]](#) Non-zero eigenvalues of a self-adjoint two-particle Laplacian $-\Delta_2$ defined on an interval $[0, l]$ with Dirichlet interactions at the endpoints and δ -type particle interactions are the values $E = k_1^2 + k_2^2 \neq 0$ with multiplicity m , where (k_1, k_2) , such that $0 \leq k_1 \leq k_2$, are solutions to the secular equations

$$Z_{\text{interval}}(k_i, k_j) = 0 \quad (3.23)$$

for $j, i \neq j \in \{1, 2\}$ with multiplicity m .

We note that, by instead choosing boundary conditions

$$A = \begin{pmatrix} 1 & -1 \\ 0 & 0 \end{pmatrix} \text{ and } B = \begin{pmatrix} 0 & 0 \\ 1 & 1 \end{pmatrix}, \quad (3.24)$$

we recover the quantisation condition for two particles on a circle deduced in [\[Yan67\]](#). Furthermore, imposing bosonic symmetry

$$\psi^\pm(x_1, x_2) = \psi^\mp(x_1, x_2) \quad (3.25)$$

we recover the result in [\[LL63\]](#).

The aim of this paper is to calculate spectra for general two-particle graphs. It turns out that systems of δ -interacting particles on graphs with more than a single edge, in general, are not exactly solvable. The task is then to establish boundary conditions on general quantum graphs which are compatible with the Bethe ansatz method. In [\[CC07\]](#), systems of particles on two-edge stars, interacting via certain non-local δ -type interactions, were shown to be exactly solvable. In what follows, we extend this approach to systems of two particles on general graphs [in order to calculate their spectra](#). Furthermore, we show that the corresponding boundary conditions are compatible with a self-adjoint Laplacian.

4 Exactly solvable two-particle equilateral stars

Before discussing general graphs it is convenient to consider a subset of graphs called equilateral stars. These graphs exhibit most of the essential features of the general case and thus act as a convenient way to introduce some key concepts. Let us define the equilateral star Γ_e as the graph $\Gamma(\mathcal{V}, \mathcal{I}, f)$ with the restrictions $l_j = l$, $f_0(i_j) = v_1$ and $f_l(i_j) = v_{j+1}$ for all $i_j \in \mathcal{I}$. The degree d of the central vertex v_1 is then the same as the number of edges.

We would like to impose boundary conditions on vectors $\Psi \in \mathcal{H}_2$ which ensure exact solvability. We note here that the total configuration space D_{Γ_e} in the equilateral star setting is the union of square subdomains

$$D_{mn} = (0, l) \times (0, l). \quad (4.1)$$

Boundary conditions at the vertices are chosen in the same way as in Section 3. Interactions between particles will be analogues of the non-local δ -type interactions imposed in [CC07] and take place when particles are situated on neighbouring edges, the same distance from the common vertex of the edges. In what follows we refer to these as $\tilde{\delta}$ -type interactions. We reiterate that the set \mathcal{N} of neighbouring edges includes pairs (i_m, i_m) . On the equilateral star, such interactions are implemented by dissecting all squares D_{mn} along the lines $x_1 = x_2$ and thus define the dissected configuration space

$$D_{\Gamma_e}^* = \bigcup_{m,n=1}^{|\mathcal{I}|} (D_{mn}^+ \cup D_{mn}^-) \quad (4.2)$$

with D_{mn}^\pm defined in an analogous way to (3.8)-(3.9). Two-particle wave functions $\Psi \in L^2(D_{\Gamma_e}^*)$ are then lists

$$\Psi = \left(\begin{array}{c} (\psi_{mn}^+)_{m=1}^{|\mathcal{I}|} \\ (\psi_{mn}^-)_{m=1}^{|\mathcal{I}|} \end{array} \right). \quad (4.3)$$

The $\tilde{\delta}$ -type interactions are characterised by the conditions

$$\psi_{mn}^+(x_1, x_2)|_{x_1=x_2^+} = \psi_{nm}^-(x_1, x_2)|_{x_1=x_2^-}; \quad (4.4)$$

$$\left(\frac{\partial}{\partial x_1} - \frac{\partial}{\partial x_2} - 2\alpha \right) \psi_{mn}^+(x_1, x_2)|_{x_1=x_2^+} = \left(\frac{\partial}{\partial x_1} - \frac{\partial}{\partial x_2} \right) \psi_{nm}^-(x_1, x_2)|_{x_1=x_2^-} \quad (4.5)$$

for almost every $x_i \in (0, l)$.

Using the method devised in [BK13c] one can introduce vectors of boundary values that reproduce the boundary conditions (4.4)–(4.5) and prove that they provide a self-adjoint realisation of the two-particle Laplacian.

We stress here that $\tilde{\delta}$ -type interactions can take place when particles are located on different edges and therefore represent rather less physical interactions than δ -type interactions. We choose these models as they permit exact solutions via the Bethe ansatz

form

$$\psi_{mn}^{\pm}(x_1, x_2) = \sum_{P \in \mathcal{W}_2} \mathcal{A}_{mn}^{(P, \pm)} e^{i(k_{P1}x_1 + k_{P2}x_2)} \quad (4.6)$$

of the components in (4.3). This form obviously leads to eigenfunctions of (3.5) with eigenvalues $E = k_1^2 + k_2^2$.

To show that the boundary conditions imposed on the ansatz (4.6) are compatible with the properties of \mathcal{W}_2 , let us first define the $d^2 \times d^2$ permutation matrix

$$\mathbb{T}_{d^2} = \begin{pmatrix} \mathbb{I}_d \otimes M_1 \\ \vdots \\ \mathbb{I}_d \otimes M_d \end{pmatrix} \quad (4.7)$$

with row vectors

$$M_j = \left(\underbrace{0 \dots 0}_{j-1} \quad 1 \quad \underbrace{0 \dots 0}_{d-j} \right). \quad (4.8)$$

It is convenient to note the properties

$$\mathbb{T}_{d^2} (\mathcal{A}_{mn}^P)_{m,n=1}^d = (\mathcal{A}_{mn}^P)_{n,m=1}^d \quad (4.9)$$

and

$$\mathbb{T}_{d^2} (M \otimes N) \mathbb{T}_{d^2} = N \otimes M \quad (4.10)$$

for any $d \times d$ matrices M and N . Then, defining the $2d^2$ -dimensional vector

$$\mathcal{A}^P = \begin{pmatrix} (\mathcal{A}_{mn}^{(P,-)})_{m,n=1}^d \\ \mathbb{T}_{d^2} (\mathcal{A}_{mn}^{(PT,+)})_{m,n=1}^d \end{pmatrix}, \quad (4.11)$$

the boundary conditions (3.12) imply

$$\begin{aligned} & (\mathbb{I}_2 \otimes A \otimes \mathbb{I}_d) \mathbb{Q} \begin{pmatrix} \mathcal{A}^P + \mathcal{A}^{PR} \\ \mathcal{A}^{PT} e^{ik_{P1}l} + \mathcal{A}^{PRT} e^{-ik_{P1}l} \end{pmatrix} \\ & + ik_{P1} (\mathbb{I}_2 \otimes B \otimes \mathbb{I}_d) \mathbb{Q} \begin{pmatrix} \mathcal{A}^P - \mathcal{A}^{PR} \\ -\mathcal{A}^{PT} e^{ik_{P1}l} + \mathcal{A}^{PRT} e^{-ik_{P1}l} \end{pmatrix} = 0 \end{aligned} \quad (4.12)$$

for all $P \in \mathcal{W}_2$ where

$$\mathbb{Q} = \begin{pmatrix} \mathbb{I}_{d^2} & 0 & 0 & 0 \\ 0 & 0 & 0 & \mathbb{T}_{d^2} \\ 0 & \mathbb{I}_{d^2} & 0 & 0 \\ 0 & 0 & \mathbb{T}_{d^2} & 0 \end{pmatrix}. \quad (4.13)$$

Equilateral stars have Dirichlet conditions at external vertices v_j , $j \geq 2$. We thus have the decomposition

$$A = \begin{pmatrix} A_1 & 0 \\ 0 & A_2 \end{pmatrix} \text{ and } B = \begin{pmatrix} B_1 & 0 \\ 0 & B_2 \end{pmatrix}, \quad (4.14)$$

where A_1, B_1 refer to the boundary conditions at the central vertex, and

$$A_2 = \mathbb{I}_d \text{ and } B_2 = 0 \quad (4.15)$$

provide the Dirichlet conditions at the external vertices. By using the properties of \mathbb{T}_{d^2} , we then have that

$$\mathbb{Q}^{-1} (\mathbb{I}_2 \otimes S_v(k) \otimes \mathbb{I}_d) \mathbb{Q} = \begin{pmatrix} \mathbb{I}_2 \otimes S_v^{(1)}(k) \otimes \mathbb{I}_d & 0 \\ 0 & -\mathbb{I}_{2d^2} \end{pmatrix}. \quad (4.16)$$

Rearranging (4.12), we can then extract the relation

$$\mathcal{A}^{PR} = (\mathbb{I}_2 \otimes S_v^{(1)}(-k_{P1}) \otimes \mathbb{I}_d) \mathcal{A}^P. \quad (4.17)$$

The $\tilde{\delta}$ -conditions (4.4)–(4.5) imply

$$\mathcal{A}^{PT} = (S_p(k_{P1} - k_{P2}) \otimes \mathbb{I}_{d^2}) \mathcal{A}^P \quad (4.18)$$

with $S_p(k)$ defined in (3.20). To prove exact solvability we need only show that relations (4.17)–(4.18) are consistent with the properties of \mathcal{W}_2 . This amounts to the requirements

1. $S_v^{(1)}(u)S_v^{(1)}(-u) = \mathbb{I}_d$;
2. $S_p(u)S_p(-u) = \mathbb{I}_2$;
3. $\left(\mathbb{I}_2 \otimes S_v^{(1)}(u) \otimes \mathbb{I}_d \right) (S_p(u+v) \otimes \mathbb{I}_{d^2}) \left(\mathbb{I}_2 \otimes S_v^{(1)}(v) \otimes \mathbb{I}_d \right) (S_p(v-u) \otimes \mathbb{I}_{d^2})$
 $= (S_p(v-u) \otimes \mathbb{I}_{d^2}) \left(\mathbb{I}_2 \otimes S_v^{(1)}(v) \otimes \mathbb{I}_d \right) (S_p(u+v) \otimes \mathbb{I}_{d^2}) \left(\mathbb{I}_2 \otimes S_v^{(1)}(u) \otimes \mathbb{I}_d \right).$

The first two conditions are easily verified by the explicit forms of $S_v^{(1)}(u)$ and $S_p(u)$. The third follows from the result in [KS06] that, for any A, B and u, v , we have the commutation relation

$$[S_v(u), S_v(v)] = 0. \quad (4.19)$$

Now we have established that the system is exactly solvable, we would like to deduce the spectrum. This can be done in a number of ways. The method we choose here generalises that used for the one-particle case in [KS06] which we presented in Section 2.2. Substituting (4.18) into (4.12) we have that

$$((\mathbb{I}_2 \otimes A \otimes \mathbb{I}_d) \mathbb{Q}X(k_{P1}, k_{P2}, l) + ik_{P1} (\mathbb{I}_2 \otimes B \otimes \mathbb{I}_d) \mathbb{Q}Y(k_{P1}, k_{P2}, l)) \begin{pmatrix} \mathcal{A}^P \\ \mathcal{A}^{PR} \end{pmatrix} = 0 \quad (4.20)$$

with

$$X(k_1, k_2, l) = \begin{pmatrix} \mathbb{I}_2 & \\ S_p(k_1 - k_2)e^{ik_1l} & S_p(-k_1 - k_2)e^{-ik_1l} \end{pmatrix} \otimes \mathbb{I}_{d^2} \quad (4.21)$$

and

$$Y(k_1, k_2, l) = \begin{pmatrix} \mathbb{I}_2 & \\ -S_p(k_1 - k_2)e^{ik_1l} & S_p(-k_1 - k_2)e^{-ik_1l} \end{pmatrix} \otimes \mathbb{I}_{d^2}. \quad (4.22)$$

Then by the properties of determinants, and since $\det(A + ikB) \neq 0$, we arrive at the condition that

$$Z_e(k_{P_1}, k_{P_2}) = 0 \quad (4.23)$$

with

$$Z_e(k_1, k_2) = \det [\mathbb{I}_{2d} + e^{2ik_1l} (S_p(k_1 - k_2)S_p(k_1 + k_2) \otimes S_v^{(1)}(k_1))] = 0 \quad (4.24)$$

is satisfied for all $P \in \mathcal{W}_2$. By using properties of determinants and the explicit forms of $S_p(k)$ and $S_v(k)$, it is easy to see that the form (4.24) is such that if (4.23) is satisfied for some $P \in \mathcal{W}_2$, then it is necessarily satisfied for elements $PR, PTRT \in \mathcal{W}_2$. With this in mind, we can state the main result of this section.

Theorem 4.1. Non-zero eigenvalues of a self-adjoint two-particle Laplacian $-\Delta_2$ defined on an equilateral star Γ_e with interactions at the central vertex specified through A_1, B_1 and $\tilde{\delta}$ -type particle interactions are the values $E = k_1^2 + k_2^2 \neq 0$ with multiplicity m , where (k_1, k_2) such that $0 \leq k_1 \leq k_2$, are solutions to the secular equations

$$Z_e(k_i, k_j) = 0 \quad (4.25)$$

for $j, i \neq j \in \{1, 2\}$ with multiplicity m .

Before we move on to general graphs, let us establish agreement with some results discussed earlier in the paper.

Firstly, choosing $d = 2$, we recover the spectrum of a system of two particles on an interval with a central impurity as solved in [CC07]. Furthermore, rather than choosing Dirichlet vertex conditions (4.15), which specified the connectivity of an equilateral star, and instead choosing Kirchhoff boundary conditions

$$A_2 = \begin{pmatrix} 1 & -1 \\ 0 & 0 \end{pmatrix} \text{ and } B_2 = \begin{pmatrix} 0 & 0 \\ 1 & 1 \end{pmatrix} \quad (4.26)$$

to establish continuity at the outer vertices, we recover the spectra of systems of two particles on a circle with an impurity also in [CC07].

Throughout this section we have used α to parameterise the strength of particle interactions. It is reasonable to expect then that by setting $\alpha = 0$, one should arrive at separable

quantisation conditions given by (2.10) for one-particle quantum graphs. Indeed, by substituting $\alpha = 0$ into the form (4.24), we recover

$$\det [\mathbb{I}_d + e^{2ikl} S_v^{(1)}(k)] = 0 \quad (4.27)$$

which is exactly the corresponding one-particle condition. It is important to point out however, that $\tilde{\delta}$ -type interactions with $\alpha = 0$ result in coupling between domains D_{mn} and D_{nm} and are thus clearly distinct from the truly non-interacting situation. For this reason we refer to such systems as *pseudo-non-interacting*. The fact the one-particle condition (4.27) is recovered in this case is a result of the specific geometry of the equilateral star. We will see in the subsequent section that this agreement does not hold for general graphs. We revisit this point in the final section of the chapter when discussing spectral statistics.

5 Exactly solvable two-particle quantum graphs

We have seen, in the previous section, how to construct exactly solvable models of two interacting particles on equilateral stars. The majority of quantum graphs literature, however, is concerned with the dynamics of single particles on graphs with, in general, different edge lengths. Indeed, in [KS97], rationally independent edge lengths are required to avoid degenerate energy levels and ensure spectral statistics following random matrix predictions. This section is concerned with extending the scope of our discussion to two particles on general compact graphs.

Again we would like to impose boundary conditions on vectors $\Psi \in \mathcal{H}_2$ which ensure exact solvability. As imposing $\tilde{\delta}$ -type interactions between particles on equilateral stars leads to exact solutions, it seems reasonable to assume that a suitable variant of such interactions in the general setting will also lead to exact solutions. However, general graphs bring added complications associated with edges of different length and distant vertices. The problem is then to choose an appropriate way to impose $\tilde{\delta}$ -type interactions in the general setting which preserves compatibility with the Bethe ansatz method. In what follows we impose that $\tilde{\delta}$ -type interactions become effective when two particles on neighbouring edges are located the same distance from their common vertex. This section is concerned with characterising these interactions in terms of self-adjoint realisations of the Laplacian and then, viewing the graph in its star representation, calculating the two-particle spectra. Let us begin by considering a pair of particles on a neighbouring edge couple $(i_m, i_n) \in \mathcal{N}$ with coordinates $x_1 \in [0, l_m]$ and $x_2 \in [0, l_n]$ respectively. The orientations of the edges could be such that $f_0(i_m) = f_0(i_n)$, $f_0(i_m) = f_l(i_n)$, $f_l(i_m) = f_0(i_n)$ or $f_l(i_m) = f_l(i_n)$. The $\tilde{\delta}$ -type interactions prescribed above in the star representation become effective when either $x_1 = x_2$, $x_1 = l_n - x_2$, $l_m - x_1 = x_2$ or $l_m - x_1 = l_n - x_2$, respectively. The interaction will then be cut off at the smaller of the two edge lengths involved. The four cases can be

mapped to the first one by changing coordinates according to

$$(\tilde{x}_1, \tilde{x}_2) = \begin{cases} (x_1, x_2) & \text{if } f_0(i_m) = f_0(i_n), \\ (x_1, l_n - x_2) & \text{if } f_0(i_m) = f_l(i_n), \\ (l_m - x_1, x_2) & \text{if } f_l(i_m) = f_0(i_n), \\ (l_m - x_1, l_n - x_2) & \text{if } f_l(i_m) = f_l(i_n). \end{cases} \quad (5.1)$$

In these coordinates interactions take place when $\tilde{x}_1 = \tilde{x}_2$. The total dissected configuration space can be written

$$D_\Gamma^* = \left(\bigcup_{(i_m, i_n) \in \mathcal{D}} D_{mn} \right) \cup \left(\bigcup_{(i_m, i_n) \in \mathcal{N}} (\tilde{D}_{mn}^+ \cup \tilde{D}_{mn}^-) \right) \quad (5.2)$$

with subdomains defined as

$$\tilde{D}_{mn}^+ = \{(\tilde{x}_1, \tilde{x}_2) \in (0, l_m) \times (0, l_n); \tilde{x}_1 > \tilde{x}_2\} \quad (5.3)$$

and

$$\tilde{D}_{mn}^- = \{(\tilde{x}_1, \tilde{x}_2) \in (0, l_m) \times (0, l_n); \tilde{x}_1 < \tilde{x}_2\}. \quad (5.4)$$

In order to define the quantum model, very much in analogy to (3.10)–(3.14), we need the Hilbert space $L^2(D_\Gamma^*)$ and a domain that is specified as a subspace of $H^2(D_\Gamma^*)$ in terms of boundary conditions. We denote the components of vectors $\Psi \in H^2(D_\Gamma^*)$ as $\psi_{mn} : D_{mn} \rightarrow \mathbb{C}$, when $(i_m, i_n) \in \mathcal{D}$ and $\psi_{mn}^\pm : D_{mn}^\pm \rightarrow \mathbb{C}$, when $(i_m, i_n) \in \mathcal{N}$. First defining vertex boundary vectors as in (3.11) we again have vertex conditions (3.12). With the definition $l_{mn}^- = \min(l_m, l_n)$ we have $\tilde{\delta}$ -type boundary conditions

$$\psi_{mn}^+(\tilde{x}_1, \tilde{x}_2)|_{\tilde{x}_1=\tilde{x}_2^+} = \psi_{nm}^-(\tilde{x}_1, \tilde{x}_2)|_{\tilde{x}_1=\tilde{x}_2^-}; \quad (5.5)$$

$$\left(\frac{\partial}{\partial \tilde{x}_1} - \frac{\partial}{\partial \tilde{x}_2} - 2\alpha \right) \psi_{mn}^+(\tilde{x}_1, \tilde{x}_2)|_{\tilde{x}_1=\tilde{x}_2^+} = \left(\frac{\partial}{\partial \tilde{x}_1} - \frac{\partial}{\partial \tilde{x}_2} \right) \psi_{nm}^-(\tilde{x}_1, \tilde{x}_2)|_{\tilde{x}_1=\tilde{x}_2^-} \quad (5.6)$$

for all $\tilde{x}_1, \tilde{x}_2 \in (0, l_{mn}^-)$ when $(i_m, i_n) \in \mathcal{N}$.

As in the equilateral star case, using the method devised in [BK13c] one can introduce vectors of boundary values that reproduce the boundary conditions (5.5)–(5.6) and prove that they provide a self-adjoint realisation of the two-particle Laplacian.

5.1 Spectra

We have seen how to establish boundary conditions which correspond to two-particle quantum graphs with $\tilde{\delta}$ -type interactions. We would now like to show that such systems are exactly solvable and calculate their spectra.

For the equilateral star graphs considered in Section 4, exact solvability was shown by substituting the ansatz (4.6) directly into boundary conditions (3.12) and (4.4)–(4.5)

defined on $D_{\Gamma_e}^*$. The spectra then followed by generalising the approach in [KS06] to two particles. While, in principle, we can use the same method in the general graph case, the extra complexity brought about by different edge lengths makes the presentation rather convoluted. To this end we will use the method presented in Section 2.3 which utilises the star representation $\Gamma^{(s)}$ of a compact graph Γ . In the compact setting, $\tilde{\delta}$ -type interactions require us to define dissections (5.3)-(5.4) of domains D_{mn} which relate to neighbouring pairs (i_m, i_n) of edges on Γ . On $\Gamma^{(s)}$, this corresponds to defining dissections of

$$D_{mn}^{(s)} = (0, \infty) \times (0, \infty), \quad (5.7)$$

with $f(e_m) = f(e_n)$, according to

$$D_{mn}^{(s,+)} = \{(x_1, x_2) \in D_{mn}^{(s)}; x_1 > x_2\}, \quad (5.8)$$

and

$$D_{mn}^{(s,-)} = \{(x_1, x_2) \in D_{mn}^{(s)}; x_1 < x_2\}. \quad (5.9)$$

In order to simplify the formalism which follows, it is convenient to artificially extend the dissections (5.8)–(5.9) to all edge pairs so that the total dissected configuration space is given by

$$D_{\Gamma^{(s)}}^* = \bigcup_{m,n=1}^{|\mathcal{E}|} (D_{mn}^{(s,+)} \cup D_{mn}^{(s,-)}). \quad (5.10)$$

The appropriate Hilbert space is then $\mathcal{H}_2^{(s,*)} = L^2(D_{\Gamma^{(s)}}^*)$ so that two-particle wave functions $\Psi^{(s)} \in \mathcal{H}_2^{(s,*)}$ are lists

$$\Psi^{(s)} = \left(\begin{array}{c} (\psi_{mn}^{(s,+)}|_{\mathcal{E}})_{m,n=1} \\ (\psi_{mn}^{(s,-)}|_{\mathcal{E}})_{m,n=1} \end{array} \right) \quad (5.11)$$

of functions $\psi_{mn}^{(s,\pm)} : D_{mn}^{(s,\pm)} \rightarrow \mathbb{C}$. We note that later in the formalism we correct for these extra dissections by imposing appropriate conditions of continuity (see Figure 1).

The task is now to specify the boundary conditions we would like to impose. Interactions at the vertices in this setting will be described by simple two-particle lifts of (2.13) and are exact analogues of (3.11)–(3.12). The $\tilde{\delta}$ -type interactions are implemented through the conditions

$$\psi_{mn}^{(s,+)}(x_1, x_2)|_{x_1=x_2^+} = \psi_{nm}^{(s,-)}(x_1, x_2)|_{x_1=x_2^-}; \quad (5.12)$$

$$\left(\frac{\partial}{\partial x_1} - \frac{\partial}{\partial x_2} - 2\alpha \right) \psi_{mn}^{(s,+)}(x_1, x_2)|_{x_1=x_2^+} = \left(\frac{\partial}{\partial x_1} - \frac{\partial}{\partial x_2} \right) \psi_{nm}^{(s,-)}(x_1, x_2)|_{x_1=x_2^-}. \quad (5.13)$$




Figure 1: Appropriate dissections in the star representation of an internal edge couple (i_m, i_n) with $f_0(i_m) = f_0(i_n)$. $\tilde{\delta}$ -type interactions are imposed along the solid diagonal. Continuity is imposed across dashed diagonals.

on edge pairs (e_m, e_n) where $f(e_m) = f(e_n)$. Continuity is reestablished in domains $D_{mn}^{(s)}$, with $f(e_m) \neq f(e_n)$, by setting

$$\psi_{mn}^{(s,+)}(x_1, x_2)|_{x_1=x_2^+} = \psi_{mn}^{(s,-)}(x_1, x_2)|_{x_1=x_2^-}; \quad (5.14)$$

$$\left(\frac{\partial}{\partial x_1} - \frac{\partial}{\partial x_2}\right) \psi_{mn}^{(s,+)}(x_1, x_2)|_{x_1=x_2^+} = \left(\frac{\partial}{\partial x_1} - \frac{\partial}{\partial x_2}\right) \psi_{mn}^{(s,-)}(x_1, x_2)|_{x_1=x_2^-}, \quad (5.15)$$

that is, for these edge pairs we require the functions and their derivatives to be continuous across diagonals.

Using the Bethe ansatz method, an eigenvector $\Psi^{(s)}$ in the star representation will be described by the collection of functions

$$\psi_{mn}^{(s,\pm)}(x_1, x_2) = \sum_{P \in \mathcal{W}_2} \mathcal{A}_{mn}^{(P,\pm)} e^{i(k_{P1}x_1 + k_{P2}x_2)} \quad (5.16)$$

on $D_{mn}^{(s,\pm)}$ (for all edge pairs). Let us define the $2|\mathcal{E}|^2$ -dimensional vector

$$\mathcal{A}^P = \begin{pmatrix} (\mathcal{A}_{mn}^{(P,-)})_{m,n=1}^{|\mathcal{E}|} \\ \mathbb{T}_{|\mathcal{E}|^2} (\mathcal{A}_{mn}^{(PT,+)})_{m,n=1}^{|\mathcal{E}|} \end{pmatrix}, \quad (5.17)$$

with \mathbb{T} as defined in (4.7). The vertex-boundary conditions then imply

$$\mathcal{A}^{PR} = (\mathbb{I}_2 \otimes S_v(-k_{P1}) \otimes \mathbb{I}_{|\mathcal{E}|}) \mathcal{A}^P \quad (5.18)$$

for all $P \in \mathcal{W}_2$. At this point, it is convenient to define the matrix $\mathbf{c} = \text{diag}(c_{mn})_{m,n=1}^{|\mathcal{E}|}$ where

$$c_{mn} = \begin{cases} 1 & \text{if } f(e_m) = f(e_n); \\ 0 & \text{else,} \end{cases} \quad (5.19)$$

which distinguishes domains with $\tilde{\delta}$ -type interactions from those where derivatives are continuous across dissections. The $\tilde{\delta}$ -conditions then imply

$$\mathcal{A}^{PT} = Y(k_{P_1} - k_{P_2})\mathcal{A}^P \quad (5.20)$$

with

$$Y(k) = S_p(k) \otimes \mathbf{c} + \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \otimes (\mathbb{I}_{|\mathcal{E}|^2} - \mathbf{c})\mathbb{T}_{|\mathcal{E}|^2}. \quad (5.21)$$

To prove exact solvability we need only show that relations (5.18) and (5.20) are consistent with the properties of \mathcal{W}_2 . This amounts to the requirements

1. $S_v(u)S_v(-u) = \mathbb{I}_{|\mathcal{E}|}$;
2. $Y(k)Y(-k) = \mathbb{I}_{2|\mathcal{E}|}$;
3. $(\mathbb{I}_2 \otimes S_v(u) \otimes \mathbb{I}_{|\mathcal{E}|}) Y(u+v) (\mathbb{I}_2 \otimes S_v(v) \otimes \mathbb{I}_{|\mathcal{E}|}) Y(v-u) = Y(v-u) (\mathbb{I}_2 \otimes S_v(v) \otimes \mathbb{I}_{|\mathcal{E}|}) Y(u+v) (\mathbb{I}_2 \otimes S_v(u) \otimes \mathbb{I}_{|\mathcal{E}|})$.

The first two conditions are easily verified by the explicit forms of $S_v(u)$ and $Y(u)$, noting that, since $c_{mn} = c_{nm}$, the properties of $\mathbb{T}_{|\mathcal{E}|^2}$ are such that

$$[\mathbf{c}, \mathbb{T}_{|\mathcal{E}|^2}] = 0. \quad (5.22)$$

Noting then the relation

$$[S_v(u) \otimes \mathbb{I}_{|\mathcal{E}|}, \mathbf{c}] = 0 \quad (5.23)$$

holds if vertex boundary conditions are local, that is $S_v(u)$ obeys the condition (2.15), and also the relation (4.19), the third condition is easily verified.

Let us bring our attention back to the original compact graph Γ . In order to turn the eigenfunctions in the star representation into eigenfunctions on the compact graph, it is sufficient to impose the relations

$$\psi_{mn}^{(s,+)}(x_1, x_2) = \psi_{(m+|\mathcal{I}|)n}^{(s,+)}(l_m - x_1, x_2) \quad \text{and} \quad \psi_{mn}^{(s,-)}(x_1, x_2) = \psi_{m(n+|\mathcal{I}|)}^{(s,-)}(x_1, l_n - x_2) \quad (5.24)$$

for all $m, n \in \{1, \dots, |\mathcal{I}|\}$ which imply

$$\mathcal{A}_{mn}^{(P,+)} = \mathcal{A}_{(m+|\mathcal{I}|)n}^{(PR,+)} e^{-ik_{P_1}l_m} \quad \text{and} \quad \mathcal{A}_{mn}^{(P,-)} = \mathcal{A}_{m(n+|\mathcal{I}|)}^{(PTRT,-)} e^{-ik_{P_2}l_n}. \quad (5.25)$$

These conditions then yield the relation

$$\mathcal{A}^P = E(-k_{P2})\mathcal{A}^{PTRT} \quad (5.26)$$

where

$$E(k) = \mathbb{I}_{4|Z|} \otimes \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \otimes e^{ikl} \quad (5.27)$$

with e^{ikl} defined as in (2.9). Applying (5.18), (5.20) and (5.26) successively we have the condition that

$$Z(k_{P1}, k_{P2}) = 0 \quad (5.28)$$

with

$$Z(k_1, k_2) = \det [\mathbb{I}_{8|Z|^2} - E(k_2)Y(k_2 - k_1) (\mathbb{I}_2 \otimes S_v(k_2) \otimes \mathbb{I}_{2|Z|}) Y(k_1 + k_2)] \quad (5.29)$$

is satisfied for all $P \in \mathcal{W}_2$. By using properties of determinants, the commutation relations established above and the explicit forms of $Y(k)$, $S_v(k)$ and $E(k)$, it is easy to see that the form (5.29) is such that if (5.28) is satisfied for some $P \in \mathcal{W}_2$, then it is necessarily satisfied for elements $PR, PTRT \in \mathcal{W}_2$. With this in mind we can state the main result of this section.

Theorem 5.1. Non-zero eigenvalues of a self-adjoint two-particle Laplacian $-\Delta_2$ defined on Γ with local vertex interactions specified through A, B and $\tilde{\delta}$ -type interactions between particles when they are located on neighbouring edges are the values $E = k_1^2 + k_2^2 \neq 0$ with multiplicity m , where (k_1, k_2) , such that $0 \leq k_1 \leq k_2$, are solutions to the secular equations

$$Z(k_i, k_j) = 0 \quad (5.30)$$

for $j, i \neq j \in \{1, 2\}$ with multiplicity m .

5.2 Recovering specific results

To finish the section, we establish agreement between the spectra of general two-particle quantum graphs presented in Theorem 5.1 and results derived and discussed earlier in the paper.

The above result matches our previous result for equilateral stars given by Theorem 4.1 by choosing the same boundary conditions at the vertices and setting all edge lengths to $l_m = l$. As, on star graphs, $\tilde{\delta}$ -type interactions are defined between all pairs of edges, one has to put

$$c_{mn} = \begin{cases} 1 & \text{if } f(e_m) = f(e_n) = v_1 \text{ or } f(e_m), f(e_n) \in \{v_2, \dots, v_{|V|}\}; \\ 0 & \text{else.} \end{cases} \quad (5.31)$$

Substituting these parameters into $Z(k_i, k_j)$ and using the properties of determinants we recover the form $Z_e(k_i, k_j)$ as required.

In Section 4, we introduced the notion of pseudo-non-interacting particles and showed that in the equilateral star setting, the appropriate quantisation condition is indeed that of the truly non-interacting case. However, in the general setting, this agreement does not hold. The spectra of such systems is calculated first by identifying the matrix

$$\lim_{\alpha \rightarrow 0} Y(k) = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \otimes (\mathbf{c} + (\mathbb{I}_{|\mathcal{E}|^2} - \mathbf{c})\mathbb{T}_{|\mathcal{E}|^2}). \quad (5.32)$$

Substitution into (5.29) then yields the quantisation condition

$$\begin{aligned} Z(k) = \det & \left[\mathbb{I}_{4|Z|^2} - (S_v(k) \otimes \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \otimes e^{ikl}) \mathbf{c} \right. \\ & \left. - (\mathbb{I}_{2|Z|} \otimes (\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \otimes e^{ikl}) S_v(k)) (\mathbb{I}_{4|Z|^2} - \mathbf{c}) \right] \end{aligned} \quad (5.33)$$

which we notice is dependent on the single momentum k .

Truly non-interacting systems are recovered by turning off all coupling between domains D_{mn} and D_{nm} . This is achieved by setting $\mathbf{c} = \mathbf{0}$. We then have that

$$Y(k)|_{\mathbf{c}=\mathbf{0}} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \otimes \mathbb{T}_{|\mathcal{E}|^2} \quad (5.34)$$

By substituting into (5.29) we recover the secular equation (2.10) for the one-particle quantum graph.

In the subsequent section we would like to analyse examples of bosons on graphs. Computationally speaking, such examples are useful as the dimension of the matrix inside the determinant $Z(k_1, k_2)$ is halved. Imposing bosonic symmetry we have

$$\psi_{mn}^{(s,\pm)}(x_1, x_2) = \psi_{nm}^{(s,\mp)}(x_2, x_1) \quad (5.35)$$

for all $(x_1, x_2) \in D_{mn}^{(s)}$ which implies the relations

$$\mathcal{A}^{(P,-)} = \mathcal{A}^{(PT,+)} \quad (5.36)$$

for all $P \in \mathcal{W}_2$. The matrix (5.21) then reduces to the form

$$Y_b(k) = \mathbb{I}_2 \otimes \left(\frac{k - i\alpha}{k + i\alpha} \mathbf{c} + (\mathbb{I}_{|\mathcal{E}|^2} - \mathbf{c})\mathbb{T}_{|\mathcal{E}|^2} \right). \quad (5.37)$$

so that from (5.29), we recover

$$Z_b(k_1, k_2) = \det \left[\mathbb{I}_{4|Z|^2} - E_b(k_2) Y_b(k_2 - k_1) (S_v(k_2) \otimes \mathbb{I}_{2|Z|}) Y_b(k_1 + k_2) \right] \quad (5.38)$$

with $E(k) = \mathbb{I}_2 \otimes E_b(k)$.

6 Spectral Statistics

In this section we calculate the spectra of certain examples of two-particle quantum graphs and analyse their statistics.

Let us first introduce the eigenvalue counting function

$$N(E) = \#\{n; E_n \leq E\}. \quad (6.1)$$

It was shown in [BK13c] that for two-particle quantum graphs with δ -type interactions, the asymptotic behaviour follows the Weyl law

$$N(E) \sim \frac{\mathcal{L}^2}{4\pi} E, \quad E \rightarrow \infty, \quad (6.2)$$

where $\mathcal{L} = \sum_{j=1}^{|\mathcal{I}|} l_j$ denotes the total length of the graph. Of course, the majority of this paper is not concerned with δ -type interactions, but with $\tilde{\delta}$ -type interactions. Nonetheless, it is still revealing to compare the eigenvalue count with (6.2). To this end, we will assign a line of best fit

$$\bar{N}(E) = aE + b\sqrt{E} + c \quad (6.3)$$

to the counting function and compare the leading term to (6.2).

One of the main motivations for the study of quantum graphs is to analyse their spectral correlations. A particularly useful statistical measure is the nearest neighbour level spacings distribution

$$\int_a^b p(s) ds = \lim_{N \rightarrow \infty} \frac{1}{N} \#\{n \leq N; a \leq \epsilon_{n+1} - \epsilon_n \leq b\}. \quad (6.4)$$

of the unfolded versions, $\epsilon_1 < \epsilon_2 < \epsilon_3 < \dots$, of the energy eigenvalues; that is the energies are rescaled such that the average spacing is equal to unity. Generic quantum systems with integrable classical limits are conjectured to have spectra with Poissonian statistics [BT77],

$$p(s) = e^{-s}, \quad (6.5)$$

while chaotic classical systems have quantum counterparts with correlations described by random matrix models. For systems with integer spin and time-reversal symmetry Gaussian orthogonal ensemble (GOE) statistics are conjectured to apply [BGS84], where the level spacings distribution can be approximated by

$$p(s) = \frac{\pi}{2} s e^{-\frac{\pi}{4}s^2}, \quad (6.6)$$

see [Haa91]. In this section we analyse the spectral statistics of certain examples by using the integrated measure (6.4).

In [KS97], nearest neighbour energy level distributions of one-particle quantum tetrahedra were shown to exhibit GOE spectral statistics and thus imply chaotic classical counterparts. In this section we analyse the spectra of two-particle quantum graphs calculated in the previous sections, looking for a potential dependence of spectral correlations on the interaction strength. We refer to the result by Srivastava et al. [STL⁺16] who analysed the spectral properties of interacting kicked rotors which individually show GOE statistics. They found a transition from Poissonian to GOE statistics as the strength of the interaction was increased.

6.1 Tetrahedron

Let us take, as an example, a system of two $\tilde{\delta}$ -interacting bosons on a tetrahedron. The appropriate spectra are calculated according to Theorem 5.1 using the secular equation (5.38). Vertex boundary conditions are determined by choosing Discrete Fourier Transform (DFT) scattering matrices $S_v^{(\eta, DFT)}$ with elements

$$(S_v^{(\eta, DFT)})_{\gamma\lambda} = \frac{1}{\sqrt{d_\eta}} e^{2\pi i \frac{n(\gamma)n(\lambda)}{d_\eta}}, \quad (6.7)$$

where $n(\cdot)$ is a bijection of the d_η neighbouring vertices of v_η onto the numbers $\{0, \dots, d_\eta - 1\}$. For the tetrahedron, appropriate DFT scattering matrices at each vertex v_η are then

$$S_\eta^{(DFT)} = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 1 & 1 \\ 1 & e^{\frac{2i\pi}{3}} & e^{\frac{4i\pi}{3}} \\ 1 & e^{\frac{4i\pi}{3}} & e^{\frac{8i\pi}{3}} \end{pmatrix}, \quad (6.8)$$

with distinct eigenvalues $\{-1, 1, i\}$. With this choice, the spectrum of the two-particle **tetrahedron** with $\tilde{\delta}$ -type interactions **and rationally independent edge lengths** is non-degenerate.

Before analysing two-particle spectra, let us consider the spectra of non-interacting systems. Figure 2 plots the nearest neighbour distributions for the single-particle spectra associated with truly non-interacting ($\mathbf{c} = \mathbf{0}$) and pseudo-non-interacting ($\alpha = 0$) particles on the tetrahedron with DFT scattering matrices. As is well-known [KS97] and confirmed in Figure 2, the one-particle spectrum follows GOE statistics. The pseudo-non-interacting system, however, shows Poissonian statistics. The crucial point here is that two-particle systems prescribed in Theorem 5.1 in fact couple systems of pseudo-non-interacting particles which individually possess spectra with Poissonian statistics, not systems of truly non-interacting particles which individually follow GOE statistics. Thus we cannot expect a transition to GOE statistics as in [STL⁺16]. Figure 3 plots the α -dependency of the lowest energy levels of a system of $\tilde{\delta}$ -interacting bosons on a tetrahedron with DFT vertex scattering matrices. There is no obvious transition to a regime of energy level repulsion as we increase α . Indeed, plots of nearest neighbour distributions reveal Poissonian statistics for all interaction strengths. Figure 4 shows these plots for interaction strengths $\alpha = 1$ and $\alpha = 10$. Figure 5 plots counting functions $N(E)$ for strengths $\alpha \in \{0, 1, 10\}$ together with quadratic lines of best fit (6.3). In each case, the leading term does not agree with the Weyl law (6.2) predicted for contact interactions.




Figure 2: Integrated level spacings distributions for systems of non-interacting and pseudo-non-interacting particles on a tetrahedron. First 50,000 eigenvalues.




Figure 3: Dependency on interaction strength of small eigenvalues of a system of two bosons on a tetrahedron.




Figure 4: Integrated level spacings distributions for systems of two bosons on a tetrahedron. First 3000 eigenvalues.




Figure 5: Counting functions $N(E)$ (solid line) with lines of best fit $\bar{N}(E)$ (dashed line) for systems of two bosons on a tetrahedron.

6.2 Equilateral star

To examine the spectral statistics of coupled chaotic systems we must look for two-particle quantum graphs for which the one-particle spectra recovered when setting $\alpha = 0$ are chaotic. We have seen that the two-particle tetrahedron does not fulfil this requirement. Let us then focus our attention on equilateral stars which we discussed in Section 4, where we showed that we recover true one-particle spectra when setting $\alpha = 0$. Thus we can discuss coupled chaotic systems in the spirit of [STL⁺16] if we can find one-particle equilateral stars which exhibit GOE statistics. Such systems are characterised by the quantisation condition (4.27) which can be written

$$e^{-2ikl} = -\mu(k), \quad (6.9)$$

where $\mu(k)$ is an eigenvalue of $S_v^{(1)}(k)$. Clearly, the multiplicity of solutions k are equal to the multiplicity these eigenvalues. For example, equilateral stars with boundary conditions characterised by the DFT scattering matrix (6.7) at the central vertex yield solutions corresponding to $\mu = \{1, -1, i, -i\}$ with degenerate values arising for $d > 3$. Clearly degenerate energy levels would obscure conclusions made in the context of spectral statistics. To navigate this issue, we must choose a scattering matrix with non-degenerate eigenvalues. In what follows, we choose a $d_{|\mathcal{I}|} \times d_{|\mathcal{I}|}$ random unitary matrix. Figure 6 plots the nearest neighbour distribution for a single particle on such an equilateral star with 9 edges. The degenerate energy level spacings arise from the imposition of equal lengths. Indeed in studies of one-particle quantum graph spectra, rationally independent lengths are chosen to avoid degenerate level spacings. We do however see approximate agreement with GOE




Figure 6: Integrated level spacings distributions for systems of non-interacting and pseudo-non-interacting particles on a 9-edge equilateral star. First 50,000 eigenvalues.




Figure 7: Dependency on interaction strength of small eigenvalues of a system of two bosons on a 9-edge equilateral star.




Figure 8: Integrated level spacings distributions for systems of two bosons on a 9-edge equilateral star. First 100 eigenvalues.




Figure 9: Dependency on interaction strength of large eigenvalues of a system of two bosons on a 9-edge equilateral star.




Figure 10: Integrated level spacings distributions for systems of two bosons on a 9-edge equilateral star with $\alpha = 10$.

statistics. In this setting we can thus investigate the coupling of two chaotic spectra by increasing α from 0.

Figure 7 plots the α -dependency of the lowest energy levels of a system of two bosons on a 9-edge equilateral star with a random unitary central scattering matrix. We clearly see a transition to level repulsion as α is increased. Figure 8 plots nearest neighbour distributions for the first 100 energy levels. There is a clear shift from Poissonian, for $\alpha = 1$, to GOE statistics, for $\alpha = 10$. We note, however, that this level repulsion becomes less apparent as we include larger energy levels; Figure 9 shows level crossing at higher energies and Figure 10 shows how the spectral statistics for the $\alpha = 10$ case tend to Poissonian as we include higher energies. Figure 11 plots counting functions for $\alpha \in \{0, 1, 10\}$ together with quadratic lines of best fit (6.3). For the non-interacting ($\alpha = 0$) case, the leading term agrees exactly with the Weyl law (6.2). As the interaction strength increases, the counting function diverges from this prediction.

References

- [ASV⁺16] F. M. Andrade, A. G. M. Schmidt, E. Vicentini, B. K. Cheng, and M. G. E. da Luz, *Green's function approach for quantum graphs: an overview*, Phys. Rep. **647** (2016), 1–46.
- [BE09] J. Bolte and S. Endres, *The trace formula for quantum graphs with general self-adjoint boundary conditions*, Ann. H. Poincaré **10** (2009), 189–223.
- [Bet31] H. Bethe, *Zur Theorie der Metalle. i. Eigenwerte und Eigenfunktionen der linearen Atomkette (German) [On the theory of metals. i. Eigenvalues and eigenfunctions of a linear chain of atoms]*, Z. Phys. **71** (1931), 205–226.
- [BGS84] O. Bohigas, M. J. Giannoni, and C. Schmit, *Characterization of chaotic quantum spectra and universality of level fluctuation laws*, Phys. Rev. Lett. **52** (1984), 1–4.




Figure 11: Counting functions $N(E)$ (solid line) with lines of best fit $\bar{N}(E)$ (dashed line) for systems of two bosons on a 9-edge equilateral star.

- [BK13a] G. Berkolaiko and P. Kuchment, *Introduction to quantum graphs*, Mathematical Surveys and Monographs, vol. 186, AMS, 2013.
- [BK13b] J. Bolte and J. Kerner, *Quantum graphs with singular two-particle interactions*, J. Phys. A: Math. Theor. **46** (2013), 045206.
- [BK13c] ———, *Quantum graphs with two-particle contact interactions*, J. Phys. A: Math. Theor. **46** (2013), 045207.
- [BK14] ———, *Many-particle quantum graphs and bose-einstein condensation*, J. Math. Phys. **55** (2014), 061901.
- [BT77] M. V. Berry and M. Tabor, *Level clustering in the regular spectrum*, Proc. R. Soc. London. Series A, Math. and Phys. Sci. **356** (1977), 375–394.
- [CC07] V. Caudrelier and N. Crampé, *Exact results for the one-dimensional many-body problem with contact interaction: Including a tunable impurity*, Rev. Math. Phys. **19** (2007), 349–370.
- [EKK⁺08] P. Exner, J. P. Keating, P. Kuchment, T. Sunada, and A. Teplyaev (eds.), *Analysis on graphs and its applications*, Proc. Symp. Pure Math., vol. 77, AMS, 2008, Papers from the program held in Cambridge, January 8–June 29, 2007.
- [FJK87] C. Flesia, R. Johnston, and H. Kunz, *Strong localization of classical waves: a numerical study*, Europhys. Lett. **3** (1987), 497–502.
- [Gau71] M. Gaudin, *Boundary energy of a Bose gas in one dimension*, Phys. Rev. A. **4** (1971), 386–394.

- [Haa91] F. Haake, *Quantum Signatures of Chaos*, Springer Series in Synergetics, vol. 54, Springer-Verlag, Berlin, 1991.
- [KN05] P. Kurasov and M. Nowaczyk, *Inverse spectral problem for quantum graphs*, J. Phys. A.: Math. Gen. **38** (2005), 49014915.
- [KS97] T. Kottos and U. Smilansky, *Quantum chaos on graphs*, Phys. Rev. Lett. **79** (1997), 4794–4797.
- [KS99] V. Kostrykin and R. Schrader, *Kirchhoff's rule for quantum wires*, J. Phys. A: Math. Gen. **32** (1999), 595–630.
- [KS06] V. Kostrykin and R. Schrader, *Laplacians on metric graphs: Eigenvalues, resolvents and semigroups*, Contemp. Math. **415** (2006), 201–225.
- [LL63] E. H. Lieb and W. Liniger, *Exact analysis of an interacting Bose gas. I. The general solution and the ground state*, Phys. Rev. (2) **130** (1963), 1605–1616.
- [Lov10] N. B. Lovett, *Universal quantum computation using the discrete time quantum walk*, Phys. Rev. A. **81** (2010), 042330.
- [MP95] Yu. B. Melnikov and B. S. Pavlov, *Two-body scattering on a graph and application to simple nanoelectronic devices*, J. Math. Phys. **36** (1995), 2813–2825.
- [Pau36] L. Pauling, *The diamagnetic anisotropy of aromatic molecules*, J. Chem. Phys. **4** (1936), 673–677.
- [Rot83] J.-P. Roth, *Spectre du laplacien sur un graphe (French) [The spectra of Laplacians on graphs]*, C. R. Acad. Sci. Paris Sér. I Math. **296** (1983), 793–795.
- [RS53] K. Ruedenberg and C. W. Scherr, *Free-electron network model for conjugated systems. I. Theory*, J. Chem. Phys. **21** (1953), 1565–1581.
- [STL⁺16] Sh. Srivastava, S. Tomsovic, A. Lakshminarayan, R. Ketzmerick, and A. Bäcker, *Universal scaling of spectral fluctuation transitions for interacting chaotic systems*, Phys. Rev. Lett. **116** (2016), 054101.
- [TM05] C. Texier and G. Montambaux, *Quantum oscillations in mesoscopic rings and anomalous diffusion*, J. Phys. A.: Math. Gen. **38** (2005), 3455–3471.
- [Yan67] C.N. Yang, *Some exact results for the many-body problem in one dimension with repulsive delta-function interaction*, Phys. Rev. Lett. **19** (1967), 1312–1315.