

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38

## The search for clues to abiogenesis on Mars

Joseph R. Michalski, University of Hong Kong  
Tullis C. Onstott, Princeton University  
Stephen J. Mojzsis, University of Colorado  
John Mustard, Brown University  
Queenie Chan, NASA Johnson Space Center  
Paul B. Niles, NASA Johnson Space Center  
Sarah Stewart Johnson, Georgetown University

### [First Paragraph]

Few traces of Earth’s geologic record are preserved from the time of life’s emergence, over 3800 million years ago. Consequently, what little we understand about abiogenesis—the origin of life on Earth—is based primarily on laboratory experiments and theory. The best geological lens for understanding the early Earth might actually come from Mars, a planet with a crust that’s overall far more ancient than our own. On Earth, surface sedimentary environments are thought to best preserve evidence of ancient life, but this is mostly because our planet has been dominated by high photosynthetic biomass production at the surface for the last ~2500 million years or more. By the time oxygenic photosynthesis evolved on Earth, Mars had been a hyperarid, frozen desert with a surface bombarded by high-energy solar and cosmic radiation for more than a billion years, and as a result, photosynthetic surface life may never have occurred on Mars. Therefore, one must question whether searching for evidence of life in martian surface sediments is the best strategy. This paper explores the possibility that the abundant hydrothermal environments on Mars might provide more valuable insights into life’s origins.

### [main text]

Following planetary accretion, early delivery via impact of extraterrestrial materials and their payload of volatiles and organic matter may have provided a vast amount of exogenous raw ingredients for abiogenesis<sup>1</sup>. Although the details of the post-accretionary impact period termed the “Late Heavy Bombardment<sup>2</sup>” are intensely debated, consensus is that large impacts were relatively common in the early inner solar system. The catastrophic effects of the impact events would have been a major impediment to the formation, evolution, and preservation of early life, particularly surface life<sup>3</sup>. Yet a mere 800 Myr after the Earth-Moon formed, at the time that the impact rate seems to have diminished, some manner of microbial life appears to have existed<sup>4</sup>.

39

40 Hints of early life on Earth are found as isotopically light graphitized carbon captured in  
41 metamorphic rocks representing the ancient seafloor in what is now Canada<sup>5</sup> and Greenland<sup>6</sup>,  
42 and conical stromatolite-like structures within slightly younger rocks<sup>7</sup>. Cryptic evidence in the  
43 form of graphite trapped in zircons from the Jack Hills region could push life's origins even  
44 earlier into the Hadean eon<sup>8</sup>. These remnants of the Archean and Hadean eons that are so  
45 relevant to understanding the temporal and taphonomic window for life's emergence  
46 comprise only about ~0.001 vol.% of the terrestrial crust<sup>9</sup> and have been intensely thermally  
47 and chemically altered due to their long crustal residence times.

48

49 Because the Earth's early geologic record is so poorly preserved, our limited understanding of  
50 how early organic chemistry may have assembled the building blocks of life is largely based  
51 upon laboratory experiments<sup>10-15</sup>. But definitive clues to the chemical steps leading to life's  
52 origins probably require empirical evidence. The fundamental question of how abiogenesis  
53 occurred on Earth may only be answerable through finding better preserved "cradle of life"  
54 chemical systems beyond Earth. Indeed, this question of how life originates is one of the  
55 fundamental drivers of international space exploration.

56

57 Which objects beyond Earth could potentially unlock the mystery of abiogenesis? Europa and  
58 Enceladus are high priority targets because they likely contain subsurface oceans even  
59 today<sup>16,17</sup>. Yet, it is not simply a subsurface ocean itself that is intriguing in terms of the  
60 origin of life perspective—it is the reaction between fluids and silicate rocks at the ocean-  
61 silicate interface<sup>18</sup> that might hold promise for energetic pathways for chemotrophic life  
62 forms<sup>19</sup>. However, all of the icy satellites are far from Earth and access to subsurface fluids  
63 will either require deep drilling, or we will be limited to collection of ejected molecules from  
64 transient cryo-volcanism<sup>20,21</sup>, which may not sample fluids from the deep rock-water interface  
65 of primary interest. In addition, there is a growing interest in the possibility that terrestrial life  
66 originated not within an ocean environment but rather in vapour-dominated inland geothermal  
67 systems, where shallow pools of fluid may have interacted with porous silicate minerals and  
68 metal sulphides<sup>22,23</sup>. While the icy worlds are clearly a high-priority target for understanding  
69 abiogenesis, Mars is the only Solar System object with an ancient, preserved, accessible crust  
70 containing clear evidence of water-rock reactions dating to the time when life appeared on  
71 Earth (Figure 1).

72

### 73 **Mars as a Rosetta Stone for Early Earth**

74 Mars, a planet without plate tectonics and with much lower weathering rates through most of  
75 its history<sup>24</sup>, contains a much older and better-preserved geologic record than the Earth

76 (Figure 2). At only ~10% of Earth's mass, Mars began with far less primordial and radiogenic  
77 heat<sup>25</sup>. By about 4000 Mya, Mars had cooled sufficiently to cause cessation of the magnetic  
78 dynamo<sup>26</sup>. The loss of the martian magnetic field marked the timing of its clearest divergence  
79 from the evolution of the Earth and its biosphere. It exposed the martian surface to punishing  
80 radiation<sup>27</sup>, and the atmosphere began to be sputtered away by solar wind<sup>28</sup>.

81

82 Mars may have been cold, arid, oxidizing and generally inhospitable at the surface for much  
83 of its history, however hydrothermal conditions in the near surface or subsurface might have  
84 been considerably more clement. Infrared remote sensing has revealed the presence of  
85 thousands of deposits of hydrated silicate minerals as well as various salts throughout the  
86 martian surface<sup>29</sup>. While essentially all of the salts and some of the hydrated silicates  
87 seemingly formed in surface environments during what may have been short lived climate  
88 excursions<sup>30</sup>, many of the deposits represent materials that were seemingly exhumed from the  
89 subsurface<sup>31</sup>. Among the exhumed phases are serpentines, Fe and Mg-rich smectite clays,  
90 chlorites, carbonates, and amorphous silica that seemingly indicate widespread subsurface  
91 hydrothermal alteration (Figure 3).

92

93 In 2008, the Spirit Rover also stumbled upon soils of nearly pure opaline silica (>90 wt %  
94 SiO<sub>2</sub>) in the vicinity of Home Plate in Gusev crater<sup>32</sup> providing compelling evidence for  
95 fumarolic hydrothermal activity. Similar materials were also detected in at least one younger  
96 caldera, that on Nili Patera<sup>33</sup>.

97

98 It is difficult to strongly constrain the timing of near surface and deep subsurface  
99 hydrothermal alteration other than to state that it was primarily in the Noachian (>3600  
100 Mya)<sup>31</sup>. Whereas some studies document impact-induced hydrothermal activity in Hesperian  
101 crater deposits (3000-3600 Mya) e.g.<sup>34</sup>, there is little to no evidence for similar alteration in  
102 craters formed in Amazonian age terranes<sup>35</sup> (<3000 Mya) (Figure 3).

103

104 The subsurface – from metres to kilometres depth - is potentially the largest, longest-lived,  
105 and most stable habitable environment on Mars<sup>36</sup>. A significant fraction of Earth's biomass  
106 consists of prokaryotic microbial life in a deep biosphere<sup>37</sup>, a habitat that was essentially  
107 disregarded more than 30 years ago and remains largely unexplored today<sup>38</sup>. The primary  
108 producers in deep subsurface ecosystems are anaerobic chemoautotrophs, or SLiMEs, that  
109 oxidize H<sub>2</sub> and reduce CO<sub>2</sub> to produce CH<sub>4</sub> (i.e. methanogens) and acetate (i.e. acetogens)<sup>39,40</sup>.  
110 The life-sustaining H<sub>2</sub> has been shown to be generated through abiotic hydrolysis of ferrous  
111 minerals in basalt<sup>41</sup> and ultramafic rock (e.g. serpentinization)<sup>42</sup> and through radiolysis of  
112 water<sup>49</sup>. Other potential sources of H<sub>2</sub> include exsolved gases from basaltic magmas,

113 decomposition of CH<sub>4</sub> at T > 600°C, reactions between CH<sub>4</sub>, H<sub>2</sub>O and CO<sub>2</sub> at elevated  
114 temperatures and silicate cataclasis. Just as important, radiolysis has been shown to generate  
115 electron acceptors such as sulphate along with H<sub>2</sub>, which can be utilized to sustain sulphate  
116 reducing bacteria indefinitely<sup>43</sup>.

117

118 On Earth, the extent of the deep biosphere is controlled not only by energy sources and  
119 nutrients, but by availability of pore space. While porosity is strongly dependent on rock type,  
120 continental rocks typically have <1-5% porosity at depth of 3-4 km. But due to the lower  
121 gravity on Mars the rocks are less compacted; similar values of porosity extend to ~10 km in  
122 depth<sup>44</sup>.

123

124 Although heat flow in the terrestrial crust is heterogeneous, geothermal gradients (10-40  
125 K/km) in continental and oceanic crustal settings suggest that the terrestrial deep biosphere  
126 likely does not extend passed ~3-4 km depth, beyond which the most tolerant thermophiles  
127 are no longer viable (~120°C)<sup>41</sup>. However, on Mars the lower surface temperature and lower  
128 crustal heat flow add up to a more favourable thermal regime within the crust. Assuming a  
129 thermal gradient of 20 K/km on Noachian Mars, the 120°C temperature limit would not have  
130 been reached until nearly twice the depth where it occurs on Earth (Figure 4).

131

132 Most of the martian crust is ultramafic or mafic, and likely contains interlayered volcanics  
133 and impactites. Given the lower temperature gradient on Mars compared to Earth, it is likely  
134 that Lost City-type<sup>45</sup> (low-temperature, alkaline) serpentinization reactions<sup>19</sup> occurred over a  
135 large range of depths on Mars, producing bioavailable H<sub>2</sub><sup>46</sup>. Although this mafic-rich crust is  
136 less radiogenic than average Earth continental crust, H<sub>2</sub> production rates from radiolysis  
137 should be as great as that for subsurface environments on the Earth because of the greater  
138 porosity of the martian subsurface<sup>47</sup>. Exhumed subsurface carbonates, and the presence of  
139 vein carbonates in martian meteorites exhumed from the subsurface<sup>48</sup> suggest that these  
140 reactions happened in the presence of CO<sub>2</sub> and may have produced abiogenic hydrocarbons.  
141 Consequently, the subsurface habitable volume and abiotic energy sources would likely have  
142 been as readily available, if not more so, on Mars as on Earth.

143

144 It is probable that fluids within alkaline crustal hydrothermal systems would have mixed with  
145 descending acidic, sulphur- (H<sub>2</sub>S, SO<sub>2</sub>) and CO<sub>2</sub>-rich fluids from surface and near-surface  
146 environments through taliks, areas of unfrozen ground surrounded by permafrost<sup>44</sup>. Likewise,  
147 alkaline fluids might have emerged in deep basins and interfaced with acidic lakes and  
148 meltwater from acidic ice deposits, resulting in mixing scenarios which may have been a

149 source of redox energy<sup>19,49</sup>. A test for such an origin of life scenario would be invaluable to  
150 earth and planetary scientists alike.

151

### 152 **Dim prospects for surface life on Mars**

153 The evolutionary innovation of oxygenic photosynthesis by cyanobacteria was a turning point  
154 in the history of life on Earth<sup>50</sup>. Although the timing remains controversial, oxygenic  
155 photosynthesis appeared late within cyanobacterial evolution<sup>51</sup>, well after their divergence at  
156 2.5 to 2.6 billion years<sup>52</sup> and after the rise of its evolutionary precursor, Mn-oxidizing  
157 phototrophy<sup>53</sup> and before the Great Oxidation Event at 2.3-2.45 Ga<sup>54,55</sup>. Production of  
158 atmospheric O<sub>2</sub> led to the formation of ozone, which shielded the immediate surface zone  
159 from harmful UV rays. The success of cyanobacteria not only led to marked increases in  
160 biomass production and deposition in shallow water environments (shelf, coastal marine, and  
161 lacustrine) where high sedimentary rates prevail, but also to the colonization of arid and cold  
162 surface environments by endolithic communities<sup>56</sup>. Our paleontologic record over the last  
163 ~3000 Mya is dominated by carbonaceous sedimentary rocks from such environments<sup>57</sup>.

164

165 On Mars, there may have never been an evolutionary drive to inhabit the surface. During the  
166 Noachian, Mars was most likely cold, arid and oxidizing<sup>58,59</sup>. Fluvial channels and most crater  
167 lakes on Noachian Mars once thought to have required some form of greenhouse atmosphere  
168 in order to stabilize liquid water over geologic time scales, are now considered by some to  
169 have formed within thousands of years<sup>60,61</sup>, perhaps under a tenuous atmosphere in the  
170 Noachian. The surface seemingly shifted from a cold but episodically wet landscape to a  
171 frozen, hyperarid desert at the Noachian-Hesperian transition ca. 3600 Mya<sup>62</sup>.

172

173 The success of surface life on Earth can be traced back to the evolution of oxygenic  
174 photosynthesis in the Archean. The most recent molecular clocks have placed the origin of  
175 photosynthesis at ~3000 Myr and the origin of oxygenic photosynthesis later than 2500 to  
176 2600 Mya on Earth. Martian phototrophs would have had to attain these evolutionary  
177 benchmarks by 3600 Mya, despite the generally frozen and arid surface conditions, fainter  
178 sunlight, and the intense radiation flux from solar UV, Solar Energetic Particles (SEPs) and  
179 Galactic Cosmic Rays (GCRs). By contrast, the evolution of methanogenesis, an important  
180 metabolic pathway for subsurface life, occurred prior to the divergence of Euryarcheota and  
181 Crenarcheota and represents one of the most ancient forms of metabolism<sup>63</sup>.

182

183 Considering these challenges, it seems prudent to consider the possibility that photosynthesis  
184 never evolved on Mars. Unless high-energy radiation could be harnessed as a form of energy,  
185 as has been reported for certain fungal species<sup>64</sup>, the radiated surface environment is an

186 impediment to the existence of surface life and an obstacle for the preservation of organic  
187 materials<sup>36</sup>. With all this in mind, it seems time to reconsider the current Mars exploration  
188 philosophy.

189

### 190 **A Mars exploration strategy focused on abiogenesis**

191 Much of the thinking about candidate landing sites for future landed missions has been aimed  
192 at maximizing taphonomic potential by targeting sedimentary environments such as lacustrine  
193 delta deposits. While this Mars exploration strategy is understandable, such an approach  
194 suffers a major epistemological problem: Mars is not Earth. We must recognize that our entire  
195 perspective on how life has evolved and how evidence of life is preserved is colored by the  
196 fact that we live on a planet where photosynthesis evolved. Even if photosynthesis did evolve  
197 on Mars, questions remain as to how successful surface life would have been, and whether  
198 evidence of that life could have been captured in the sedimentary record.

199

200 Considering that some of the most ancient analogue habitats on Earth, hydrothermal and  
201 subsurface environments, are mirrored on Mars, it is logical to search for the signs of  
202 primitive life there in settings analogous to where it may have emerged here. We thereby not  
203 only maximize our chances of finding chemotrophic life, but also of finding the evidence of  
204 prebiotic chemistry that might have led to the formation of life in a sustained habitable  
205 setting.

206

207 The search for biosignatures in hydrothermal deposits must also be questioned. For example,  
208 silica sinter deposits of the type found in Gusev crater<sup>32</sup> are widely considered deposits with  
209 high preservation potential for textural and chemical biomarkers on Earth. But, a significant  
210 fraction of the biomass and biosignatures associated with silica sinters correspond to  
211 photosynthetic bacteria that thrive in fluid mixing zones<sup>65</sup> and therefore the effects of a  
212 possible absence of photosynthesis on biosignature preservation should also be considered in  
213 this context.

214

215 Potential biosignatures in exhumed deep crustal rocks include the following: 1) isotopic  
216 signatures of gasses (e.g. CH<sub>4</sub>) trapped in fluid inclusions, 2) isotopic signatures of minerals,  
217 fluids and organic matter trapped in veins and diagenetic replacements<sup>66</sup>, 3) metal or  
218 carbonate accumulations at redox gradients—especially indicating disequilibrium conditions,  
219 4) biotextures in fractures and pores, 5) microfossils preserved in mineralized veins<sup>66</sup> or  
220 diagenetic cements and concretions, and 6) important organic molecules such as nucleic  
221 acids, lipids, and amino acids in fractures, fluid inclusions, and within mineral aggregates<sup>67,68</sup>.

222 The detection of disequilibrium chemistry implicating life may perhaps be less satisfying than

223 the detection of fossilized microbial mats in lacustrine sediments, but such an approach might  
224 actually teach us more about the origin of life. Because the chemical signatures from the  
225 dawn of life have been entirely obliterated on Earth, finding these clues on Mars, a unique site  
226 within the Solar System, would provide an invaluable window into our own history.

227

228 Given how little we understand about the origin of life on Earth, it makes sense to adopt a  
229 broader plan to seek signs of life. In other words, it is perhaps more logical to seek evidence  
230 of prebiotic chemistry that might have led to the formation of life in sustained habitable  
231 settings rather than searching directly for evolved forms of surface life in ephemeral  
232 environments. We could search for the signs of primitive life on Mars in settings analogous to  
233 where it may have formed on Earth.

234

235 While concerns about the preservation potential of biosignatures in rocks from hydrothermal  
236 and subsurface martian environments are important to consider, it is clear that preservation  
237 potential does not present an ultimate stumbling block. The preservation of biomolecules  
238 associated with hydrothermal activity in the extraterrestrial context has been validated by  
239 their common occurrence in hydrous meteorites with signs of ancient hydrothermal  
240 processing ( $\leq 150^{\circ}\text{C}$ )<sup>69</sup>. Upon the cessation of the hydrothermal event, plunging temperatures  
241 in martian environment would be ideal for preserving biosignatures (e.g. amino acid  
242 enantiomeric ratio)<sup>70</sup>. Silica has been recognized for its significance in microfossil  
243 preservation, and iron-silicate biomineralization in hot spring environments has been shown  
244 to serve as a potent shield to UV radiation<sup>71</sup>. Biomarker preservation in subsurface  
245 environments is a field that has hardly been explored, but biomarkers from Cretaceous  
246 subsurface environments clearly demonstrate that preservation is possible<sup>67</sup>.

247

248 By focusing our search on non-photosynthetic life, we not only maximize our chances of  
249 finding biosignatures on Mars but also uncovering clues to abiogenesis, an aspect that should  
250 be a key part of our exploration strategy. The quest to understand life's origins could be  
251 described as "Follow the energy sources<sup>46</sup>: sulphur, iron and H<sub>2</sub>." That mantra would lead us  
252 to Mars, an iron and sulphur-rich planetary crust with abundant evidence for ancient  
253 hydrothermal activity and H<sub>2</sub> production that could have fuelled an early chemosynthetic  
254 biosphere.

255

256

257

258

259

260 **Methods**

261 The map of hydrothermal and subsurface mineral deposits on Mars (Figure 3) was derived  
262 from multiple sources. The primary data sources include mineral detections by Carter et al.<sup>29</sup>  
263 and Ehlmann et al.<sup>31</sup>, which were created with significant input from the science instrument  
264 teams for the Observatoire pour la minéralogie, l'eau, les glaces et l'activité (OMEGA)<sup>72</sup> and  
265 the Compact Reconnaissance Imaging Spectrometer for Mars (CRISM)<sup>73</sup>. These instruments  
266 have produced 1000s of detections of hydrated minerals on Mars, many of which correspond  
267 to contexts in surface environments and many of which correspond to deposits exhumed from  
268 the subsurface. All of the detections shown in Figure 3 correspond to detections that have  
269 seemingly been exhumed from the subsurface by impact or erosion.

270

271 The edited global-scale datasets of Carter and Ehlmann were supplemented with other  
272 information pertaining to the detection of subsurface, surface or near-surface hydrothermal  
273 deposits. Subsurface carbonate detections were supplemented with data from studies of  
274 exhumed carbonates<sup>74</sup> and a global carbonate<sup>75</sup> study. Serpentine deposits include those  
275 described in a global search for serpentized rocks<sup>76</sup>. Fumarolic silica corresponds to silica  
276 detected by the Spirit rover<sup>77</sup> and with CRISM. Seafloor-type clays correspond to Fe- and  
277 Mg-rich phyllosilicates and carbonates within the Eridania basin on Mars, which was the site  
278 of a large inland sea when the deposits formed >3800 Mya<sup>78</sup>.

279

280

281

282

283

284

285

286

287

288

289

290

291

292

293

294

295

296 **Figure Captions:**

297

298 **Figure 1: A comparison of the age of planetary crust.** Lines represent the best estimate  
299 limits of oldest preserved crust. Dashed lines represent significant uncertainties. The crust of  
300 Mars might provide the best window into the time when abiogenesis occurred on Earth.

301

302 **Figure 2: A comparison of key events in the histories of the Earth and Mars.** The area of  
303 each time line is an approximation of the amount of crust preserved from over different  
304 epochs. The generally unmetamorphosed and well-preserved geologic record of early Mars is  
305 an invaluable window into the geology and prebiotic chemistry of the early Earth.

306

307 **Figure 3: Hydrothermal and exhumed, altered subsurface deposits on Mars.** The global  
308 occurrence of alteration minerals formed in deep crustal or surface hydrothermal  
309 environments detected by infrared remote sensing. See Methods for explanation of data  
310 included in the map.

311

312 **Figure 4: A comparison of the average porosity of thermal gradients of the crusts of**  
313 **Earth and Mars.** For similar rock types and surface porosities, the martian crust contains  
314 significantly more porosity to greater depth than that of the Earth (left). Estimated thermal  
315 gradients for Noachian ( $\phi_N$ ) and modern ( $\phi_m$ ) Mars are lower than that of the modern  
316 continental ( $\phi_c$ ) or oceanic ( $\phi_o$ ) crust of Earth (right). A hypothetical 120°C limit is  
317 encountered at 3-4 km depth on Earth, where the porosity is 1-2%. The same temperature  
318 limit would not be encountered until ~6 km depth on Noachian Mars or much deeper on  
319 modern Mars.

320

321

322

323

324

325

## 326 **References**

327

- 328 1. Herd, C. D. K. *et al.* Origin and evolution of prebiotic organic matter as  
329 inferred from the Tagish Lake meteorite. *Science* **332**, 1304–1307 (2011).
- 330 2. Morbidelli, A., Marchi, S., Bottke, W. F. & Kring, D. A. A sawtooth-like  
331 timeline for the first billion years of lunar bombardment. *Earth Planet. Sci.*  
332 *Lett.* **355–356**, 144–151 (2012).
- 333 3. Sleep, N. H. & Zahnle, K. Refugia from asteroid impacts on early Mars and  
334 the early Earth. *J. Geophys. Res.* **103**, 28529–28544 (1998).
- 335 4. Rosing, M. T. & Frei, R. U-rich Archaean sea-floor sediments from  
336 Greenland - indications of > 3700 Ma oxygenic photosynthesis. *Earth*  
337 *Planet. Sci. Lett.* **217**, 237–244 (2004).
- 338 5. Dodd, M. S. *et al.* Evidence for early life in Earth's oldest hydrothermal vent  
339 precipitates. *Nature* **543**, 60–64 (2017).
- 340 6. Mojzsis, S. J. *et al.* Evidence for life on Earth before 3,800 million years ago.  
341 *Nature* **384**, 55–59 (1996).
- 342 7. Nutman, A. P., Bennett, V. C., Friend, C. R. L., Van Kranendonk, M. J. &

- 343 Chivas, A. R. Rapid emergence of life shown by discovery of 3,700-million-  
344 year-old microbial structures. *Nature* **537**, 535–538 (2016).
- 345 8. Bell, E. A., Boehnke, P., Harrison, T. M. & Mao, W. L. Potentially biogenic  
346 carbon preserved in a 4.1 billion-year-old zircon. *Proc. Natl. Acad. Sci.* **112**,  
347 14518–14521 (2015).
- 348 9. Abramov, O. & Mojzsis, S. J. Thermal effects of impact bombardments on  
349 Noachian Mars. *Earth Planet. Sci. Lett.* **442**, 108–120 (2016).
- 350 10. MILLER, S. L. & UREY, H. C. Organic compound synthesis on the primitive  
351 earth. *Science* **130**, 245–51 (1959).
- 352 11. Szostak, J. W. *et al.* Synthesizing life. *Nature* **409**, 387–390 (2001).
- 353 12. Powner, M. W., Gerland, B. & Sutherland, J. D. Synthesis of activated  
354 pyrimidine ribonucleotides in prebiotically plausible conditions. *Nature*  
355 **459**, 239–242 (2009).
- 356 13. Oró, J. & Kimball, a. P. Synthesis of purines under possible primitive earth  
357 conditions. I. Adenine from hydrogen cyanide. *Arch. Biochem. Biophys.* **94**,  
358 217–227 (1961).
- 359 14. Ricardo, A., Carrigan, M. A., Olcott, A. N. & Benner, S. A. Borate Minerals  
360 Stabilize Ribose. *Science (80-. )*. **303**, 196 LP-196 (2004).
- 361 15. Hanczyc, M. M., Fujikawa, S. M. & Szostak, J. W. Experimental Models of  
362 Primitive Cellular Compartments: Encapsulation, Growth, and Division.  
363 *Science (80-. )*. **302**, 618 LP-622 (2003).
- 364 16. Schmidt, B. E., Blankenship, D. D., Patterson, G. W. & Schenk, P. M. Active  
365 formation of ‘chaos terrain’ over shallow subsurface water on Europa.  
366 *Nature* **479**, 502–5 (2011).
- 367 17. Patthoff, D. A. & Kattenhorn, S. A. A fracture history on Enceladus provides  
368 evidence for a global ocean. *Geophys. Res. Lett.* **38**, (2011).
- 369 18. Vance, S. *et al.* Hydrothermal systems in small ocean planets. *Astrobiology*  
370 **7**, 987–1005 (2007).
- 371 19. Russell, M. J. *et al.* The drive to life on wet and icy worlds. *Astrobiology* **14**,  
372 308–43 (2014).
- 373 20. Běhouňková, M. *et al.* Timing of water plume eruptions on Enceladus  
374 explained by interior viscosity structure. *Nat. Geosci.* **8**, 1–6 (2015).
- 375 21. Sparks, W. B. *et al.* Probing for Evidence of Plumes on Europa with  
376 HST/STIS. *Astrophys. J.* **829**, 121 (2016).
- 377 22. Mulikdjanian, a. Y., Bychkov, a. Y., Dibrova, D. V., Galperin, M. Y. & Koonin,  
378 E. V. PNAS Plus: Origin of first cells at terrestrial, anoxic geothermal fields.  
379 *Proc. Natl. Acad. Sci.* **109**, E821–E830 (2012).
- 380 23. Deamer, D. W. & Georgiou, C. D. Hydrothermal Conditions and the Origin of  
381 Cellular Life. *Astrobiology* **15**, 1091–1095 (2015).
- 382 24. Golombek, M. P. *et al.* Erosion rates at the Mars Exploration Rover landing  
383 sites and long-term climate change on Mars. *J. Geophys. Res.* **111**, 1–14  
384 (2006).
- 385 25. Zuber, M. T. The crust and mantle of Mars. *Nature* **412**, 220–7 (2001).
- 386 26. Lillis, R. J., Robbins, S., Manga, M., Halekas, J. S. & Frey, H. V. Time history of  
387 the Martian dynamo from crater magnetic field analysis. *J. Geophys. Res. E*  
388 *Planets* **118**, 1488–1511 (2013).
- 389 27. Kminek, G. & Bada, J. L. The effect of ionizing radiation on the preservation  
390 of amino acids on Mars. *Earth Planet. Sci. Lett.* **245**, 1–5 (2006).
- 391 28. Lammer, H. *et al.* Outgassing history and escape of the martian atmosphere

- 392 and water inventory. *Space Sci. Rev.* **174**, 113–154 (2013).
- 393 29. Carter, J., Poulet, F., Bibring, J. P., Mangold, N. & Murchie, S. Hydrous  
394 minerals on Mars as seen by the CRISM and OMEGA imaging  
395 spectrometers: Updated global view. *J. Geophys. Res. E Planets* **118**, 831–  
396 858 (2013).
- 397 30. Carter, J., Loizeau, D., Mangold, N., Poulet, F. & Bibring, J. P. Widespread  
398 surface weathering on early Mars: A case for a warmer and wetter climate.  
399 *Icarus* **248**, 373–382 (2014).
- 400 31. Ehlmann, B. L. *et al.* Subsurface water and clay mineral formation during  
401 the early history of Mars. *Nature* **479**, 53–60 (2011).
- 402 32. Ruff, S. W. & Farmer, J. D. ARTICLE Silica deposits on Mars with features  
403 resembling hot spring biosignatures at El Tatio in Chile. *Nat. Commun.* **7**,  
404 1–10 (2016).
- 405 33. Skok, J. R., Mustard, J. F., Ehlmann, B. L., Milliken, R. E. & Murchie, S. L. Silica  
406 deposits in the Nili Patera caldera on the Syrtis Major volcanic complex on  
407 Mars. *Nat. Geosci.* **3**, 838–841 (2010).
- 408 34. Marzo, G. A. *et al.* Evidence for Hesperian impact-induced  
409 hydrothermalism on Mars. *Icarus* **208**, 667–683 (2010).
- 410 35. Turner, S. M. R., Bridges, J. C., Grebbly, S. & Ehlmann, B. L. Hydrothermal  
411 activity recorded in post Noachian-aged impact craters on Mars. *J. Geophys.*  
412 *Res. Planets* **121**, 608–625 (2016).
- 413 36. Cockell, C. S. *et al.* Habitability: A Review. *Astrobiology* **16**, 89–117 (2016).
- 414 37. Whitman, W. B., Coleman, D. C. & Wiebe, W. J. Prokaryotes: The unseen  
415 majority. *Proc. Natl. Acad. Sci. U. S. A.* **95**, 6578–6583 (1998).
- 416 38. Onstott, T. C., Colwell, F. S., Kieft, T. L., Murdoch, L. & Phelps, T. J. New  
417 Horizons for Deep Subsurface Microbiology. *Microbe* **4**, 499–505 (2009).
- 418 39. Stevens, T. O. Subsurface lithoautotrophic microbial ecosystems (SLMEs)  
419 in igneous rocks: Prospects for detection. *Instruments, Methods, Mission.*  
420 *Investig. Extraterr. Microorg.* **3111**, 358–365530 (1997).
- 421 40. Pedersen, K. Microbial life in deep granitic rock. in *FEMS Microbiology*  
422 *Reviews* **20**, 399–414 (1997).
- 423 41. Stevens, T. O. & Mckinley, J. P. Lithoautotrophic microbial ecosystems in  
424 deep basalt aquifers. *Science (80-. ).* **270**, 450–454 (1995).
- 425 42. Coveney, R. M., Goebel, E. D., Zeller, E. J., Dreschhoff, G. A. M. & Angino, E. E.  
426 Serpentinization and the origin of hydrogen gas in Kansas ( USA).  
427 *American Association of Petroleum Geologists Bulletin* **71**, 39–48 (1987).
- 428 43. Lillis, R. J., Manga, M., Mitchell, D. L., Lin, R. P. & Acuna, M. H. Unusual  
429 magnetic signature of the Hadriaca Patera volcano: Implications for early  
430 Mars. *Geophys. Res. Lett.* **33**, (2006).
- 431 44. Michalski, J. R. *et al.* Groundwater activity on Mars and implications for a  
432 deep biosphere. *Nat. Geosci.* **6**, 133–138 (2013).
- 433 45. Kelley, D. S. *et al.* A serpentinite-hosted ecosystem: The Lost City  
434 hydrothermal field. *Science (80-. ).* **307**, 1428–1434 (2005).
- 435 46. Parnell, J., Boyce, A. J. & Blamey, N. J. F. Follow the methane: the search for  
436 a deep biosphere, and the case for sampling serpentinites, on Mars. *Int. J.*  
437 *Astrobiol.* **9**, 193–200 (2010).
- 438 47. Onstott, T. C. *et al.* Martian CH<sub>4</sub>: Sources, flux, and detection. *Astrobiology*  
439 **6**, 377–395 (2006).
- 440 48. Niles, P. B. *et al.* Geochemistry of Carbonates on Mars: Implications for

- 441 Climate History and Nature of Aqueous Environments. *Space Sci. Rev.* **174**,  
442 301–328 (2013).
- 443 49. Russell, M. J., Hall, A. J. & Martin, W. Serpentinization as a source of energy  
444 at the origin of life. *Geobiology* **8**, 355–371 (2010).
- 445 50. Dismukes, G. C. *et al.* The origin of atmospheric oxygen on Earth: the  
446 innovation of oxygenic photosynthesis. *Proc. Natl. Acad. Sci. U. S. A.* **98**,  
447 2170–5 (2001).
- 448 51. Soo, R. M., Hemp, J., Parks, D. H., Fischer, W. W. & Hugenholtz, P. On the  
449 origins of oxygenic photosynthesis and aerobic respiration in  
450 Cyanobacteria. *Science (80-. )*. **355**, 1436–1440 (2017).
- 451 52. Shih, P. M., Hemp, J., Ward, L. M., Matzke, N. J. & Fischer, W. W. Crown  
452 group Oxyphotobacteria postdate the rise of oxygen. *Geobiology* **15**, 19–29  
453 (2017).
- 454 53. Johnson, J. E. *et al.* Manganese-oxidizing photosynthesis before the rise of  
455 cyanobacteria. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 11238–11243 (2013).
- 456 54. Farquhar, J., Bao, H. & Thiemens, M. Atmospheric Influence of Earth's  
457 Earliest Sulfur Cycle. *Science (80-. )*. **289**, 756–758 (2000).
- 458 55. Lyons, T. W., Reinhard, C. T. & Planavsky, N. J. The rise of oxygen in Earth's  
459 early ocean and atmosphere. *Nature* **506**, 307–15 (2014).
- 460 56. Davila, A. F. & Schulze-Makuch, D. The Last Possible Outposts for Life on  
461 Mars. *Astrobiology* **16**, ast.2015.1380 (2016).
- 462 57. Veizer, J. & Mackenzie, F. T. in *Treatise on Geochemistry: Second Edition* **9**,  
463 399–435 (2013).
- 464 58. Wordsworth, R. D., Kerber, L., Pierrehumbert, R. T., Forget, F. & Head, J. W.  
465 Comparison of 'warm and wet' and 'cold and icy' scenarios for early Mars  
466 in a 3-D climate model. *J. Geophys. Res. Planets* **120**, 1201–1219 (2015).
- 467 59. Wordsworth, R. *et al.* Global modelling of the early martian climate under a  
468 denser CO<sub>2</sub> atmosphere: Water cycle and ice evolution. *Icarus* **222**, 1–19  
469 (2013).
- 470 60. De Villiers, G., Kleinhans, M. G. & Postma, G. Experimental delta formation  
471 in crater lakes and implications for interpretation of Martian deltas. *J.*  
472 *Geophys. Res. E Planets* **118**, 651–670 (2013).
- 473 61. Fassett, C. I. & III, J. W. H. Fluvial sedimentary deposits on Mars: Ancient  
474 deltas in a crater lake in the Nili Fossae region. *Geophys. Res. Lett.* **32**, 1–5  
475 (2005).
- 476 62. Fassett, C. I. & Head, J. W. Sequence and timing of conditions on early Mars.  
477 *Icarus* **211**, 1204–1214 (2011).
- 478 63. Vanwonterghem, I. *et al.* Methylophilic methanogenesis discovered in the  
479 archaeal phylum Verstraetearchaeota. *Nat. Microbiol.* **1**, 16170 (2016).
- 480 64. Dadachova, E. & Casadevall, A. Ionizing radiation: how fungi cope, adapt,  
481 and exploit with the help of melanin. *Current Opinion in Microbiology* **11**,  
482 525–531 (2008).
- 483 65. Kato, K. *et al.* Microbial Mat Boundaries between Chemolithotrophs and  
484 Phototrophs in Geothermal Hot Spring Effluents. *Geomicrobiol. J.* **21**, 91–98  
485 (2004).
- 486 66. Pedersen, K. *et al.* Evidence of ancient life at 207 m depth in a granitic  
487 aquifer. *Geology* **25**, 827–830 (1997).
- 488 67. Klein, F. *et al.* Fluid mixing and the deep biosphere of a fossil Lost City-type  
489 hydrothermal system at the Iberia Margin. *Proc. Natl. Acad. Sci. U. S. A.* **112**,

- 490 12036–41 (2015).
- 491 68. Ringelberg, D. B., Sutton, S. & White, D. C. Biomass, bioactivity and  
492 biodiversity: Microbial ecology of the deep subsurface: Analysis of ester-  
493 linked phospholipid fatty acids. in *FEMS Microbiology Reviews* **20**, 371–377  
494 (1997).
- 495 69. Ehrenfreund, P., Glavin, D. P., Botta, O., Cooper, G. & Bada, J. L.  
496 Extraterrestrial amino acids in Orgueil and Ivuna: tracing the parent body  
497 of CI type carbonaceous chondrites. *Proc. Natl. Acad. Sci. U. S. A.* **98**, 2138–  
498 2141 (2001).
- 499 70. Bada, J. L. & McDonald, G. D. Amino Acid Racemization on Mars:  
500 Implications for the Preservation of Biomolecules from an Extinct Martian  
501 Biota. *Icarus* **114**, 139–143 (1995).
- 502 71. Phoenix, V. R., Konhauser, K. O., Adams, D. G. & Bottrell, S. H. Role of  
503 biomineralization as an ultraviolet shield: Implications for Archean life.  
504 *Geology* **29**, 823–826 (2002).
- 505 72. Poulet, F. *et al.* Martian surface mineralogy from Observatoire pour la  
506 Mineralogie, l’Eau, les Glaces et l’Activite on board the Mars Express  
507 spacecraft (OMEGA/MEx): Global mineral maps. *J. Geophys. Res.* **112**,  
508 (2007).
- 509 73. Murchie, S. *et al.* Compact reconnaissance Imaging Spectrometer for Mars  
510 (CRISM) on Mars Reconnaissance Orbiter (MRO). *J. Geophys. Res.* **112**,  
511 (2007).
- 512 74. Michalski, J. R. & Niles, P. B. Deep crustal carbonate rocks exposed by  
513 meteor impact on Mars. *Nat. Geosci.* **3**, 751–755 (2010).
- 514 75. Wray, J. J. *et al.* Orbital evidence for more widespread carbonate-bearing  
515 rocks on Mars. *J. Geophys. Res. E Planets* **121**, 652–677 (2016).
- 516 76. Ehlmann, B. L., Mustard, J. F. & Murchie, S. L. Geologic setting of serpentine  
517 deposits on Mars. *Geophys. Res. Lett.* **37**, (2010).
- 518 77. Squyres, S. W. *et al.* Detection of Silica-Rich Deposits on Mars. *Science (80-*  
519 *)*. **320**, 1063–1067 (2008).
- 520 78. Michalski, J. R., Dobrea, E. Z. N., Niles, P. B. & Cuadros, J. Ancient  
521 hydrothermal seafloor deposits in Eridania basin on Mars. **8**, 15978  
522 (2017).
- 523
- 524







