

Simon D, Bradley C, Gönder-Frederick L, Eschwège EM, de Pablos-Velasco P, Vandenberghe H, Bouzamondo H and Parhofer KG (2010) The PANORAMA pan-European survey: Hypoglycaemia associated with different pharmacological treatments for type 2 diabetes. *Value in Health* 13 (7) A297 (PDB73).

Poster presented at the ISPOR 13th Annual European Congress: November 6-9, 2010 Prague, Czech Republic.

THE PANORAMA PAN-EUROPEAN SURVEY: HYPOGLYCAEMIA ASSOCIATED WITH DIFFERENT PHARMACOLOGICAL TREATMENTS FOR TYPE 2 DIABETES.

Simon D¹, Bradley C², Gönder-Frederick L³, Eschwège EM⁴, de Pablos-Velasco P⁵, Vandenberghe H⁶, Bouzamondo H⁷, Parhofer KG⁸

¹Hôpital de la Pitié/INSERM U-1018, Paris, France, ²Royal Holloway, University of London, Surrey, United Kingdom, ³University of Virginia Health System, Charlottesville, VA, USA, ⁴INSERM, Villejuif, France, ⁵Dr Negrin Hospital, Las Palmas University, Las Palmas de Gran Canaria, Spain, ⁶AstraZeneca, Zaventem, Belgium, ⁷Bristol-Myers Squibb, Paris, France, ⁸Klinikum der Universität München, Munich, Germany

OBJECTIVES: Hypoglycaemia can be a side effect of glucose-lowering treatment in patients with type 2 diabetes (T2D) that may counterbalance the beneficial effects of diabetes control. PANORAMA is a large (n=5156) pan-European cross-sectional survey (NCT00916513) of patients assessing patient reported outcomes and glycaemic control. This subgroup analysis compared rates of severe and non-severe hypoglycaemic events in patients taking different pharmacological treatment regimens.

METHODS: Patients with T2D were randomly or consecutively selected from medical practices in eight countries. Patients were aged ≥ 40 years, with T2D diagnosed >1 year and a clinic medical record available >1 year. All patients received dietary/exercise advice and most were also taking either oral antidiabetic drugs (OADs) and/or injectables (insulin and/or GLP-1 receptor agonists). Patients included in this subgroup analysis had been taking the same pharmacological treatment regimen for ≥ 12 months. Patient-reported frequency of severe (symptomatic episodes requiring external assistance) and non-severe hypoglycaemic episodes in the past year were examined. **RESULTS:** In this subgroup analysis 3106 patients were evaluated including: 1346 taking only OADs without secretagogues; 1452 taking only OADs including secretagogues (sulphonylurea/glinides) and 308 on insulin alone. The percentages of patients experiencing >1 non-severe hypoglycaemic episode in each treatment group were: 8.9% for patients taking OADs without secretagogues; 17.5% for patients taking OADs including secretagogues and 47.4% for patients using insulin alone. The differences between these three treatment categories (pair-wise comparisons) were highly significant ($p < 0.001$). The percentage of patients reporting ≥ 1 severe hypoglycaemic episode was greater for OADs including secretagogues versus no secretagogues (3.0% versus 1.3%; $p = 0.011$) and for insulin alone versus OADs including secretagogues (13.7% versus 3.0%; $p < 0.001$). **CONCLUSIONS:** Among patients with T2D on glucose-lowering medication, rates of non-severe and severe hypoglycaemic episodes were lowest amongst patients treated with OADs not including secretagogues and highest among patients treated with insulin alone.