

Three months' withdrawal of growth hormone (GH) therapy from GH-deficient adults does not affect cognitive functioning

CV McMillan¹, C Bradley¹, J Gibney², DL Russell-Jones³ and PH Sonksen²

¹Health Psychology Research, Dept of Psychology, Royal Holloway, University of London, Egham, Surrey TW20 0EX, UK, ²St Thomas' Hospital, London, ³Royal Surrey County Hospital, Guildford

Background

- Adults with untreated GH deficiency (GHD) may complain of poor cognitive functioning: difficulties in short and long-term memory, word-finding, concentration and decreased attention span¹.
- Some studies^{2,3} have found improvement in cognitive functioning with GH treatment but others⁴ no improvement.
- No known previous study has investigated effects on cognitive functioning of discontinuing GH treatment from GH-treated adults.

Objective

To determine the effect on cognitive functioning of discontinuation of GH treatment from adults with GHD.

Method

- Design: randomised double-blind placebo-controlled trial.
- GH-replacement therapy discontinued for 3 months from 12 of 21 adults, with 9 patients continuing on GH.
- Cognitive tests completed at baseline and end-point, except Test 8 (see Measures) completed once only.
- Semi-structured interviews plus questionnaires.

Measures

Criteria for selection of cognitive tests

- Cover main areas of cognitive functioning.
- Short.
- 2 equivalent forms available, or sufficient numbers of stimuli to split into two equivalent sets (Tests 1 and 4 below).


Cognitive tests

- 1 Warrington Recognition Memory Test⁵. (Visual recognition memory for faces and words).
- 2 Rey Auditory Verbal Learning Test⁶ (AVLT). (Short-term memory, learning, delayed recall and executive function).
- 3 Trail making⁷. (Executive function and attention).
- 4 Graded Naming Test⁸. (Word-naming).
- 5 Benton Visual Retention Test⁹. (Visual recall).
- 6 Controlled Oral Word Association Test¹⁰ (COWAT). (Executive function).
- 7 Digit span test, a sub-test of Wechsler Adult Intelligence Scale¹¹. (Short-term memory).
- 8 National Adult Reading Test¹² (NART). (Pre-morbid intelligence).

Questionnaires included:

General Health Questionnaire¹³ (GHQ), a measure of change in psychological well-being. Range 0-90.

Fig. 1: Some cognitive test scores at baseline (T1) and end-point (T2)



Results

Cognitive tests

- No significant between-group differences in pre-morbid intelligence
- No significant change (nor any trends in data) in either treatment group between baseline and end-point. (See Fig. 1 for a selection of test results: higher scores indicate better cognitive function).
- No significant between-group differences in scores at end-point when baseline scores had been partialled out.

Interviews

Three placebo- and 2 GH-treated patients reported perceived worsening of memory and concentration over the withdrawal period. Four of these 5 patients also reported increased depression (confirmed by scores on the GHQ, mean change of 23.5 +/- 14.2).

Conclusions

Some studies have reported negative effects of GHD on cognitive functioning, with improvements after GH-replacement therapy. The present study found no detrimental effects on cognitive functioning after discontinuation of GH treatment in GH-deficient adults. Patient reports of perceived poorer cognitive functioning in relation to untreated GHD may be associated with increased depression.

References

- 1 Han SM et al. (1993) Preliminary report on the development of a disease specific instrument for assessing quality of life of adults with growth hormone deficiency. *Acta Endocrinologica* 128, (Suppl. 2): 37-40.
- 2 Ruzman P et al. (1995) Quality of life in adults with growth hormone (GH) deficiency: Response to treatment with recombinant human GH in a 3-year study. *Journal of Clinical Endocrinology and Metabolism* 81, 1985-90.
- 3 Toegen JB et al. (1993) Cognitive changes during growth hormone replacement in adult men. *Psychoneuroendocrinology* 23(1), 45-55.
- 4 Baum HB et al. (1978) Effects of physiological growth hormone therapy on cognition and quality of life in patients with untreated GH deficiency. *Journal of Clinical Endocrinology and Metabolism* 48 (2), 319-49.

- 5 Warrington E. (1984) *The Recognition Memory Test*. Windsor: NFER-Nelson.
- 6 Rey A. (1964) *Levasser subteste on psychologie*. Paris: Presses Universitaires de France.
- 7 Trail Making Test. Tucson, Arizona: Reitan Neuropsychology Laboratory.
- 8 Nicklison P and Warrington E. (1963) *Graded Naming Test*. Windsor: NFER-Nelson.
- 9 Benton AL. (1975) *Revised Visual Retention Test* (4th ed.). San Antonio: Psychological Corp.
- 10 Benton AL and Hamsher K de S. (1989) *Multilingual Aphasia Examination*. Iowa City, IJA Associates.
- 11 Wechsler D. (1985) *The Wechsler Adult Intelligence Scale*. New York: Psychological Corporation.

¹³Nelson HE. (1971) *National Adult Reading Test Manual*. Windsor: NFER-Nelson.
¹⁴Gulberg DP. (1972) *The detection of psychiatric illness by questionnaire*. Maudsley Monograph no. 21. Oxford: Oxford University Press.

Acknowledgements

Cecyln McMillan was funded by a research studentship from Lilly Industries Ltd. We also acknowledge with thanks, help received from colleagues at St Thomas' Hospital, London and from the patients who contributed to the studies.